Preprint typeset in JHEP style - HYPER VERSION

arXiv:1103.2598v1 [hep-th] 14 Mar 2011

Wall-Crossing in Coupled 2d-4d Systems

Davide Gaiotto[1], **Gregory W. Moore**[2], **Andrew Neitzke**[3]

[1] *School of Natural Sciences, Institute for Advanced Study,*
Princeton, NJ 08540, USA
[2] *NHETC and Department of Physics and Astronomy, Rutgers University,*
Piscataway, NJ 08855–0849, USA
[3] *Department of Mathematics, University of Texas at Austin,*
Austin, TX 78712, USA

`dgaiotto@ias.edu`, `gmoore@physics.rutgers.edu`, `neitzke@math.utexas.edu`

Abstract: We introduce a new wall-crossing formula which combines and generalizes the Cecotti-Vafa and Kontsevich-Soibelman formulas for supersymmetric 2d and 4d systems respectively. This 2d-4d wall-crossing formula governs the wall-crossing of BPS states in an $\mathcal{N} = 2$ supersymmetric 4d gauge theory coupled to a supersymmetric surface defect. When the theory and defect are compactified on a circle, we get a 3d theory with a supersymmetric line operator, corresponding to a hyperholomorphic connection on a vector bundle over a hyperkähler space. The 2d-4d wall-crossing formula can be interpreted as a smoothness condition for this hyperholomorphic connection. We explain how the 2d-4d BPS spectrum can be determined for 4d theories of class $\mathcal{S}$, that is, for those theories obtained by compactifying the six-dimensional $(0, 2)$ theory with a partial topological twist on a punctured Riemann surface C. For such theories there are canonical surface defects. We illustrate with several examples in the case of A_1 theories of class $\mathcal{S}$. Finally, we indicate how our results can be used to produce solutions to the A_1 Hitchin equations on the Riemann surface C.

Contents

1. Introduction: A guided tour of the paper **3**

2. Formal statement of wall-crossing formulae **11**
- 2.1 The 2d Cecotti-Vafa wall-crossing formula 11
- 2.2 The 4d Kontsevich-Soibelman wall-crossing formula 13
- 2.3 The 2d-4d wall-crossing formula 14
  - 2.3.1 Automorphisms of wall-crossing identities 18
- 2.4 Examples 19

3. A physical interpretation **23**
- 3.1 Surface defects: kinematics 23
- 3.2 Examples of surface defects 24
- 3.3 Low energy (IR) description of the 2d-4d system 25
  - 3.3.1 Massive 2d theories in the IR 25
  - 3.3.2 4d gauge theories in the IR 26
  - 3.3.3 2d-4d theories in the IR: Local picture, Gukov-Witten parameters, and the torsors Γ_{ij} 28
  - 3.3.4 Superpotentials and their shifts: the torsors Γ_i 30
  - 3.3.5 2d-4d theories in the IR: Global picture 31
  - 3.3.6 Algebraic integrable systems, mirror symmetry, and surface defects 33
- 3.4 Surface defect (twisted) chiral rings: periods as superpotentials 34
  - 3.4.1 A potential anomaly 36
  - 3.4.2 Example: The GW surface defect in pure $SU(2)$ gauge theory 37
  - 3.4.3 An example of the anomaly 44
- 3.5 BPS degeneracies and wall-crossing 46
- 3.6 Cancellation of surface defect global anomalies by gauging flavor symmetries 48
  - 3.6.1 Flavor holonomies 50
  - 3.6.2 Anomaly cancellation 51

4. Line defects, interfaces, framed BPS states, and a spin 2d-4d wall-crossing formula **53**
- 4.1 Interfaces 53
- 4.2 Review of the 4d case: Framed BPS states and halos 54
- 4.3 Analog in two dimensions 56
- 4.4 Framed indices in the 2d-4d case 57
- 4.5 Ring of line defects 58
- 4.6 A 2d-4d spin wall-crossing formula 60
- 4.7 Halo picture 61
  - 4.7.1 Crossing soliton walls 61

4.7.2 Crossing 4d walls 61
4.7.3 Determining the generalized quantum dilogarithms $\Phi_{\mathbb{S},\gamma}$ 63

5. Compactification to 3d and hyperholomorphic bundles 64
5.1 The dimensionally reduced action 64
5.2 Globalizing 65
5.2.1 Line bundles 65
5.2.2 Vector bundles 66
5.2.3 Twisted vector bundles and mirror symmetry 66
5.3 Constraints from supersymmetry: hyperkähler and hyperholomorphic 68
5.4 The semiflat connection is hyperholomorphic in the semiflat geometry 68
5.5 Physical interpretation of V: Expectation values 70
5.6 Integral equations 71
5.7 Wall-crossing formula as a consistency condition 75
5.8 $\mathcal{Y}_{\gamma_i}$ and framed BPS states 76

6. Local models and resolution of singularities 76
6.1 Massless 4d and 2d particles on a one-dimensional Coulomb branch 77
6.1.1 Local system and BPS degeneracies 77
6.1.2 Review of the PTN geometry 79
6.1.3 The semiflat line bundle 83
6.1.4 The quantum-corrected bundle V 84
6.1.5 Deriving the connection 85
6.1.6 Analyzing the singularities 86
6.1.7 Mirror manifolds and mirror branes 87
6.2 A massless 2d particle with 4d gauge charges only 87
6.3 A massless 2d soliton 89

7. Example: A type theories and Hitchin systems 92
7.1 Review of pure 4d data 93
7.2 2d-4d data 94
7.3 3d compactification and Hitchin's equations 98
7.4 Lagrangian descriptions of A_1 theories and their surface defects 100
7.5 A brief review of the WKB analysis and its extension to 2d-4d 102
7.5.1 4d review 102
7.5.2 Extension to 2d-4d 105

8. Examples of 2d-4d wall-crossing in some A_1 theories 110
8.1 Simple Argyres-Douglas-type theories 111
8.1.1 $N = 1$ 112
8.1.2 $N = 2$ 115
8.1.3 Larger N 121
8.2 The $\mathbb{CP}^1$ sigma model 124
8.3 The canonical surface defect in pure $SU(2)$ gauge theory 128

8.3.1 Preliminaries on local systems 129
8.3.2 The soliton spectrum at strong coupling 130
8.3.3 The soliton spectrum for weak 4d coupling and strong 2d coupling 134
8.3.4 The soliton spectrum for weak 4d coupling and weak 2d coupling 139

9. An application: solving Hitchin systems by integral equations 143

10. Future directions 144

A. Relation of 2d and 4d superalgebras and their multiplets 147

B. A simple Hitchin system 149

C. Two-dimensional bound state radius in Landau-Ginzburg theories 150

D. Physical models for the affine linearity of $\omega(\gamma, \gamma_i)$ 152
D.1 A supersymmetric quantum mechanics model 152
D.2 Probe particle in the presence of a solenoid and dyon 153

E. Twistor construction of hyperholomorphic connections 155

F. $\mathcal{Y}_{\gamma_{ij'}}$ in A_1 theories 157
F.1 Defining $\mathcal{Y}$ 158
F.2 Multiplication laws 158
F.3 Morphism: crossing a separating WKB curve 159
F.4 Morphism: crossing an edge 161
F.5 Morphism: flips and juggles 161

G. A review: hidden flavor symmetries in 3d Coulomb branches 161

---

1. Introduction: A guided tour of the paper

The present paper is a continuation of a project revolving around the intertwined themes of wall-crossing, hyperkähler geometry, BPS states, and their application to the rich mathematical physics associated with four-dimensional field theories with $\mathcal{N} = 2$ supersymmetry [1, 2, 3]. In this paper we introduce several new ingredients into the emerging structure. All of these new ingredients are associated in one way or another with *surface defects.*

Several developments over the past decade have amply demonstrated that the inclusion of defects of decreasing codimension leads to physical systems which possess increasing richness and beauty, albeit at the price of increasing complexity. For examples, in 2-dimensional conformal field theory the inclusion of defect lines enhances the structure wonderfully. In four dimensional $\mathcal{N} = 4$ supersymmetric field theory the line defects shed much light on S-duality [4] and play a central role in the geometric Langlands program [5].

Deeper developments along this direction have come with the inclusion of surface defects [6, 7, 8]. Finally, in topological field theory, a whole hierarchy of defects leads to a beautiful mathematical structure involving higher category theory [9, 10, 11, 12, 13, 14].

The present paper together with [15, 16, 17] may be viewed as part of an analogous line of development for theories with $\mathcal{N} = 2$ supersymmetry.[1] For example, in [3] we considered defects of codimension three — that is, *line defects* in $\mathcal{N} = 2$ theories. This led naturally to a new class of BPS states, the framed BPS states. The consistency of framed BPS indices gave a very simple and natural derivation of the "motivic" Kontsevich-Soibelman wall-crossing formula for the "vanilla" BPS states. (As in [3], we refer to the BPS states of the original four-dimensional theory without defects as "vanilla.") In addition it led to a natural quantization of the algebra of holomorphic functions on Seiberg-Witten moduli spaces as well as to new connections with the mathematics of laminations of punctured Riemann surfaces, quantized Teichmüller theory, cluster algebras, and cluster varieties. In the present paper, as we have said, we move on to investigate some aspects of codimension two defects, also known as surface defects. Not surprisingly, this introduces a new layer of complexity. The remainder of this introduction is an expository account of the paper.

The surface defects we will consider preserve four of the eight supersymmetries of the vanilla 4-dimensional theory (§3.1 and Appendix A). Various UV constructions of such defects are reviewed in §3.2. One representative construction proceeds by coupling a 1+1 dimensional theory $\mathcal{T}_{2d}$ with (2,2) supersymmetry on a surface $\mathbb{S}$ to an ambient four-dimensional theory $\mathcal{T}_{4d}$. We assume that $\mathcal{T}_{2d}$ has a compact global symmetry group G and moreover we assume that in isolation $\mathcal{T}_{2d}$ has a finite number of massive vacua, labeled $i \in \mathcal{V}[\mathbb{S}]$.[2] The G-symmetry can then be gauged by coupling to a G-gauge theory in $\mathcal{T}_{4d}$. A second UV construction, due to Gukov and Witten, involves just a 4d theory with G gauge symmetry but with reduced structure group along the surface $\mathbb{S}$ [6]. In addition we can consider domain walls welding together different theories $\mathcal{T}_{2d}$ and $\mathcal{T}'_{2d}$, or more generally different defects on $\mathbb{S}$ and $\mathbb{S}'$. The domain wall can be viewed as a line defect in four dimensions embedded in a surface. We will refer to it as a supersymmetric interface. We refer to the general systems we have just described as *coupled 2d-4d systems* or 2d-4d systems for short.

2d-4d systems can support novel and nontrivial dynamics, qualitatively different from the dynamics exhibited by, say, $\mathcal{T}_{2d}$ and $\mathcal{T}_{4d}$ in isolation. They have a rich set of BPS states. Besides the 4d vanilla BPS states, there are BPS states of the type associated to the 2d theory $\mathcal{T}_{2d}$. Such BPS states were studied extensively by Cecotti and Vafa and collaborators in the two-dimensional context [18, 19, 20]. The coupling to a 4d theory $\mathcal{T}_{4d}$ has an effect akin to the introduction of 2d twisted masses for the G-symmetry [21, 22, 23]. ("Twisted masses" refers to the parameters introduced in 2d $(2,2)$ models in [24].) For example, there can be infinitely many solitons interpolating between distinct vacua of the surface defect and carrying four-dimensional gauge and "flavor charges."[3] There can also be BPS states localized on $\mathbb{S}$ which do not interpolate between different vacua but rather reside in a single vacuum. These states too can carry four-dimensional gauge and flavor charge. Finally, in the presence of interfaces there are again framed BPS states.

---

[1] This of course suggests that we should go on to investigate domain walls in $\mathcal{N} = 2$ theories, but we will not address that interesting topic in the present work.

[2] The symbol $\mathbb{S}$ denotes the location of the surface defect, but also sometimes the "theory" on the defect. It has no meaning without specifying the ambient four-dimensional theory.

In order to exhibit the curious antics of these new BPS states we must turn to a low energy effective field theory description of the dynamics. There is a description of the low energy dynamics of 2d-4d systems generalizing the renowned solution of pure four-dimensional theories initiated by Seiberg and Witten [25, 26] (For reviews see, for example, [27, 28].) This is described in some detail in §3, following and developing further the story in [17]. The four-dimensional theory has a Coulomb branch of vacua $\mathcal{B}$.[4] In 2d-4d systems this is generalized to a ramified cover $\mathcal{B}_{\mathbb{S}}$ whose sheets are in correspondence with vacua $\mathcal{V}[\mathbb{S}]$. The dynamics at a vacuum $(u, i) \in \mathcal{B}_{\mathbb{S}}$ is described by an effective four-dimensional abelian $\mathcal{N} = 2$ gauge theory coupled to a twisted chiral effective 2d $(2, 2)$ superpotential $\mathcal{W}$, localized on $\mathbb{S}$. (See equation (3.12).) Thus, the description of the low-energy dynamics of four-dimensional theories by a prepotential $\mathcal{F}$ is replaced in 2d-4d systems by a description in terms of a pair $(\mathcal{F}, \mathcal{W})$. There is a simple IR picture of the ground states of the surface defect in a definite charge sector: the defect looks like a solenoid, outside of which there is a flat abelian gauge field. If we describe the 4d IR abelian gauge theory by a self-dual gauge field $\mathbb{F} = \mathrm{d}\mathbb{A}$, then the holonomy of $\mathbb{A}$ around the solenoid is specified by a symplectic vector ν (equation (3.19)). Upon choosing a duality frame, ν has electric and magnetic components η and α, related to $\mathcal{W}$ by equation (3.15):

$$t = \eta + \tau\alpha = \frac{\partial \mathcal{W}}{\partial a}. \tag{1.1}$$

We refer to (η, α) as (infrared) Gukov-Witten parameters.

When two defects are welded together by an interface the analogous physical picture of the ground state in a definite charge sector is that of a pair of half-solenoids glued together on a BPS dyon, rather like a nunchuk, or a boa which has swallowed an elephant [29]. See Figure 12. The dyon allows flux to escape from the solenoids, and hence the field outside is no longer flat. Moreover, the dyon charge need not satisfy Dirac quantization. Indeed we can regard the worldsheets of the solenoids as those of Dirac strings which have materialized into observable objects.

If we wish to describe our ground states with more precision we must be careful about the description of the 4d charges of our BPS states. It turns out that this is no simple matter. We begin with the local system over $\mathcal{B}$ of electromagnetic and flavor charges Γ, and its quotient system Γ_g of electromagnetic charges. The latter is a local system of symplectic lattices. (See equation (3.6).) Associated to this is the Seiberg-Witten moduli space

---

[3] In this paper flavor charges play an important role. We define a "flavor symmetry" of a 4d $\mathcal{N} = 2$ theory to be a global symmetry commuting with the supersymmetries which is not spontaneously broken on the Coulomb branch of vacua. States and operators of the theory will transform in representations of the flavor symmetry group and "flavor charges" are characters of this group. We will in general limit our considerations to flavor symmetry groups which are abelian.

[4] We do not venture onto Higgs branches in this paper.

$$\mathcal{M} := \Gamma_g^* \otimes \mathbb{R}/(2\pi\mathbb{Z}), \tag{1.2}$$

one of the central actors in [1, 2, 3]. Geometrically, the holonomy $\tilde{\nu}$ of $\mathbb{A}$ should be viewed as a "section" of the "mirror-dual" moduli space $\widetilde{\mathcal{M}} = \Gamma_g \otimes \mathbb{R}/(2\pi\mathbb{Z})$, but it will be important to consider a lift $\nu \in \Gamma_g \otimes \mathbb{R}$, thus assigning a definite flux to the interior of the solenoid. Of course, since Γ_g is a nontrivial local system with monodromy there will be no globally well-defined section ν, only a multisection. Generically the possible values of ν above (u, i) will be a "torsor" for Γ_u [5] and monodromies will act in an affine-linear way on ν. It follows then from (1.1) that the superpotential must have monodromy, and that the set of possible effective superpotentials is a torsor for Γ. (This assumes that the "mass parameters" dual to the flavor charges are generic.) Physically this monodromy can be traced back to the fact that at singular loci $\mathcal{B}^{\text{sing}}$ not only will four-dimensional BPS states become massless, but also chiral multiplets on $\mathbb{S}$ will become massless. In §3.5, Witten's famous computation of effective twisted superpotentials in gauged linear sigma models [30, 31] is reinterpreted as computing an extension of the central charge function $Z : \Gamma \to \mathbb{C}$ to the torsors associated to $\mathbb{S}$. A surprising physical implication is that the number of chiral multiplets on the surface defect is not an absolute invariant but rather more like a gauge choice in the description of the 2d-4d system. This is intimately related to the fact that a 2d chiral multiplet can mix with 4d hypermultiplets with the same gauge and flavor charges. We will denote the torsor associated to a vacuum i as Γ_i. The Γ_i form a local system of torsors over $\mathcal{B}_{\mathbb{S}}$.[6]

If we consider 2d solitons between vacua i and j, charge quantization is controlled by the difference in the holonomies $\tilde{\nu}_i$ and $\tilde{\nu}_j$ associated to the two vacua, and hence the 4d charges of 2d solitons form a Γ-torsor Γ_{ij}. With interfaces these are further generalized to $\Gamma_{ij'}$, where i, j' are vacua of the two surface defects welded by the interface. Charges in Γ_{ij} will be denoted by symbols like γ_{ij}, and satisfy addition rules like $\gamma_{ij} + \gamma_{jk} = \gamma_{ik}$. Mathematically this means we define a *groupoid of vacua* $\mathbb{V}$ which will prove to be a useful concept when we discuss wall-crossing. See §2.3 for the formal construction.

The charge torsors described above grade the Hilbert spaces of 2d solitons as well as other BPS states, framed and unframed. There are BPS bounds in all these charge sectors and hence associated to our new BPS states are a host of new BPS indices. These include:

- $\mu(\gamma_{ij})$: These are the degeneracies of 2d solitons with 4d gauge charge γ_{ij}. They generalize the degeneracies μ_{ij} studied by Cecotti and Vafa in the pure 2d context [20]. See §3.5 and Appendix A.

---

[5] A *torsor* for a group G is a principal homogeneous space $\mathbf{S}$ for G. This means that there is a transitive G-action on $\mathbf{S}$ and moreover there are no nontrivial stabilizers. That is, for any two elements $s_1, s_2 \in \mathbf{S}$ there is a $g \in G$ so that $s_2 = g \cdot s_1$, and moreover, if $s = g \cdot s$ for any s then $g = 1$. Thus, $\mathbf{S}$ is "a copy of G," but there is no distinguished element of $\mathbf{S}$ corresponding to the identity element of G. A typical example of a G-torsor is the fiber of a principal G-bundle.

[6] Actually, truth be told, we will face situations where the torsors Γ_i of effective superpotentials suffer global twisting analogous to the notion of "twisted vector bundles." We will return to this subtlety later in this overview. Mercifully, the Γ_{ij} introduced in the next paragraph, corresponding to differences of superpotentials, will always be honest torsors.

- $\omega(\gamma, \gamma_i)$: These are the degeneracies of 2d BPS states in vacuum i on $\mathbb{S}$ with 4d gauge charge γ. Their definition is extremely subtle, for reasons explained in §3.5, and these subtleties show that the degeneracy also depends on the choice of γ_i, that is, the "gauge choice" of superpotential on $\mathbb{S}$, or equivalently, of the flux within the solenoid. The dependence on γ_i is affine-linear:

$$\omega(\gamma, \gamma_i + \gamma') = \omega(\gamma, \gamma_i) + \Omega(\gamma)\langle\gamma, \gamma'\rangle. \tag{1.3}$$

In addition to μ and ω there are also generalizations of framed indices. Moreover, all these have further generalizations which describe the response of BPS states to rotation around the surface defect. This leads to a "spin 2d-4d wall-crossing formula," analogous to the "motivic" wall-crossing formula for vanilla BPS states. All of this is described in §4.

All of our new BPS states and their indices undergo wall-crossing phenomena analogous to those already known for pure 2d and 4d theories. In §2.3 we state a precise 2d-4d wall-crossing formula for 2d-4d systems, whose existence was first suggested in [17]. Our formula combines and generalizes the previous wall-crossing formulae of Cecotti and Vafa [32] and Kontsevich and Soibelman [33]. The formula is a kind of "matrix generalization" of the Kontsevich-Soibelman formula. The degeneracies μ are associated with certain finite-dimensional non-diagonal matrices, called $\mathcal{S}$-factors $\mathcal{S}^{\mu}$ (equation (2.27)), while the degeneracies ω are associated with diagonal matrices of symplectomorphisms leading to $\mathcal{K}$-factors $\mathcal{K}^{\omega}$ (equation (2.30)). Technically, these are automorphisms of a noncommutative algebra associated to the vacuum groupoid. They belong to a semidirect product of a group of matrix gauge transformations and a group of Poisson morphisms, acting on matrix-valued functions on an algebraic Poisson torus $\Gamma^* \otimes \mathbb{C}^{\times}$. After stating our wall-crossing formula in §2.3, we systematically analyze its consequences in §2.4, and find that the 2d and 4d degeneracies influence each other's wall-crossing in intricate ways. (When we come to apply these identities to some physical examples we find that they lead to extremely rich spectra of BPS states even in basic examples like those of §8.3.)

The 2d-4d wall-crossing-formula is a mathematically natural synthesis of the known formulae for 2d and 4d theories, but it remains to give it precise physical meaning and justification. The physical meaning of the relevant quantities that enter the formula has already been sketched above and is discussed in detail in §3. The justification for the wall-crossing formula can, as with the pure 4d case, be given in two ways:

- First, one can study interfaces and their framed BPS states. As in [3], invoking the "halo technique" of [34, 3, 35], the consistency of the wall-crossing of the framed BPS indices implies the wall-crossing formula for the unframed indices μ and ω. This story can be found in §4, which also draws on material in Appendices C and D for developing the halo picture. Moreover, §4 begins the development of a spin version of the 2d-4d WCF, although some details of that generalization remain to be worked out.

- Second, one can compactify the theory on a circle and study the geometry implied by constraints of supersymmetry.

We now turn to a detailed discussion of the second justification of the 2d-4d WCF.

This approach was developed for purely 4-dimensional theories in [1]. Let us briefly recall the reasoning here. The argument begins with the observation of Seiberg and Witten [36] that compactification of $\mathcal{T}_{4d}$ on $\mathbb{R}^3 \times S^1$ produces an effective low energy sigma model whose target space is the space $\mathcal{M}$ of (1.2). Supersymmetry implies that $\mathcal{M}$ carries a hyperkähler metric. If the circle has radius R, then when R is large the hyperkähler metric on $\mathcal{M}$ can be computed to exponentially good accuracy to be a semiflat metric. (See §2.4 of [1] or (6.23) below.) The existence of BPS states in the 4-dimensional theory leads, at finite but large values of R, to exponentially small corrections to this semiflat metric. The continuity of these corrections across walls of marginal stability in $\mathcal{B}$ implies the Kontsevich-Soibelman WCF. A key technical step in the argument is the observation that there is a distinguished set of coordinates $\mathcal{Y}_\gamma$ on $\mathcal{M}$, obeying the (twisted) algebra of functions on the algebraic torus $\Gamma^* \otimes \mathbb{C}^\times$, in terms of which the holomorphic symplectic form on $\mathcal{M}$ can be simply written: $\{\mathcal{Y}_\gamma, \mathcal{Y}_{\gamma'}\} \sim \langle\gamma, \gamma'\rangle \mathcal{Y}_\gamma \mathcal{Y}_{\gamma'}$. In particular, upon choosing a basis for Γ we obtain a system of Darboux coordinates on $\mathcal{M}$. It turns out that formulating the quantum corrections of BPS states to the semiflat geometry can be formulated as a Riemann-Hilbert problem whose solution is an integral equation for the $\mathcal{Y}_\gamma$. (See equation (5.21).) This equation is formally identical to the Thermodynamic Bethe Ansatz equation of Zamolodchikov and, although in general no known integrable system is associated to it, we will refer to this equation and its generalizations below as a TBA equation.

The line of argument we have just reviewed can also be generalized to 2d-4d systems, and this we proceed to do in §§5 and 6, which follow in outline much of the development of [1]. We compactify 2d-4d systems by putting $\mathcal{T}_{4d}$ on $\mathbb{R}^3 \times S^1$ as before, but now we wrap one of the dimensions of $\mathbb{S}$ on the circle to produce a 3d sigma model coupled to a 1d line defect. The quantum mechanical system on the line defect has a vector space of ground states which define a vector bundle $V_{\mathbb{S}}$ over $\mathcal{M}$. Constraints of supersymmetry demand that this vector bundle carry a "Berry connection" which is hyperholomorphic. The latter epithet means that the curvature is of type $(1,1)$ for all of the complex structures of the hyperkähler manifold $\mathcal{M}$. This is a generalization of the notion of an instanton on a four-dimensional hyperkähler manifold, which has been studied in the mathematics literature; see, for example, [37, 38, 39, 40, 41, 42]. Once again, the geometry simplifies in the semiflat limit $R \to \infty$. In this case each of the vacua $i \in \mathcal{V}[\mathbb{S}]$ defines a single line bundle $V_i \to \mathcal{M}$ and the Berry connection on $V = \oplus_i V_i$ becomes a diagonal semiflat connection (equation 5.4). The semiflat connection has a natural geometrical interpretation in terms of the relative Poincaré connection on the fiber product of $\mathcal{M} \times_{\mathcal{B}} \widetilde{\mathcal{M}}$ over $\mathcal{B}$. At finite R the semiflat connection receives quantum corrections from all our BPS states: the vanilla BPS states of $\mathcal{T}_{4d}$ as well as the new ones with degeneracies ω and μ. For example, the corrections associated with μ come from the worldlines of solitons wrapping the circle in the cylindrical surface $\mathbb{S}$. Once again, analogs of "Darboux coordinates" $\mathcal{Y}_{\gamma_i}$ and $\mathcal{Y}_{\gamma_{ij'}}$ can be introduced. As in [3] they can be interpreted physically in terms of expectation values of interfaces. They can also be interpreted geometrically as sections of V and $\mathrm{Hom}(V, V')$, respectively, which are holomorphic in all complex structures. Finally, they satisfy a Riemann-Hilbert problem analogous to that in [1]. As in that case we can solve the RH problem through a system of integral equations, (5.21)-(5.25), allowing us to construct $\mathcal{Y}_{\gamma_i}$ and $\mathcal{Y}_{\gamma_{ij'}}$ as an expansion around the explicitly known semiflat sections (5.16). With these in hand we can go on to construct the hyperholomorphic connection, as explained in Appendix E. As in the case of the hyperkähler metric on $\mathcal{M}$, the 2d-4d WCF follows as a consistency condition on the system of equations, (5.21)-(5.25), ensuring smoothness of the metric and connection.

In [1] an important role in the physical argument was played by a local analysis of the quantum corrections coming from mutually local light BPS states. That analysis is repeated and refined in the case of 2d-4d systems in §6. A classification of the possible singularities due to mutually local massless 2d and 4d particles in a single vacuum i is spelled out in §6.1, which also contains a very explicit example of the construction of hyperholomorphic connections on periodic Taub-NUT space. We must point out that the argument of [1] also relied on a general analysis of anomalous Ward identities. The analogous argument for 2d-4d systems is omitted here, but we feel this gap can be easily filled.

There is an interesting and important aspect of the geometry which has thus far been suppressed in this exposition. The bundles $V_{\mathbb{S}}$ are, strictly speaking, in general not bundles at all but rather "twisted bundles." These objects have been encountered previously by physicists, notably in the context of D-branes in the presence of B-fields. There are analogous topological subtleties (also suppressed in our exposition thus far) related to the definitions of the charge torsors Γ_i. The essential point is that the monodromy of the superpotentials, or equivalently of the ν_i sometimes turns out to be fractional, leading to ill-defined Aharonov-Bohm (AB) phases for test particles transported around the surface defect. These "anomalies" are discussed abstractly in §3.4.1 and illustrated concretely in §3.4.3. In the geometrical construction of $V_{\mathbb{S}}$ the twisting of Γ_i leads to a construction of twisted bundles over $\mathcal{M}$, as explained in §5.2.3. In order to have well-defined amplitudes we follow a strategy outlined in §3.6: By gauging a suitable finite abelian group of flavor symmetries we can twist the surface defects so that particles transported around them pick up compensating phases, leading to well-defined AB phases. The cancellation mechanism is described in detail in §3.6.2 and illustrated in a concrete class of models in §7.4. In addition, the relation of global symmetries of three-dimensional sigma models to B-fields and connections on twisted bundles is discussed in Appendix G, which might prove to be of independent interest.

The somewhat abstract structure we have outlined is realized concretely in the theories of class $\mathcal{S}$ introduced in [43, 2, 44]. These are partially twisted $(2,0)$ superconformal theories compactified on $\mathbb{R}^4 \times C$, where C is a punctured Riemann surface and certain codimension two defects are inserted at the punctures. As in our previous papers [2] and [3] our later sections, §§7, 8, and 9, are devoted to the A_1 theories of class $\mathcal{S}$. We leave, yet once more, the full generalization to the higher rank A_n cases for a future occasion. The theories of class $\mathcal{S}$ have the distinguishing property that the Seiberg-Witten moduli spaces $\mathcal{M}$ may be identified with moduli spaces of Hitchin systems on C. (See §7.3.) The Seiberg-Witten curve Σ is a ramified covering of C and is identified with the spectral curve of the Hitchin system, and thus sits in T^*C. The Seiberg-Witten differential λ is then the restriction of the canonical trivialization of the symplectic form on T^*C.

As in [45, 2, 3] there are geometrical realizations of the new BPS states in terms of "WKB curves" or, more generally, "WKB networks." These are certain networks of curves on C, satisfying a first-order differential equation determined by λ. See §7.1. The precise expression of the data of the 2d-4d systems discussed above in terms of the geometry of Σ, C, and WKB networks is explained in §7.2. As in [2] the basic phenomenon of wall-crossing can be understood in terms of the discontinuous change of WKB foliations of C, a topic discussed in §7.5. In 2d-4d systems there is a new wrinkle: Associated to a point $z \in C$ is a canonical surface defect $\mathbb{S}_z$. (If we view theories of class $\mathcal{S}$ as decoupling limits of M5-branes on $\mathbb{R}^4 \times C$, then the surface defects arise from open membranes ending on $\mathbb{S} \times \{z\}$.) In addition to the dependence of our BPS states on $u \in \mathcal{B}$ we can also study their dependence on $z \in C$. There is wall-crossing, and certain special WKB curves known as critical WKB curves play the role of walls of marginal stability. Moreover, the "vector bundles" $V_{\mathbb{S}_z}$ turn out to be the canonically defined universal bundles over $\mathcal{M} \times C$, restricted to z. Actually, as discussed in §7.3, they are in fact twisted bundles. The anomaly in AB phases from the twisting can be cancelled using the mechanism of gauging discrete flavor symmetries, as explained in §7.4.

After explaining the general rules for understanding the BPS spectrum and its wall-crossing for A_1-type examples in §7 we proceed in §8 to look at some detailed examples. We first consider Argyres-Douglas type superconformal theories in §8.1. Even the most trivial examples lead to interesting wall-crossing of framed BPS states, as shown in §8.1.1. Many generic aspects of 2d-4d systems are nicely illustrated in the relatively simple example of the $N = 2$ AD theory in §8.1.2. In particular, this carefully worked example illustrates the compatibility of the 2d-4d wall-crossing formula with the monodromy of the local systems Γ_{ij}. Indeed, one could use the 2d-4d wall-crossing formula to *derive* the monodromy of the local system, without going through the difficult task of analytic continuation of periods! In §8.1.3 we briefly look at 2d-4d wall-crossing for AD theories with $N > 2$. We argue that, while the spectrum of BPS states can be very intricate, the essential qualitative wall-crossing phenomena have already been captured in the examples with small N. In §8.2 we turn to the pure two-dimensional $\mathbb{CP}^1$ sigma model, where we gain new insights though the four-dimensional theory is empty. In particular, we reproduce known results on the BPS spectrum of this model [21] and clarify the relation of the marginal stability phenomena in this model to those occurring in four-dimensional $SU(2)$ theory. Indeed, this two-dimensional example is a useful warmup for the analysis in §8.3 of the $\mathbb{CP}^1$ model coupled to the four-dimensional $SU(2)$ theory without matter. Even this relatively simple example turns out to be rather nontrivial. There are two parameters which control the BPS spectrum: u, which controls the strength of the four-dimensional coupling, and z, which controls the strength of the two-dimensional coupling. The BPS spectrum is a very complicated function of (u, z) and we require the full power of the 2d-4d WCF to conquer it. When u corresponds to strong 4d coupling the BPS spectrum is finite, and, as a function of z, is similar to what one finds in the AD theories. However, when u corresponds to weak 4d coupling the spectrum depends sensitively on z. When z corresponds to strong 2d coupling there are infinitely many walls of marginal stability and infinitely many chambers with different soliton spectra. In each chamber the soliton spectrum is finite, but there is no upper bound on the spectrum. When z corresponds to weak 2d coupling there is an explosion in the complexity of the walls of marginal stability: there are countably many such walls, but uncountably many chambers. It would be very interesting to check these predictions with a weak-coupling field theory analysis.

An application of our work is an algorithm for solving Hitchin's equations (7.13),

$$\begin{aligned} F + R^2[\varphi, \bar{\varphi}] &= 0, \\ \bar{\partial}_A \varphi &= 0, \\ \partial_A \bar{\varphi} &= 0, \end{aligned} \tag{1.4}$$

with rank 1 gauge group on a punctured Riemann surface C. (See §7.3 for notation.) One begins with the spectral double cover $\lambda^2 + \phi_2(z) = 0$, where $\phi_2(z)$ is a meromorphic quadratic differential on C. From the analysis of WKB curves described in §7, (or, in principle, using the spectrum generator (7.38)) one derives the BPS spectrum μ and ω. Then one writes down the integral equations of §5.6. At large values of R the integral equations can be solved by a rapidly convergent iteration. Then, as explained in §9, one expresses the solutions to the integral equations as solutions to the linear problem $(\mathrm{d} + \mathcal{A})\Psi = 0$, where $\mathcal{A}$ is given by (7.15):

$$\mathcal{A} = R\frac{\varphi}{\zeta} + A + R\zeta\bar{\varphi}. \tag{1.5}$$

From Ψ one can determine $\mathcal{A}$, and from $\mathcal{A}$ one clearly obtains a solution to the Hitchin equations. It would be a very useful and interesting check on our logic to work through this algorithm in some explicit examples. We have not yet done this.

We have indicated just a few of the possible future lines of inquiry in §10. We have said this before and might say it again: Despite the length of this paper, we feel we have just scratched the surface of our topic. Time and again, the reader, who will require stamina and dedication to read what follows, will find that we are forced to call a halt to an interesting line of exploration, leaving unexplored fertile fields for future cultivation.

2. Formal statement of wall-crossing formulae

In this section we will formulate a new wall-crossing formula which combines features from both the Cecotti-Vafa wall-crossing formula for 2d $(2,2)$ theories and the Kontsevich-Soibelman wall-crossing formula for 4d $\mathcal{N} = 2$ theories.

2.1 The 2d Cecotti-Vafa wall-crossing formula

The data which enters the CV wall-crossing formula is a finite set $\mathcal{V}$, a "central charge function" $Z : \mathcal{V} \to \mathbb{C}$, and a "degeneracy" $\mu_{ij} \in \mathbb{Z}$ associated to any pair $i, j \in \mathcal{V}$ with $i \neq j$. A physical interpretation of these data will be discussed in §3, where elements of $\mathcal{V}$ will be identified with vacua in an $\mathcal{N} = 2$ supersymmetric $1 + 1$ dimensional QFT.

We consider the function Z as allowed to vary, and the μ_{ij} as functions of Z, i.e. of the values Z_i for $i \in \mathcal{V}$. The dependence of the μ_{ij} on Z is piecewise constant: they are only allowed to jump when Z crosses a "wall of marginal stability," where the values Z_i,

Z_j, Z_k for i, j, k distinct become collinear as points of $\mathbb{C}$. The precise way in which the μ_{ij} jump as Z crosses such a wall is dictated by the wall-crossing formula, which we now describe.

We begin by introducing formal variables X_i and X_{ij}, with the rule that X_{ij} acts by left-multiplication on the X_i as

$$X_{ij} X_k = \delta_{jk} X_i. \tag{2.1}$$

(One could easily realize these product laws in terms of explicit matrices, but our abstract notation here will be convenient for the rest of the paper.) Then we define the "$\mathcal{S}$-factor" $\mathcal{S}^{\mu}_{ij}$ to be a transformation which acts on the X_k as left-multiplication by $1 - \mu_{ij} X_{ij}$, i.e.

$$\mathcal{S}^{\mu}_{ij} : X_k \to X_k - \delta_{jk} \mu_{ij} X_i. \tag{2.2}$$

Next, to each pair (i, j) with $\mu_{ij} \neq 0$ we associate a "BPS ray" ℓ_{ij} in the complex plane, with slope given by the phase of $Z_{ij} := Z_i - Z_j$:

$$\ell_{ij} = Z_{ij} \mathbb{R}_- \subset \mathbb{C}. \tag{2.3}$$

Now choose some convex angular sector $\lhd$ in the complex plane, with apex at the origin. The basic actor in the wall-crossing formula is the composition $A(\lhd)$ of all the $\mathcal{S}$-factors corresponding to BPS rays lying in $\lhd$:

$$A(\lhd) =: \prod_{i,j:\ell_{ij}\subset\lhd} \mathcal{S}^{\mu}_{ij} : \quad . \tag{2.4}$$

Here the normal-ordering symbols mean that the product is ordered by the phase of Z_{ij}, i.e., reading from left to right in the product we encounter the factors associated with the BPS rays ℓ_{ij} successively in the counterclockwise direction. See Figure 1. The statement of the wall-crossing formula is: $A(\lhd)$ *is constant under variation of* Z*, as long as no BPS rays cross the boundary of* $\lhd$.


ℓ_{23}
ℓ_{13}
ℓ_{12}
$\lhd$
$A(\lhd) = \mathcal{S}^{\mu_{12}}_{12} \mathcal{S}^{\mu_{13}}_{13} \mathcal{S}^{\mu_{23}}_{23}$

Figure 1: The definition of $A(\lhd)$ as a product of $\mathcal{S}$-factors associated to the BPS rays ℓ_{ij}.

Let us see how this WCF works in practice. Suppose we have three rays ℓ_{ij}, ℓ_{ik}, ℓ_{jk} which are close together in the complex plane, ordered say counterclockwise (with i, j, k

pairwise distinct). Choose the sector $\lhd$ to include these three and no others. Now suppose we vary Z until we reach a wall of marginal stability, where Z_{ij} and Z_{jk} (and hence Z_{ik}) have the same phase. Upon crossing this wall the rays ℓ_{ij}, ℓ_{ik}, ℓ_{jk} switch from counterclockwise to clockwise ordering. Then the WCF says we should compare the composition $\mathcal{S}^{\mu}_{ij}\mathcal{S}^{\mu}_{ik}\mathcal{S}^{\mu}_{jk}$ on one side of the wall and $\mathcal{S}^{\mu'}_{jk}\mathcal{S}^{\mu'}_{ik}\mathcal{S}^{\mu'}_{ij}$ on the other side:

$$\begin{aligned}\mathcal{S}^{\mu}_{ij}\mathcal{S}^{\mu}_{ik}\mathcal{S}^{\mu}_{jk} &= (1-\mu_{ij}X_{ij})(1-\mu_{ik}X_{ik})(1-\mu_{jk}X_{jk})\\ &= 1-\mu_{ij}X_{ij}-(\mu_{ik}-\mu_{ij}\mu_{jk})X_{ik}-\mu_{jk}X_{jk}, \qquad (2.5)\\ \mathcal{S}^{\mu'}_{jk}\mathcal{S}^{\mu'}_{ik}\mathcal{S}^{\mu'}_{ij} &= (1-\mu'_{jk}X_{jk})(1-\mu'_{ik}X_{ik})(1-\mu'_{ij}X_{ij})\\ &= 1-\mu'_{ij}X_{ij}-\mu'_{ik}X_{ik}-\mu'_{jk}X_{jk}. \qquad (2.6)\end{aligned}$$

Hence we find

$$\mu'_{ij} = \mu_{ij}, \qquad (2.7)$$

$$\mu'_{ik} = \mu_{ik}-\mu_{ij}\mu_{jk}, \qquad (2.8)$$

$$\mu'_{jk} = \mu_{jk}. \qquad (2.9)$$

This is precisely the wall-crossing formula of Cecotti and Vafa [20]. A simple example is $\mu_{ij}=\mu_{jk}=\mu_{ik}=1$, which gives $\mu'_{ij}=\mu'_{jk}=1$ but $\mu'_{ik}=0$.

Since μ_{ij} enters the wall-crossing formula only through $\mu_{ij}X_{ij}$ it is sometimes useful to give an equivalent presentation of the wall-crossing formula, where some signs are shifted from the μ_{ij} to the definition of X_{ij}.

2.2 The 4d Kontsevich-Soibelman wall-crossing formula

The data which enters the KS wall-crossing formula is a lattice Γ of "charges" with an integer-valued bilinear antisymmetric pairing $\langle\cdot,\cdot\rangle$ (the "intersection product"), a central charge function $Z:\Gamma\to\mathbb{C}$, linear on Γ, and a set of "degeneracies" $\Omega(\gamma)\in\mathbb{Z}$ associated to charges $\gamma\in\Gamma$.

As in the 2d case, the $\Omega(\gamma)$ depend on the function Z in a piecewise constant fashion. They may jump only when Z crosses a "wall of marginal stability" where the phases of Z_γ and $Z_{\gamma'}$ for two linearly independent γ, γ' become aligned. The wall-crossing formula dictates precisely how the $\Omega(\gamma)$ jump when Z crosses such a wall.

In order to formulate the wall-crossing formula it is useful to introduce formal variables X_γ for $\gamma\in\Gamma$, with a twisted multiplication rule

$$X_\gamma X_{\gamma'} = (-1)^{\langle\gamma,\gamma'\rangle}X_{\gamma+\gamma'}. \qquad (2.10)$$

Then we introduce the "$\mathcal{K}$-factor" $\mathcal{K}^{\Omega}_{\gamma}$, which acts on the $X_{\gamma'}$ by[7]

$$\mathcal{K}^{\Omega}_{\gamma}: X_{\gamma'}\to(1-X_\gamma)^{\langle\gamma',\gamma\rangle\Omega(\gamma)}X_{\gamma'}. \qquad (2.11)$$

---

[7] In previous papers on the subject we defined $\mathcal{K}_\gamma$ directly rather than $\mathcal{K}^{\Omega}_{\gamma}$. We do not do this here because $\mathcal{K}_\gamma$ itself does not have a natural extension to the 2d-4d setting; $\mathcal{K}^{\Omega}_{\gamma}$ turns out to be the more natural object.

To each γ such that $\Omega(\gamma) \neq 0$ we associate a “BPS ray” ℓ_γ in the complex plane, with slope given by the phase of Z_γ:

$$\ell_\gamma = Z_\gamma \mathbb{R}_- \subset \mathbb{C}. \tag{2.12}$$

The basic actor in the wall-crossing formula is the composition $A(\sphericalangle)$ of all the $\mathcal{K}$-factors corresponding to BPS rays lying the sector $\sphericalangle$ of the complex plane:

$$A(\sphericalangle) = :\prod_{\gamma : \ell_\gamma \subset \sphericalangle} \mathcal{K}^{\Omega}_{\gamma} : \ . \tag{2.13}$$

As in the 2d case, the ordering is such that if $\ell_{\gamma_1}, \ell_{\gamma_2}$ are in counterclockwise order then $\mathcal{K}^{\Omega}_{\gamma_1}$ is to the left of $\mathcal{K}^{\Omega}_{\gamma_2}$. The statement of the wall-crossing formula [33] is: $A(\sphericalangle)$ *is constant under variation of* Z*, as long as no BPS rays cross the boundary of* $\sphericalangle$. All this is completely parallel to the 2d case.

For convenience of our discussion in §§7 and 8 we now give two standard examples of this wall-crossing formula in action. Consider the simple situation where we begin with two BPS rays ℓ_γ and $\ell_{\gamma'}$, with $\Omega(\gamma) = \Omega(\gamma') = 1$, and no BPS rays lying between ℓ_γ and $\ell_{\gamma'}$. Then we adjust Z until the two BPS rays collide and cross one another. After the collision we will have a more complicated set of BPS rays and degeneracies. What we get precisely depends heavily on the value of $\langle\gamma, \gamma'\rangle$. First, suppose $\langle\gamma, \gamma'\rangle = 1$. In this case we have

$$\mathcal{K}^{\Omega}_{\gamma'} \mathcal{K}^{\Omega}_{\gamma} = \mathcal{K}^{\Omega'}_{\gamma} \mathcal{K}^{\Omega'}_{\gamma+\gamma'} \mathcal{K}^{\Omega'}_{\gamma'}, \tag{2.14}$$

where $\Omega'(\gamma) = \Omega'(\gamma') = \Omega'(\gamma+\gamma') = 1$. This identity is simple enough to be checked directly; just consider the action of $\mathcal{K}^{\Omega}_{\gamma'} \mathcal{K}^{\Omega}_{\gamma}$,

$$(X_\gamma; X_{-\gamma'}) \overset{\mathcal{K}^{\Omega}_{\gamma}}{\mapsto} (X_\gamma; X_{-\gamma'} + X_{\gamma-\gamma'}) \overset{\mathcal{K}^{\Omega}_{\gamma'}}{\mapsto} (X_\gamma + X_{\gamma+\gamma'}; X_{-\gamma'} + X_{\gamma-\gamma'} + X_\gamma), \tag{2.15}$$

and compare it with the action of $\mathcal{K}^{\Omega'}_{\gamma} \mathcal{K}^{\Omega'}_{\gamma+\gamma'} \mathcal{K}^{\Omega'}_{\gamma'}$,

$$\begin{aligned}(X_\gamma; X_{-\gamma'}) &\overset{\mathcal{K}^{\Omega'}_{\gamma'}}{\mapsto} (X_\gamma + X_{\gamma+\gamma'}; X_{-\gamma'}) \overset{\mathcal{K}^{\Omega'}_{\gamma+\gamma'}}{\mapsto} (X_\gamma + X_{2\gamma+\gamma'} + X_{\gamma+\gamma'}; X_{-\gamma'} + X_\gamma) \\ &\overset{\mathcal{K}^{\Omega'}_{\gamma}}{\mapsto} (X_\gamma + X_{\gamma+\gamma'}; X_{-\gamma'} + X_{\gamma-\gamma'} + X_\gamma). \end{aligned} \tag{2.16}$$

The fact that the two agree is equivalent to (2.14). For a more interesting example, suppose instead $\langle\gamma, \gamma'\rangle = 2$. Then

$$\mathcal{K}^{\Omega}_{\gamma'} \mathcal{K}^{\Omega}_{\gamma} = \mathcal{K}^{\Omega'}_{\gamma} \mathcal{K}^{\Omega'}_{2\gamma+\gamma'} \mathcal{K}^{\Omega'}_{3\gamma+2\gamma'} \cdots \mathcal{K}^{\Omega'}_{\gamma+\gamma'} \cdots \mathcal{K}^{\Omega'}_{2\gamma+3\gamma'} \mathcal{K}^{\Omega'}_{\gamma+2\gamma'} \mathcal{K}^{\Omega'}_{\gamma'} \tag{2.17}$$

where $\Omega'((n+1)\gamma + n\gamma') = \Omega'(n\gamma + (n+1)\gamma') = 1$ for all $n \geq 0$, $\Omega'(\gamma+\gamma') = -2$. This identity is less trivial to prove; see Appendix A of [1] for an elementary proof.

2.3 The 2d-4d wall-crossing formula

Now we are ready to give the new wall-crossing formula which is the subject of this paper. It combines the features of the wall-crossing formulae of §§2.1 and 2.2, but it involves some new features. Heuristically, it is a kind of "fiber-product" of the two previously known wall-crossing formulae.

Our data include both the finite set $\mathcal{V}$ and the lattice Γ with integral antisymmetric pairing $\langle\cdot,\cdot\rangle$. In addition, we will need new charge "lattices," an extension of the central charge function to these "lattices," a new set of BPS degeneracies, and, finally, a new $\mathbb{Z}_2$-valued twisting function. The physical interpretation of these data is explained in §3. We now describe the new data more precisely.

The first piece of new data is a set of distinct Γ-*torsors* Γ_i, one for each $i \in \mathcal{V}$. We will usually use the notation γ_i to denote a generic element of Γ_i (but the reader should be warned that on several occasions γ_i will refer to some special or specific element of Γ_i). Each Γ_i is both a left- and right- Γ-torsor, with $\gamma + \gamma_i = \gamma_i + \gamma$.

We also introduce additional sets Γ_{ij}, defined as follows: an element $\gamma_{ij} \in \Gamma_{ij}$ is a formal difference $\gamma_i - \gamma_j$ of elements of Γ_i and Γ_j, modulo the equivalence $(\gamma_i + \gamma) - (\gamma_j + \gamma) = \gamma_i - \gamma_j$. Γ_{ij} is again a Γ-torsor, with $\gamma + \gamma_{ij} = \gamma_{ij} + \gamma$. A convenient notation for this definition is

$$\Gamma_{ij} = \Gamma_i - \Gamma_j. \tag{2.18}$$

One can identify Γ_{ii} canonically with Γ, and it will sometimes be convenient to do so. [8] Then, to an element $\gamma_{ij} \in \Gamma_{ij}$ we can associate a unique element $(-\gamma_{ij}) \in \Gamma_{ji}$ so that $\gamma_{ij} + (-\gamma_{ij}) = 0 \in \Gamma_{ii}$.

There are naturally defined addition operations $\Gamma_{ij} \times \Gamma_j \to \Gamma_i$ and $\Gamma_{ij} \times \Gamma_{jk} \to \Gamma_{ik}$. We will write these operations as $+$, thus $\gamma_{ij} + \gamma_j \in \Gamma_i$, but the reader should be warned that $+$ is not symmetric; for example, $\gamma_j + \gamma_{ij}$ is not defined when $i \neq j$.



Figure 2: A partial depiction of the groupoid $\mathbb{V}$, in case $\mathcal{V}$ has three elements, here labeled i, j, k. Each morphism space is a torsor for Γ, of which we have shown only a single element.

---

[8] This leads to the slightly awkward notational point that γ_{ii} is not zero. It is generically $\gamma_i - \gamma_i'$ where γ_i, γ_i' are both in Γ_i, i.e. it is a generic element of Γ.

The situation is nicely summarized by declaring that the first piece of data is a groupoid $\mathbb{V}$ with the following properties. For each $i \in \mathcal{V}$, $\mathbb{V}$ has a corresponding object. The set of morphisms $\mathrm{Hom}(i,j)$ from i to j is a Γ-torsor and identified with Γ_{ij}. The composition of morphisms is given by the addition laws we have written above: In addition there is one more object denoted o,[9] making $\mathbb{V}$ into a pointed groupoid. We identify $\mathrm{Hom}(i,o) = \Gamma_i$. Note that $\mathrm{Hom}(o,i)$ is another torsor which gives independent meaning to elements $-\gamma_i \in -\Gamma_i$ so that $\gamma_i + (-\gamma_j) \in \Gamma_{ij}$. Thus, e.g. the composition of $\gamma_j \in \Gamma_j = \mathrm{Hom}(j,o)$ and $\gamma_{ij} \in \Gamma_{ij} = \mathrm{Hom}(i,j)$ is $\gamma_{ij} + \gamma_j \in \Gamma_i = \mathrm{Hom}(i,o)$ while $-\gamma_i + \gamma_{ij} \in -\Gamma_j$. The automorphism group $\mathrm{Hom}(i,i)$ of each object (including $i = o$) is canonically isomorphic to Γ. See Figure 2.

Our second new piece of data is an extension of the central charge function Z from Γ to all the Γ_i, which is compatible with the Γ-action. Thus we have $Z : \amalg_i \Gamma_i \to \mathbb{C}$ which is affine-linear in the sense that $Z_{\gamma+\gamma_i} = Z_\gamma + Z_{\gamma_i}$. Once we have this extension, we automatically get a further extension of Z to the Γ_{ij}, namely, if $\gamma_{ij} = \gamma_i - \gamma_j$ then we define $Z_{\gamma_{ij}} := Z_{\gamma_i} - Z_{\gamma_j}$. Then we have

$$Z_{a+b} = Z_a + Z_b \tag{2.19}$$

whenever $a+b$ is defined. (Here and below, we use the notation $a, b, c, \dots$ in statements which are true when $a, b, c, \dots$ are valued in any of Γ, Γ_i or Γ_{ij}.) In the categorical language mentioned above, giving Z is equivalent to giving a homomorphism from the groupoid $\mathbb{V}$ to $\mathbb{C}$.

Our third new piece of data is a new collection of "BPS degeneracies". As in the 4d case we have "4d degeneracies" $\Omega(\gamma) \in \mathbb{Z}$ for $\gamma \in \Gamma$. It is convenient to define a function $\omega : \Gamma \times \Gamma \to \mathbb{Z}$ by

$$\omega(\gamma, \gamma') := \Omega(\gamma)\langle \gamma, \gamma' \rangle. \tag{2.20}$$

We then introduce new "mixed degeneracies" $\omega : \Gamma \times \amalg_i \Gamma_i \to \mathbb{Z}$, [10] subject to the condition

$$\omega(\gamma, \gamma_i + \gamma') = \omega(\gamma, \gamma_i) + \Omega(\gamma)\langle \gamma, \gamma' \rangle. \tag{2.21}$$

(Because of this condition, $\omega(\gamma, \cdot)$ is determined once we give a single $\omega(\gamma, \gamma_i)$ for each i.) We also define $\omega : \Gamma \times \amalg_{i,j} \Gamma_{ij} \to \mathbb{Z}$ by

$$\omega(\gamma, \gamma_i - \gamma_j) := \omega(\gamma, \gamma_i) - \omega(\gamma, \gamma_j). \tag{2.22}$$

Then for any two "charges" a, b, each belonging to Γ, Γ_i, or Γ_{ij}, we have

$$\omega(\gamma, a+b) = \omega(\gamma, a) + \omega(\gamma, b) \tag{2.23}$$

so long as $a+b$ is defined. In the categorical language, for each γ, giving $\omega(\gamma, \cdot)$ is equivalent to giving a homomorphism $\mathbb{V} \to \mathbb{Z}$.

Turning to 2d particles, we do not simply take over a set of "2d degeneracies" $\mu_{ij} \in \mathbb{Z}$ as in the pure 2d story: rather we promote these to a set of $\mu(\gamma_{ij}) \in \mathbb{Z}$, defined for each $\gamma_{ij} \in \Gamma_{ij}$ where $i \neq j$.

---

[9] o will correspond physically to the "empty surface defect."

[10] In some situations we should relax the requirement that ω is $\mathbb{Z}$-valued; see the end of this subsection.

As in the pure 2d and 4d cases, the BPS degeneracies μ, ω depend on Z, but this dependence is completely determined by a wall-crossing formula, to be stated below.

Our fourth, and final, piece of data is a "twisting" function $\sigma(a,b)$, defined whenever $a+b$ is defined, and valued in $\{\pm 1\}$. We require σ to obey a cocycle condition,[11]

$$\sigma(a,b)\sigma(a+b,c) = \sigma(a,b+c)\sigma(b,c). \tag{2.24}$$

In order to formulate the wall-crossing formula we introduce formal variables X_a, where, as above, a is an element of Γ, Γ_i or Γ_{ij}. We define a twisted multiplication rule:

$$X_a X_b = \begin{cases} \sigma(a,b) X_{a+b} & \text{if } a+b \text{ is defined,} \\ 0 & \text{otherwise.} \end{cases} \tag{2.25}$$

This product is associative thanks to (2.24). In general $a+b$ and $b+a$ are not simultaneously defined, so the product is *noncommutative*. However, if $a+b$ and $b+a$ are both defined we assume $\sigma(a,b) = \sigma(b,a)$ so in this case $X_a X_b = X_b X_a$. We will also assume that $\sigma(\gamma,\gamma') = (-1)^{\langle\gamma,\gamma'\rangle}$ to insure compatibility with the pure 4d case. In the categorical language mentioned above, we are essentially considering the twisted groupoid algebra $\mathbb{C}[\mathbb{V}]$.[12]

The wall-crossing formula controls the dependence of the BPS degeneracies μ and ω on Z. The degeneracies are piecewise constant as functions of Z. They jump only at the "walls of marginal stability" where the phases of Z_a and Z_b become equal for some pair (a,b) with $a \neq b$, where each of a, b separately belongs either to Γ or to some Γ_{ij} (with the proviso that if both a and b belong to Γ then they are linearly independent.) We define rays ℓ_γ and $\ell_{\gamma_{ij}}$ in the complex plane, with slopes given respectively by the phases of Z_γ or $Z_{\gamma_{ij}}$:

$$\ell_\gamma = Z_\gamma \mathbb{R}_-, \qquad \ell_{\gamma_{ij}} = Z_{\gamma_{ij}} \mathbb{R}_-. \tag{2.26}$$

We call ℓ_γ a *BPS $\mathcal{K}$-ray* if $\omega(\gamma,\cdot) \neq 0$, and likewise $\ell_{\gamma_{ij}}$ a *BPS $\mathcal{S}$-ray* if $\mu_{\gamma_{ij}} \neq 0$. BPS degeneracies can jump only when Z is such that two BPS rays become coincident.

As in the Cecotti-Vafa and Kontsevich-Soibelman WCF reviewed above, we will attach certain "factors" (automorphisms of $\mathbb{C}[\mathbb{V}]$) to the BPS rays. The factors associated to the $\mathcal{S}$-rays $\ell_{\gamma_{ij}}$, called "$\mathcal{S}$-factors" and denoted $\mathcal{S}^\mu_{\gamma_{ij}}$, are defined by

$$\mathcal{S}^\mu_{\gamma_{ij}} : X_a \to (1 - \mu(\gamma_{ij}) X_{\gamma_{ij}}) X_a (1 + \mu(\gamma_{ij}) X_{\gamma_{ij}}). \tag{2.27}$$

In particular the $\mathcal{S}$-factor acts trivially on X_γ and

$$\mathcal{S}^\mu_{\gamma_{ij}} : X_{\gamma_k} \mapsto \begin{cases} X_{\gamma_k} & k \neq j \\ X_{\gamma_j} - \mu(\gamma_{ij})\sigma(\gamma_{ij},\gamma_j) X_{\gamma_{ij}+\gamma_j} & k = j \end{cases} \tag{2.28}$$

---

[11] σ is very similar to the data one would associate to a twisting of K theory on the groupoid $\mathbb{V}$. However, although $\mathbb{V}$ is equivalent to $(pt)/\!\!/\Gamma$ as a groupoid, the extra structure of Z does not respect this equivalence of categories.

[12] The X_γ as defined so far do not strictly belong to $\mathbb{C}[\mathbb{V}]$, but one can identify X_γ with $X_{\gamma_{oo}} + \sum_i X_{\gamma_{ii}} \in \mathbb{C}[\mathbb{V}]$, where each $\gamma_{ii} \in \Gamma_{ii}$ (including $i = o$) is the one corresponding to $\gamma \in \Gamma$. This identification is compatible with everything we will do below.

$$\mathcal{S}^{\mu}_{\gamma_{ij}} : X_{-\gamma_k} \mapsto \begin{cases} X_{-\gamma_k} & k \neq i \\ X_{-\gamma_i} + \mu(\gamma_{ij})\sigma(-\gamma_i, \gamma_{ij}) X_{-\gamma_i + \gamma_{ij}} & k = i \end{cases} \tag{2.29}$$

which the reader should compare with (2.2). The factors associated to the $\mathcal{K}$-rays ℓ_γ, called "$\mathcal{K}$-factors" and denoted $\mathcal{K}^{\omega}_{\gamma}$, are defined by

$$\mathcal{K}^{\omega}_{\gamma} : X_a \to (1 - X_\gamma)^{-\omega(\gamma, a)} X_a. \tag{2.30}$$

Note that when $a = \gamma'$ this transformation reduces to (2.11): this is the reason why our wall-crossing formula will reduce to the 4d wall-crossing formula as far as the 4d degeneracies Ω are concerned.

We can now state our new wall-crossing formula in a way closely analogous to the 2d and 4d cases. For an angular sector $\sphericalangle$ we define an automorphism of the groupoid algebra,

$$A(\sphericalangle) = : \prod_{\gamma : \ell_\gamma \subset \sphericalangle} \mathcal{K}^{\omega}_{\gamma} \prod_{\gamma_{ij} : \ell_{\gamma_{ij}} \subset \sphericalangle} \mathcal{S}^{\mu}_{\gamma_{ij}} : \tag{2.31}$$

where the normal ordering symbol indicates that the factors — be they of type $\mathcal{K}$ or $\mathcal{S}$ — are ordered so that reading from left to right we encounter factors associated with rays successively in the counterclockwise direction. Then the new wall-crossing formula says: *$A(\sphericalangle)$ is constant, as long as no BPS rays cross the boundary of $\sphericalangle$.*

In later sections of the paper we will describe physical realizations of this set of axioms. We will also encounter a slightly "twisted" version of them, which originates from the observation that one can consistently study the wall-crossing of the quantities $\mu(\gamma_{ij})$, $\Omega(\gamma)$ and $\omega(\gamma, \gamma_{ij})$ as $Z_{\gamma_{ij}}$ vary, with no reference to the Z_{γ_i}, $\omega(\gamma, \gamma_i)$ or even Γ_i. Indeed, in some physical examples, certain anomalies can prevent us from defining Γ_i, Z_{γ_i}, $\omega(\gamma, \gamma_i)$ unambiguously, even though Γ_{ij}, $Z_{\gamma_{ij}}$, $\omega(\gamma, \gamma_{ij})$, $\mu(\gamma_{ij})$, $\Omega(\gamma)$ are perfectly well-defined and satisfy the 2d-4d wall-crossing axioms. See §3.4.1 below. In these situations it appears to be appropriate simply to require integrality of the $\omega(\gamma, \gamma_{ij})$ and in fact it might not be possible to define $\omega(\gamma, \gamma_i)$ as integers.

2.3.1 Automorphisms of wall-crossing identities

If α is an automorphism or anti-automorphism of the groupoid algebra $\mathbb{C}[\mathbb{V}]$ then, given a wall-crossing identity, we can conjugate the factors by α to produce another wall-crossing identity.

The simplest example of this is to consider the transformation $\alpha(X_a) = (-1)^{c_a} X_a$, which in general changes $\sigma(a, b)$ by a coboundary but will be an automorphism if $c_{a+b} = c_a + c_b \bmod 2$. In particular, if degeneracies μ, ω satisfy the wall-crossing formula then so do $\tilde{\mu}, \omega$ where

$$\tilde{\mu}(\gamma_{ij}) = (-1)^{c_i - c_j} \mu(\gamma_{ij}) \tag{2.32}$$

for integer c_i.

As a second example, consider the anti-automorphism of $\mathbb{C}[\mathbb{V}]$ defined by $\alpha : X_a \mapsto X_{-a}$. (Note that if $a + b$ is composable then $(-b) + (-a)$ is composable, so α must be an anti-automorphism.) One can check that $\alpha \mathcal{K}^{\omega}_{\gamma} \alpha^{-1} = \mathcal{K}^{\tilde{\omega}}_{-\gamma}$ and $\alpha \mathcal{S}^{\mu}_{\gamma_{ij}} \alpha^{-1} = \mathcal{S}^{\tilde{\mu}}_{-\gamma_{ij}}$ with

$$\tilde{\mu}(\gamma_{ij}) = -\mu(-\gamma_{ij}), \qquad \tilde{\omega}(\gamma, a) = \omega(-\gamma, -a), \tag{2.33}$$

giving a second automorphism of the wall-crossing formula.

In the physical realizations, the second automorphism above is closely related to CPT invariance. CPT takes $a \to -a$, but also takes $Z \to \bar{Z}$, where Z is the central charge. Therefore, CPT reverses the clockwise ordering of BPS rays. On the other hand, given a wall-crossing identity, written as an equality of a certain word in $\mathcal{K}$ and $\mathcal{S}$ factors with the identity operator, we may always take the inverse of that word to produce another identity. Then the order of the BPS rays will be reversed. Thus we may define the CPT conjugate of a wall-crossing identity as the one obtained by taking the inverse and conjugating by $\alpha : X_a \mapsto X_{-a}$.

2.4 Examples

In this section we give a few simple examples of wall-crossing formulae. The general wall of marginal stability occurs when a ray of the form $\ell_{\gamma_{ij}}$ becomes collinear with a ray of the form $\ell_{\gamma_{kl}}$. We can consider several examples according to how the indices i, j, k, l are related.

First, if none of the indices i, j, k, l coincide, then the charges $n\gamma_{ij} + m\gamma_{kl}$ are not defined, so there are only two BPS rays in the game. We associate two commuting $\mathcal{S}$-factors with these two rays, and the wall-crossing is trivial:

$$\mathcal{S}^{\mu}_{\gamma_{ij}} \mathcal{S}^{\mu}_{\gamma_{kl}} = \mathcal{S}^{\mu'}_{\gamma_{kl}} \mathcal{S}^{\mu'}_{\gamma_{ij}} \tag{2.34}$$

with $\mu' = \mu$.

Next, suppose there are three distinct indices and one pair of indices coincides. The cases $j = k$ and $i = l$ are related by wall-crossing, and in this case the wall-crossing formula is essentially identical to the pure 2d case: we must take into account a third BPS ray associated to the charge $\gamma_{il} := \gamma_{ij} + \gamma_{jl}$, and then we have

$$\mathcal{S}^{\mu}_{\gamma_{ij}} \mathcal{S}^{\mu}_{\gamma_{il}} \mathcal{S}^{\mu}_{\gamma_{jl}} = \mathcal{S}^{\mu'}_{\gamma_{jl}} \mathcal{S}^{\mu'}_{\gamma_{il}} \mathcal{S}^{\mu'}_{\gamma_{ij}} \tag{2.35}$$

with $\mu'(\gamma_{jl}) = \mu(\gamma_{jl})$, $\mu'(\gamma_{ij}) = \mu(\gamma_{ij})$ and $\mu'(\gamma_{il}) = \mu(\gamma_{il}) - \sigma(\gamma_{ij}, \gamma_{jl})\mu(\gamma_{ij})\mu(\gamma_{jl})$.

If instead there are three distinct indices with $i = j$ or $j = l$, there cannot be a third BPS ray, and the commuting $\mathcal{S}$-factors pass through each other as in (2.34).

Next, suppose there are three distinct indices and, say, $i \neq j \neq k = l$, so we can set $\gamma_{kk} = \gamma \in \Gamma$. Now there can be intermediate rays associated to charges $n\gamma + \gamma_{ij}$ for $n > 0$. The wall-crossing formula then states that

$$\mathcal{K}^{\omega}_{\gamma} \prod_{n\searrow} \mathcal{S}^{\mu}_{\gamma_{ij}+n\gamma} = \prod_{n\nearrow} \mathcal{S}^{\mu'}_{\gamma_{ij}+n\gamma} \mathcal{K}^{\omega'}_{\gamma} \tag{2.36}$$

where the range on n is from 0 to $+\infty$ and the product $\prod_{n\nearrow}$ means that as we read from left to right n is increasing.[13] Applying this identity on the generators X_{γ_k} for $k \neq j$ shows that $\omega'(\gamma, \gamma_k) = \omega(\gamma, \gamma_k)$ for $k \neq j$ and $k \neq -i$. Then applying the formula to X_{γ_j} we find that in fact $\omega' = \omega$ always and the μ' are determined in terms of ω and μ. To state the result it is useful to define a power series in X_γ:

---

[13] In fact, in this example the $\mathcal{S}$-factors actually commute with each other.

$$\Sigma_{ij} := \sum_{n=0}^{\infty} \frac{\mu(\gamma_{ij} + n\gamma)}{\sigma(\gamma_{ij}, \gamma)^n} X_\gamma^n, \tag{2.37}$$

with a similar definition for Σ'_{ij}. Then we have

$$\Sigma'_{ij} = \Sigma_{ij}(1 - X_\gamma)^{-\omega(\gamma, \gamma_{ij})}. \tag{2.38}$$

A few special cases are worth noting separately. Suppose for example that we take $\mu(\gamma_{ij}) = 1$ and $\omega(\gamma, \gamma_{ij}) = -1$ with all other μ, ω vanishing. Then $\Sigma_{ij} = 1$, and (2.38) says $\Sigma'_{ij} = 1 - X_\gamma$: so (2.36) specializes to

$$\mathcal{K}_\gamma^\omega \mathcal{S}_{\gamma_{ij}}^\mu = \mathcal{S}_{\gamma_{ij}}^{\mu'} \mathcal{S}_{\gamma_{ij}+\gamma}^{\mu'} \mathcal{K}_\gamma^{\omega'} \tag{2.39}$$

where $\mu'(\gamma_{ij}) = 1$, $\mu'(\gamma_{ij} + \gamma) = -\sigma(\gamma_{ij}, \gamma)$, and $\omega' = \omega$. We will use this example, in the form (8.2a) and (8.2b), extensively in §8.

Let us now suppose that there are only two distinct indices, i.e. two pairs of indices coincide. There are three cases to consider. If we have $\ell_{\gamma_{ij}}$ collinear with $\ell_{\gamma'_{ij}}$, then since $\gamma'_{ij} = \gamma_{ij} + \gamma$ for some $\gamma \in \Gamma$, it follows that Z_γ or $Z_{-\gamma}$ is collinear with $Z_{\gamma_{ij}}$, and we return to the case of (2.36).

If $i = j$ and $k = l$, then we have BPS rays ℓ_{γ^1} and ℓ_{γ^2} crossing for $\gamma^1, \gamma^2 \in \Gamma$. In this case all the BPS rays are $\mathcal{K}$-rays and the wall-crossing formula says

$$\prod_{\frac{n}{m}\nearrow} \mathcal{K}^\omega_{n\gamma^1 + m\gamma^2} = \prod_{\frac{n}{m}\searrow} \mathcal{K}^{\omega'}_{n\gamma^1 + m\gamma^2}. \tag{2.40}$$

Here $\sigma(a, b)$ will have to be generic, as $\langle \gamma^1, \gamma^2 \rangle$ can be odd. This will not affect the present calculation. Acting on X_γ for $\gamma \in \Gamma$ this reduces precisely to the 4d wall-crossing formula, and hence implies the same relations between Ω and Ω' that we had in the pure 4d discussion. As we have noted, though, ω and ω' contain a bit more information than Ω and Ω'. To determine the relation between ω and ω' we should act with both sides of (2.40) on some X_{γ_i}. For example, consider again the case we discussed around (2.14), where $\langle \gamma^1, \gamma^2 \rangle = 1$ and $\Omega(\gamma^1) = \Omega(\gamma^2) = 1$, with all other $\Omega(n\gamma_1 + m\gamma_2) = 0$. Then the 4d wall-crossing formula says that $\Omega'(\gamma^1) = \Omega'(\gamma^2) = \Omega'(\gamma^1 + \gamma^2) = 1$, with all other $\Omega'(n\gamma_1 + m\gamma_2) = 0$. Suppose moreover that the only nonvanishing ω are $\omega(\gamma^1, \cdot)$ and $\omega(\gamma^2, \cdot)$. Then (2.40) is satisfied if and only if ω' are

$$\omega'(\gamma^1, \cdot) = \omega(\gamma^1, \cdot), \tag{2.41}$$

$$\omega'(\gamma^2, \cdot) = \omega(\gamma^2, \cdot), \tag{2.42}$$

$$\omega'(\gamma^1 + \gamma^2, \cdot) = \omega(\gamma^1, \cdot) + \omega(\gamma^2, \cdot) \tag{2.43}$$

with all other $\omega'(m\gamma^1 + n\gamma^2, \cdot)$ vanishing. To prove this, first note that given (2.21) and our knowledge of Ω and Ω', these relations are equivalent to the ones where $\cdot$ is replaced by a single $\gamma_i^0 \in \Gamma_i$. Then choose γ_i^0 with $\omega(\gamma^1, \gamma_i^0) = \omega(\gamma^2, \gamma_i^0) = 0$. (Concretely, such a γ_i^0 can be produced by starting with any generic γ_i and then forming $\gamma_i^0 = \gamma_i - \omega(\gamma^1, \gamma_i)\gamma^2 + \omega(\gamma^2, \gamma_i)\gamma^1$.) The check is then trivial, by comparing the action of both sides of (2.40) on $X_{\gamma_i^0}$. In many concrete examples a similar trick can be employed: find a convenient reference vector in each Γ_i which simplifies ω.

The third case with two distinct indices occurs when $\ell_{\gamma_{ij}}$ coincides with $\ell_{\gamma_{ji}}$. Note that $\gamma_{ij} + \gamma_{ji} = \gamma \in \Gamma$ so that all the rays for $\gamma_{ij} + n\gamma$ and $\gamma_{ji} + n\gamma$ with $n \geq 0$ coincide, and generically there are no other collinear BPS rays. The wall-crossing formula therefore becomes



Figure 3: The configuration of BPS rays which participate in the wall-crossing formula (2.44). As we approach the wall, all of the rays shown here become aligned. On the other side of the wall their ordering is reversed.

$$\prod_{n\nearrow} \mathcal{S}^{\mu}_{\gamma_{ij}+n\gamma} \prod_{m=1}^{\infty} \mathcal{K}^{\omega}_{m\gamma} \prod_{n\searrow} \mathcal{S}^{\mu}_{\gamma_{ji}+n\gamma} = \prod_{n\nearrow} \mathcal{S}^{\mu'}_{\gamma_{ji}+n\gamma} \prod_{m=1}^{\infty} \mathcal{K}^{\omega'}_{m\gamma} \prod_{n\searrow} \mathcal{S}^{\mu'}_{\gamma_{ij}+n\gamma} \tag{2.44}$$

Here we assumed γ primitive, but the derivation goes through with minor modifications in the general case. The general solution of (2.44) can be given as follows. Applying (2.44) to X_{γ_k} for $k \neq i, j$ reveals that $\omega'(m\gamma, \gamma_k) = \omega(m\gamma, \gamma_k)$ for $k \neq i, j$. Now choose some $\gamma_i \in \Gamma_i$ and define $\gamma_j := \gamma_{ji} + \gamma_i$. Then applying (2.44) to X_{γ_i} and X_{γ_j} shows that

$$\begin{aligned} \Pi_i' &= \Pi_i + \sigma(\gamma_{ij}, \gamma_{ji}) \Sigma_{ij} \Sigma_{ji} X_\gamma \Pi_j, \\ \Sigma_{ij}' \Pi_i' &= \Sigma_{ij} \Pi_j, \\ \Sigma_{ji}' \Pi_i' &= \Sigma_{ji} \Pi_j, \\ \Pi_j' &= \Pi_j - \sigma(\gamma_{ij}, \gamma_{ji}) \Sigma_{ij}' \Sigma_{ji}' X_\gamma \Pi_i', \end{aligned} \tag{2.45}$$

where we recall that Σ_{ij} was defined in (2.37) (with similar definitions for $\Sigma_{ij}', \Sigma_{ji}, \Sigma_{ji}'$) and we introduced the notation

$$\Pi_i := \prod_{m=1}^{\infty} (1 - X_\gamma^m)^{-\omega(m\gamma,\gamma_i)} \tag{2.46}$$

(with similar definitions for Π_i', Π_j, Π_j'.) Given μ and ω, these equations determine μ' and ω' as follows. The first line can be viewed as an upper triangular set of equations for $\omega'(m\gamma,\gamma_i)$, $m = 1, 2, \ldots$. Once we know that, the next two equations determine $\mu'(\gamma_{ij}+n\gamma)$ and $\mu'(\gamma_{ji}+n\gamma)$. Finally, the fourth equation determines $\omega'(m\gamma,\gamma_j)$. In fact, it is easy to see that it determines

$$\omega'(m\gamma,\gamma_i) + \omega'(m\gamma,\gamma_j) = \omega(m\gamma,\gamma_i) + \omega(m\gamma,\gamma_j). \tag{2.47}$$

If, as is often the case, we are only interested in the differences $\omega(\cdot,\gamma_{ij})$, then the transformation formulae can be written more directly as a transformation reminiscent of modular forms:

$$\begin{aligned} \Pi_{ji}' &= \Delta^{-2}\Pi_{ij}, \\ \Sigma_{ij}' &= \Delta^{-1}\Sigma_{ij}, \\ \Sigma_{ji}' &= \Delta^{-1}\Sigma_{ji}, \end{aligned} \tag{2.48}$$

where $\Delta := \Pi_{ji} + \sigma(\gamma_{ij},\gamma_{ji})\Sigma_{ij}\Sigma_{ji}X_\gamma$.

Two special cases of the above solution are worth noting:

1. If $\sigma(\gamma_{ij},\gamma_{ji})\Sigma_{ij}\Sigma_{ji} = 1$ and $\omega(\gamma,\gamma_i) = \omega(\gamma,\gamma_j) - 1$ and $\omega(m\gamma,\gamma_i) = \omega(m\gamma,\gamma_j) = 0$ for $m > 1$, then we have
$$\mathcal{S}^{\mu}_{\gamma_{ij}}\mathcal{K}^{\omega}_{\gamma}\mathcal{S}^{\mu}_{\gamma_{ji}} = \mathcal{S}^{\mu'}_{\gamma_{ji}}\mathcal{K}^{\omega'}_{\gamma}\mathcal{S}^{\mu'}_{\gamma_{ij}} \tag{2.49}$$
with $\Sigma_{ij}' = \Sigma_{ij}$, $\Sigma_{ji}' = \Sigma_{ji}$ and $\omega'(\gamma,\gamma_j) = \omega(\gamma,\gamma_i)$ and $\omega'(\gamma,\gamma_i) = \omega(\gamma,\gamma_j)$.

2. If $\omega(\gamma,\gamma_i) = \omega(\gamma,\gamma_j) = 0$ and $\omega(m\gamma,\gamma_i) = \omega(m\gamma,\gamma_j) = 0$ for $m > 1$ and also $\sigma(\gamma_{ij},\gamma_{ji})\Sigma_{ij}\Sigma_{ji} = -1$, then we have $\omega'(\gamma,\gamma_i) = -1$, $\omega'(\gamma,\gamma_j) = 1$, $\omega'(m\gamma,\gamma_i) = \omega'(m\gamma,\gamma_j) = 0$ for $m > 1$ and
$$\begin{aligned} \Sigma_{ij}' &= \mu(\gamma_{ij})(1 - X_\gamma)^{-1} \\ \Sigma_{ji}' &= \mu(\gamma_{ji})(1 - X_\gamma)^{-1} \end{aligned} \tag{2.50}$$
corresponding to
$$\mathcal{S}^{\mu}_{\gamma_{ij}}\mathcal{S}^{\mu}_{\gamma_{ji}} = \prod_{\nearrow} \mathcal{S}^{\mu'}_{\gamma_{ji}+n\gamma}\mathcal{K}^{\omega'}_{\gamma}\prod_{\searrow}\mathcal{S}^{\mu'}_{\gamma_{ij}+n\gamma}. \tag{2.51}$$
Some physical realizations of this formula are discussed in §8.2 below.

The cases where 3 or 4 indices i, j, k, l coincide are subsumed by the above examples.

3. A physical interpretation

So far we have written out an abstract wall-crossing formula which we called the 2d-4d WCF. In this section we describe a natural physical setting where the 2d-4d WCF arises, and thus justify its name. The main idea is to study supersymmetric surface defects in $d = 4$, $\mathcal{N} = 2$ field theories.

In §§3.1 and 3.2 we give basic definitions of the relevant surface defects in the UV theory. Then §3.3 discusses in detail the IR description of surface defects. The main upshot is to give a physical realization of the vacuum groupoid $\mathbb{C}[\mathbb{V}]$ and in particular of the charge torsors Γ_i. Then §3.4 narrows the focus to certain classes of surface defects, and illustrates some important subtleties that can arise in trying to define the charge torsors Γ_i in important physical situations. These subtleties include certain "anomalies" which can prevent the surface defects from being physically sensible. In §3.5 we return to the general theory and give definitions of the BPS data μ and ω in our physical context. Finally, in §3.6 we return to the potential anomalies in defining surface defects and illustrate an anomaly cancellation mechanism.

3.1 Surface defects: kinematics

We consider an $\mathcal{N} = 2$ gauge theory in the presence of a surface defect $\mathbb{S}$, satisfying the same conditions as spelled out in [17]. The surface defect is localized at $x^1 = x^2 = 0$ in Minkowski space. In the ultraviolet it preserves a subalgebra of the superconformal $\mathcal{N} = 2$ algebra. Similarly to the case of the line defects studied in [3], the preserved subalgebra is the fixed subalgebra of an involution. In this case we combine reflection in the x^1-x^2 plane, i.e. a rotation by π around the x^3 axis, with a rotation by π around some axis in "R-symmetry space." The bosonic part of the superconformal algebra is $so(2,2) \oplus so(2)_{12} \oplus (u(1)_r \oplus u(1)_R)$, where the first summand corresponds to conformal transformations in the x^0-x^3 plane, the second summand corresponds to rotations around the surface defect, and the third summand is a surviving $u(1) \oplus u(1)$ from the $su(2) \oplus u(1)_R$ R-symmetry with $u(1)_r \subset su(2)$. The $u(1)_R$ subalgebra is in general anomalous. In the conventions of Appendix A of [3], the preserved Poincaré supersymmetries are Q^1_1, Q^2_2, and their respective Hermitian conjugates $\bar{Q}^2_{\dot{1}}$ and $\bar{Q}^1_{\dot{2}}$.

Thus the surface defect preserves $(2,2)$ SUSY on its "worldsheet." The $d = 4$ supermultiplets can be decomposed into multiplets of this $d = 2$ $(2,2)$ superalgebra. Of central importance in what follows is that part of the $d = 4$ vectormultiplet can be used to define a twisted chiral multiplet for this superalgebra. See Appendix A for details.

In the literature it is common to use the term "surface operator," but we believe that "surface defect" is a better name. It is true that a surface defect wrapping a compact Euclidean surface acts as an operator on the Hilbert space of the bulk theory. However, this operator carries less information than the surface defect itself. For example, the operator does not detect 2d degrees of freedom living at the defect which do not interact with the bulk theory.

Finally, as in [17], we assume that in the IR the 4d Abelian gauge fields exhaust all massless degrees of freedom. In particular there are no massless degrees of freedom confined to $\mathbb{S}$. Throughout this paper we let n denote the number of vacua of $\mathbb{S}$, and r the rank of the 4d gauge group.

3.2 Examples of surface defects

How can surface defects of the desired sort be described?

The bulk vectormultiplet scalars restricted to a two-dimensional subspace transform as twisted chiral fields under the $(2,2)$ SUSY algebra (the bulk $u(1)_R$ becomes the axial $u(1)_A$ for the 2d SUSY algebra). So one way to write a Lagrangian describing a surface defect is simply to build a twisted superpotential $\mathcal{W}$ out of these twisted chiral fields, and integrate it over two dimensions. Another way of defining surface defects is to introduce some purely 2d degrees of freedom, with $(2,2)$ supersymmetry and some continuous global "flavor" symmetries. These degrees of freedom can then be coupled to the 4d theory as follows: the restriction of the 4d gauge fields to two dimensions gives 2d gauge fields, and we can use these gauge fields to gauge the flavor symmetries. In this case the 4d vectormultiplet scalars behave as twisted masses from the 2d point of view.

Naively the $\mathcal{W}$ we use would have to be gauge invariant. However, there is an interesting alternative [6]. One can reduce the gauge group G at the defect down to a subgroup H whose center contains at least one $U(1)$ factor (more properly, a "Levi subgroup"). For simplicity here let us consider just the case of a single $U(1)$. One can then write a simple FI-like twisted superpotential, linear in the $U(1)$ part $\Phi^{U(1)}$ of the bulk vectormultiplet scalars:

$$\mathcal{W} = t_{\mathrm{uv}}\, \Phi^{U(1)}. \tag{3.1}$$

The effect of this superpotential can be conveniently expressed in terms of two real couplings $(\alpha,\, \eta)$, the "Gukov-Witten parameters" [6]. They are related to t_{uv} by

$$t_{\mathrm{uv}} = \eta_{\mathrm{uv}} + \tau_{\mathrm{uv}}\alpha_{\mathrm{uv}} \tag{3.2}$$

where τ_{uv} is the complexified gauge coupling in the $U(1)$ we are considering. (If G is a simple group then τ_{uv} is just the coupling for G.) η_{uv} is a 2d theta angle coupled to the magnetic field in the $U(1)$ factor. α_{uv} fixes a boundary condition for the $U(1)$ part of the gauge connection near the defect, of the form

$$A^{U(1)} \sim \alpha_{\mathrm{uv}}\, \mathrm{d}\varphi, \tag{3.3}$$

where φ is the angle in a system of polar coordinates in the plane transverse to the defect.

So far we have implicitly worked with a cutoff field theory and not considered the constraint of UV completeness. Requiring the surface defect to have a good continuum limit turns out to be a significant further restriction. The twisted superpotential $\mathcal{W}$, for example, has to have dimension 1, and hence can only be linear in the 4d vectormultiplet scalars: if the 4d theory is nonabelian, $\mathcal{W}$ cannot include even the simplest gauge invariant operators such as $\mathrm{Tr}\,\Phi^2$. The Gukov-Witten defect just described gets around this problem by breaking the gauge symmetry, and does make good sense in the UV. Another possibility is to take a 2d $(2,2)$ theory which is defined in the UV (say, a gauged linear sigma model) and couple it to the restrictions of the 4d gauge fields.

In the original context of [6], $\mathcal{N}=4$ SYM, both $\eta_{\rm uv}$ and $\alpha_{\rm uv}$ are periodic, but in the current context of surface operators in $\mathcal{N}=2$ theories the parameter space is modified by 2d instanton effects controlled by a 2d instanton factor $e^{2\pi i t_{\rm uv}}$. Since the couplings run, and in particular $\tau_{\rm uv}$ goes to $i\infty$, some renormalization is needed in defining $t_{\rm uv}$.

Finally, we should mention that the coupling of the surface defect to the bulk hypermultiplets might also be important. Unfortunately, it is also poorly understood. When expanded in terms of 2d fields, the bulk hypermultiplets give rise only to chiral multiplets, which cannot directly enter the twisted superpotential couplings which are usually the focus of our computations. Still, the bulk hypermultiplets can affect the definition and properties of the surface defect in the UV in more subtle ways. Basic examples based on brane constructions suggest that some extra discrete structure is hidden in the definition of a "Gukov-Witten defect" in the presence of matter hypermultiplets. We will come back to some of these issues when we specialize to theories and surface defects with a six-dimensional construction, in §7 (see the introduction to §7 and §7.4). The possibility of gauging discrete flavor symmetries will play an important role.

3.3 Low energy (IR) description of the 2d-4d system

Now let us discuss what these combined 2d-4d systems look like in the IR. We begin by recalling the 2d and 4d stories separately and then explain how they can be combined.

3.3.1 Massive 2d theories in the IR

We begin by briefly recalling some of the physics of 2d $\mathcal{N}=(2,2)$ theories with no massless degrees of freedom in the IR, and discrete vacua labeled by $i=1,\dots,n$.

In such a theory one can consider solitons which interpolate between vacuum i and vacuum j (ij-solitons for short). Such a soliton obeys a BPS bound of the form $M \geq |Z_{ij}|$, where Z_{ij} is the central charge, given by

$$Z_{ij} = \mathcal{W}_i - \mathcal{W}_j \tag{3.4}$$

where $\mathcal{W}_i$ and $\mathcal{W}_j$ are the values of the IR superpotential in vacua i and j respectively. In particular, BPS ij-solitons have mass exactly $|Z_{ij}|$.

If the theory includes flavor symmetries, then there is a generalization of the above, which will play an important role in this paper: in addition to the topological charge determined by the pair (i,j) of vacua, particles can carry charge under the flavor symmetries. Letting γ denote an element in the lattice Γ of flavor charges, the total charge of a particle is given by a triple (i,j,γ).

We now introduce a bit of superfluous-looking notation which will be very convenient when we treat coupled 2d-4d systems: we let Γ_{ij} denote the set of all charges $\gamma_{ij}=(i,j,\gamma)$. Of course, Γ_{ij} is trivially isomorphic to Γ, via the map $(i,j,\gamma)\mapsto\gamma$.

Once we have flavor symmetries, the theory can be deformed by "twisted masses" [24], whose IR effect is to change the central charge to

$$Z_{\gamma_{ij}} = \mathcal{W}_i - \mathcal{W}_j + m\cdot\gamma, \tag{3.5}$$

where m is some linear functional on Γ. Twisted masses can be rather important for the physics of a 2d system. For example, in the presence of twisted masses we might have BPS ii-solitons (or more simply "BPS particles in vacuum i"), carrying only the flavor charges. These particles play an important role in the wall-crossing phenomena. This contrasts with the behavior of 4d particles with pure flavor charge, which are invisible to the 4d IR dynamics and wall-crossing.

When we consider 2d-4d coupling, we will find an analogous but more intricate situation, where the lattice Γ includes 4d charges as well as 2d flavor charges.

3.3.2 4d gauge theories in the IR

We now quickly review some standard facts about the IR behavior of 4-dimensional $\mathcal{N} = 2$ gauge theories.

The vacua of the 4d theory are parameterized by the Coulomb branch $\mathcal{B}$, an r-dimensional complex manifold. (Other branches of the moduli space will play no role in our discussion.) For a generic vacuum $u \in \mathcal{B}$, there is an unbroken abelian gauge group $\mathcal{G}_{\text{ab}} \simeq U(1)^r$. There is a "singular locus" in $\mathcal{B}$ where the description in terms of pure $\mathcal{G}_{\text{ab}}$ gauge theory breaks down; let $\mathcal{B}^*$ be its complement. There might also be a continuous group of flavor symmetries; for generic values of the mass parameters of the theory, this flavor group will be of the form $U(1)^{n_f}$.[14]

There is a local system of lattices $\Gamma \to \mathcal{B}^*$, of rank $2r + n_f$. Γ is the "charge lattice" of the theory.[15] It arises as an extension

$$0 \to \Gamma_f \to \Gamma \to \Gamma_g \to 0, \tag{3.6}$$

where Γ_f and Γ_g are respectively the lattices of charges under the continuous flavor and gauge symmetries. Γ_f is fibered trivially over $\mathcal{B}^*$, but Γ_g generally is not: the gauge charges can experience monodromy $\gamma_g \to M(\gamma_g)$, for some $M \in \text{Aut}(\Gamma_g)$, under parallel transport around a nontrivial loop in $\mathcal{B}^*$. Moreover, the sequence (3.6) can be split locally but perhaps not globally. What this means in more physical terms is that the splitting of a charge γ into flavor and gauge charges, $\gamma = \gamma_f \oplus \gamma_g$, is not globally well-defined: monodromy around a loop in $\mathcal{B}^*$ takes $\gamma_f \to \gamma_f + N(\gamma_g)$ for some $N \in \text{Hom}(\Gamma_g, \Gamma_f)$. These phenomena were noted already at the birth of Seiberg-Witten theory [26].

The lattice Γ_g is equipped with the Dirac-Schwinger-Zwanziger pairing of electric and magnetic charges: this is an integral, anti-symmetric, monodromy invariant pairing which we write as $\langle \cdot, \cdot \rangle$. We will assume that Γ_g is self-dual under this pairing.[16] We define an integral antisymmetric pairing $\langle \cdot, \cdot \rangle$ on Γ, whose annihilator is the sublattice Γ_f, by projecting both arguments to Γ_g and using the nondegenerate pairing on Γ_g.

---

[14] In general, the full flavor group could have several disconnected components. We elaborate on that in §3.6.1.

[15] As explained in [3] this should be taken to be the lattice of all IR charges Γ_L where L ranges over a maximal set $\mathcal{L}$ of mutually local line defects in the theory. This lattice was denoted $\Gamma_{\mathcal{L}}$ in [3].

[16] This assumption can be justified in Lagrangian $\mathcal{N} = 2$ theories at weak coupling. See in particular equation (2.13) of [3]. We believe it could be relaxed with some minor adjustments to our analysis.

When writing local formulae it is often useful to choose a symplectic basis e_I, e^I for Γ_g, with $\langle e^I, e_J\rangle = \delta^I{}_J$, $I, J = 1, \dots r$. We then write IR gauge charges $\gamma_g \in \Gamma_g$ as $\gamma_g = p^I e_I + q_I e^I$, with $p^I, q_I \in \mathbb{Z}$.

An essential piece of IR data in the 4d theory is a globally defined section $Z \in \mathrm{Hom}(\Gamma, \mathbb{C})$, known as the central charge function. For each local section γ of Γ, we thus have a function Z_γ on a patch of $\mathcal{B}^*$, obeying

$$Z_{\gamma+\gamma'} = Z_\gamma + Z_{\gamma'}. \tag{3.7}$$

BPS states of charge γ have mass $|Z_\gamma|$.

The Z_{γ_f} for $\gamma_f \in \Gamma_f$ are actually *constant* functions on the whole $\mathcal{B}^*$. They can be organized as $Z_{\gamma_f} = m \cdot \gamma_f$, where m is a (u-independent) complex-valued linear functional on Γ_f, i.e. $m \in \Gamma_f^* \otimes_{\mathbb{Z}} \mathbb{C}$. m is determined by the choice of mass parameters in the theory. If we choose a duality basis and a splitting of (3.6), we may decompose $\gamma = p^I e_I + q_I e^I + \gamma_f$, and correspondingly write

$$Z_\gamma(u) = p^I a_{D,I}(u) + q_I a^I(u) + m \cdot \gamma_f, \tag{3.8}$$

for some local holomorphic functions a^I and $a_{D,I}$.

As u varies in $\mathcal{B}$, $Z(u)$ varies in a Lagrangian subspace of $\Gamma^* \otimes_{\mathbb{Z}} \mathbb{C}$. Once a duality basis has been chosen, we can write a local generating function $\mathcal{F}(a^I)$ for this subspace (also known as a prepotential), i.e. we have the standard relation

$$a_{D,I} = \frac{\partial \mathcal{F}}{\partial a^I}. \tag{3.9}$$

The a^I have a simple physical interpretation, as follows. Having chosen a duality basis over some patch of $\mathcal{B}^*$, we have also fixed a specific Lagrangian description of the effective IR theory in that patch, as a $U(1)^r$ gauge theory. The a^I are the vacuum expectation values of the scalar components of the r $U(1)$ vector multiplets appearing in this Lagrangian. The Lagrangian can be written explicitly as

$$\begin{aligned} S &= \int \mathrm{d}^4\theta \, \mathrm{d}^4 x \, \mathcal{F} \\ &= -\frac{1}{4\pi} \int \left(Y_{IJ} F^I \star F^J + X_{IJ} F^I F^J - Y_{IJ} D^I \star D^J \right) + \cdots \end{aligned} \tag{3.10}$$

where in the first line $\mathcal{F}$ is evaluated on the superfields corresponding to a^I, in the second line we have dropped terms which will not be used in explicit computations below, and the matrix functions X, Y of a^I are the real and imaginary parts of the coupling

$$\tau_{IJ} = \frac{\partial a_{D,I}}{\partial a^J} = X_{IJ} + \mathrm{i} Y_{IJ}. \tag{3.11}$$

The auxiliary fields D^I_{AB} form an $su(2)_R$ triplet. Since we will eventually add a surface defect which breaks $su(2)_R \to u(1)_r$, one component, namely D^I_{12}, is distinguished, and is denoted above simply by D^I.

3.3.3 2d-4d theories in the IR: Local picture, Gukov-Witten parameters, and the torsors Γ_{ij}

Now we discuss the IR structure of the combined 2d-4d system.

A vacuum for the combined system is determined by a pair (u, i) of a point in the Coulomb branch $\mathcal{B}$ of the bulk 4d theory and a vacuum for the surface defect. In a neighborhood of any specific (u, i), it is possible to write an IR Lagrangian for the system in terms of a holomorphic prepotential $\mathcal{F}$ and a holomorphic twisted superpotential $\mathcal{W}_i$, extending (3.10):

$$S = \int \mathrm{d}^4\theta \, \mathrm{d}^4x \, \mathcal{F} + \left(\frac{1}{2} \int \mathrm{d}^2\tilde{\vartheta} \, \mathrm{d}x^3 \, \mathrm{d}x^0 \, \mathcal{W}_i + \text{c.c.} \right). \tag{3.12}$$

The amount of supersymmetry we have here also allows unprotected D-terms integrated over the defect. We will restrict ourselves to considering objects which depend on the twisted F-terms only.[17]

The twisted superpotential $\mathcal{W}_i$ is a function of the associated twisted chiral multiplet Υ, defined in (A.4). It might also depend on the mass parameters of the 4d theory, and possibly extra twisted mass parameters associated to flavor symmetries of the UV surface defect. Indeed, some flavor charges might well be carried only by the 2d degrees of freedom, and be absent in the 4d theory. The corresponding mass parameters are twisted masses.

Notice that $\mathcal{W}_i$ might become singular (and the simple Lagrangian description break down) at new singular loci in $\mathcal{B}^*$, where the 2d physics becomes more intricate. We will from now on redefine $\mathcal{B}^*$ to exclude these loci.

One important place where the twisted superpotential plays a role is in the physics of BPS solitons. So let us consider an ij-soliton. We introduce the symbol γ_{ij} for its total charge, and let Γ_{ij} denote the set of all possible charges γ_{ij}. As we reviewed in §3.3.1, in a pure 2d theory one could decompose any γ_{ij} into the "topological charge" (i, j) plus a pure flavor charge γ. Here we have a similar story: γ_{ij} can be decomposed into the "topological charge" (i, j) plus a charge γ, which now can include 4d gauge charge as well as flavor charge. BPS solitons obey a mass formula of the form $M = |Z|$, with Z written as a sum of two pieces (cf. (3.5)):

$$Z_{\gamma_{ij}} = \mathcal{W}_i - \mathcal{W}_j + Z_\gamma. \tag{3.14}$$

However, in contrast to the pure 2d case, we will discover momentarily that the map from Γ_{ij} to Γ taking $\gamma_{ij} \mapsto \gamma$ is not canonically defined, and neither is the decomposition (3.14).

Let us try to understand the precise relation between Γ_{ij} and Γ. Suppose we bring a bulk line defect, with some charge γ', close to a soliton with charge γ_{ij}. When viewed from a long distance the resulting object looks like a soliton carrying charge $\gamma' + \gamma_{ij}$; so there is an addition operation $\Gamma \times \Gamma_{ij} \to \Gamma_{ij}$ (and similarly in the other order). Moreover, bringing an ij soliton and a jk soliton together shows that there is an addition operation

---

[17] The F-term integration measure can be written a bit more covariantly as the four-form in superspace

$$n_{IJ} \mathrm{d}\theta^I_\alpha \mathrm{d}\theta^J_\beta (C\gamma_{\mu\nu})^{\alpha\beta} \mathrm{d}x^\mu \mathrm{d}x^\nu, \tag{3.13}$$

where I, J are $SU(2)_R$ doublet indices and n_{IJ} is the direction of the $u(1)_V$ subalgebra in $SU(2)_R$.

$\Gamma_{ij} \times \Gamma_{jk} \to \Gamma_{ik}$. Finally, consider "*ii* solitons," i.e. particles in the vacuum i. If we have such a particle we may put it onto a surface defect in vacuum i with its spatial direction compactified. At long distance we can neglect the circumference of this cylinder; it thus looks like a line defect, so the charge it carries must lie in Γ (which by definition is the lattice of all charges of line defects in the 4d theory). So we conclude that Γ_{ii} is naturally isomorphic to Γ, for every i. It follows that each Γ_{ij} is a *torsor* for Γ.[18] The reader should note well that the properties of Γ_{ij} we have just verified are exactly the ones which were used in formulating the 2d-4d WCF in §2.3.

We emphasize that in the above the charge γ might include flavor contributions as well as gauge charges. In particular it might include charges for flavor symmetries which act only on the surface defect $\mathbb{S}$, not on the 4d theory. We always extend the lattices Γ_f and Γ to include such 2d flavor charges as necessary; Z_γ then includes the corresponding contribution from twisted mass couplings on $\mathbb{S}$.

The charges measured by Γ_{ij} are the same kind of charges measured by the bulk charge lattice Γ (they would be obtained by integrating the appropriate currents). From this point of view the difference between Γ_{ij} and Γ is only that the charges in Γ_{ij} need not be properly quantized. In other words, there is an embedding $\Gamma_{ij} \subset \Gamma_{\mathbb{R}}$; so Γ_{ij} looks like a shifted copy of the charge lattice Γ.

Now we should explore some consequences of the Lagrangian couplings (3.12). Very similarly to (3.9), differentiating $\mathcal{W}_i$ once yields

$$\frac{\partial \mathcal{W}_i}{\partial a^I} = t_I =: \eta_I + \tau_{IJ}\alpha^J, \tag{3.15}$$

where η_I, α^I are real functions of the a^I, and we suppress their i-dependence. We claim that these parameters should be viewed as coordinates in a symplectic vector space, so the object

$$\nu := \alpha^I e_I + \eta_I e^I \tag{3.16}$$

is duality invariant. Moreover, we claim that α^I, η_I should be interpreted as IR Gukov-Witten parameters for the surface defect.[19]

In order to establish our claims we introduce the duality invariant gauge field $\mathbb{F} = e_I F^I + e^I G_I$ (where e_I, e^I are a symplectic frame as introduced in §3.3.2). Then the relevant terms in the action (3.12) can be written as

$$\begin{aligned} &-\frac{1}{4\pi}\int \left(Y_{IJ}F^I \star F^J + X_{IJ}F^I F^J - Y_{IJ}D^I \star D^J\right) + \\ &+ \int \mathrm{d}x^0 \mathrm{d}x^3 \left((\eta_I + X_{IJ}\alpha^J)F^I_{03} + \alpha^I Y_{IJ}(F^J_{12} - D^J)\right). \end{aligned} \tag{3.17}$$

The equations of motion following from (3.17) admit a cylindrically symmetric solution $F^I_{12} = 2\pi\delta\alpha^I$ and $F^I_{03} = 0$, where $\delta := \delta(x^1)\delta(x^2)$. This implies that α^I determines the holonomy of the gauge field A^I according to the Gukov-Witten prescription. Moreover, properly dualizing the gauge field in the presence of electric sources gives

---

[18] To see this, just choose any $\gamma_{ji} \in \Gamma_{ji}$: then we get a map $\Gamma_{ij} \to \Gamma_{ii} = \Gamma$ defined by adding γ_{ji}, which is injective and commutes with the Γ actions on both sides.

[19] We emphasize that this is true whether or not the surface defect was defined as a Gukov-Witten defect in the UV. Moreover, even if it were so defined, the UV Gukov-Witten parameters are very different from the ones we see in the IR unless everything is weakly coupled and instanton corrections are suppressed.

$$\begin{aligned} G_{12I} &= Y_{IJ}F^J_{03} - X_{IJ}F^J_{12} + 2\pi\delta(\eta_I + X_{IJ}\alpha^J), \\ G_{03I} &= -Y_{IJ}F^J_{12} - X_{IJ}F^J_{03} + 2\pi\delta Y_{IJ}\alpha^J, \end{aligned} \tag{3.18}$$

and hence the duality invariant gauge field is

$$\mathbb{F}_{12} = 2\pi\delta\nu, \tag{3.19}$$

with all other components $\mathbb{F}_{mn} = 0$. In particular, η_I determines the holonomy of the dual gauge field $A_{D,I}$. Integrating out the auxiliary term and substituting the solution (3.19) one finds that the on-shell action is *precisely zero*, thus showing that the configuration is in fact supersymmetric. Thus, in the IR the surface defect appears to be an infinitely thin solenoid with field configuration (3.19).

3.3.4 Superpotentials and their shifts: the torsors Γ_i

We must stress that the Gukov-Witten parameters used in (3.15) and (3.19) are real valued rather than periodic, that is, ν is a local section of $\Gamma_g \otimes \mathbb{R}$ over $\mathcal{B}^*$. This raises an apparent puzzle since large gauge transformations outside of the solenoid induce shifts

$$\nu \to \nu + \gamma_g \tag{3.20}$$

with $\gamma_g \in \Gamma_g$, and hence our superpotentials are not gauge invariant.

Usually, the shift (3.20) is not considered a puzzle. It is usually said that the physical effects of the solenoid are only measurable through Aharonov-Bohm phases of test particles transported around loops linking the solenoid. Recall that the wavefunction of a test particle with charge γ^{test} which is adiabatically transported around the surface defect picks up a phase $e^{2\pi \mathrm{i}\langle\nu,\gamma_g^{\text{test}}\rangle}$ where γ_g^{test} is the projection of γ^{test} in (3.6).[20] Of course, the Aharonov-Bohm phase determines (α^I, η_I) only up to shifts $(\alpha^I, \eta_I) \to (\alpha^I + m^I, \eta_I + n_I)$ with $m^I, n_I \in \mathbb{Z}$, or equivalently up to (3.20).

In our story, by contrast, we need $\nu \in \Gamma_g \otimes \mathbb{R}$ and by (3.15) it follows that the shifts (3.20) correspond to shifting $\mathcal{W}_i \to \mathcal{W}_i + n_I a^I + m^I a_{D,I}$, i.e.

$$\mathcal{W}_i \to \mathcal{W}_i + Z_{\gamma'} \tag{3.21}$$

where we have now lifted $\gamma'_g \in \Gamma_g$ to some $\gamma' \in \Gamma$, i.e. we have included a further possible constant shift coming from the flavor charges. The resolution of the apparent puzzle is that a specific choice of superpotential is like a specific gauge choice.

How does this ambiguity of $\mathcal{W}_i$ affect the central charge functions as given in (3.14)? For a fixed charge γ_{ij}, $Z_{\gamma_{ij}}$ is a physical quantity which should have no ambiguity. The shift (3.21) leaves $Z_{\gamma_{ij}}$ invariant only if it is compensated by shifting the γ in (3.14) by $\gamma \to \gamma - \gamma'$. This makes sense: by shifting ν we are changing our prescription for the flux through the solenoid, but the flux sourced by the BPS soliton outside the solenoid should be invariant, so we need to shift the (quantized) total flux. This shift explains our earlier remark that the map from Γ_{ij} to Γ which takes $\gamma_{ij} \mapsto \gamma$ is not canonical: it depends on how we choose to fix the ambiguities of $\mathcal{W}_i$ and $\mathcal{W}_j$.

---

[20] For the moment we are assuming that test particles carrying pure flavor charge, $\gamma = \gamma_f$, do not pick up any phase; we will loosen this restriction in §§3.3.6 and 3.6.1 below.

If one is considering the physics of massive BPS particles charged under the abelian gauge fields, the shifts $(\alpha^I, \eta_I) \to (\alpha^I + m^I, \eta_I + n_I)$ change the boundary conditions for the wavefunctions of the charged particles at the solenoid. We discuss the consequences in more detail in §D.

The precise form (3.21) of the ambiguity of $\mathcal{W}_i$ suggests a notation which will turn out to be very useful below. We let Γ_i be a set parameterizing possible choices of $\mathcal{W}_i$. Then for any $\gamma_i \in \Gamma_i$, we denote the corresponding $\mathcal{W}_i$ as Z_{γ_i}. The possibility of a shift (3.21) then just means that for any $\gamma_i \in \Gamma_i$ and $\gamma \in \Gamma$ we can define a new object $\gamma_i + \gamma \in \Gamma_i$, and we have an equation parallel to (3.7),

$$Z_{\gamma_i+\gamma} = Z_{\gamma_i} + Z_{\gamma}. \tag{3.22}$$

These are not the only possible modifications of $\mathcal{W}_i$: after all, we can always shift all the $\mathcal{W}_i$ by the same constant, without changing any of the physical central charges $Z_{\gamma_{ij}}$. For our purposes to follow, though, we will not need to include all possible choices of $\mathcal{W}_i$ in Γ_i. We will only need that Γ_i is closed under all allowed shifts (3.21). We will therefore take Γ_i to be a Γ-torsor.

Earlier we considered another Γ-torsor Γ_{ij} which keeps track of the charges of ij-solitons. It follows from (3.14) that the Γ_i we just introduced is simply related to Γ_{ij}:

$$\Gamma_{ij} = \Gamma_i - \Gamma_j. \tag{3.23}$$

(We defined this notation around (2.18).) This suggests a nice physical interpretation of Γ_i: it is a set of possible charges for *boundaries* of the surface defect $\mathbb{S}$ in vacuum i. At least as far as the IR charges are concerned, then, (3.23) says that a general interface between $\mathbb{S}$ and $\mathbb{S}'$ could be obtained by bringing a boundary of $\mathbb{S}$ close to a boundary of $\mathbb{S}'$. We will explore the properties of boundaries and interfaces further in §4.

3.3.5 2d-4d theories in the IR: Global picture

It is well known that a single physical system can generally be described by many possible prepotentials $\mathcal{F}$, corresponding to different electromagnetic duality frames. We have seen that a similar phenomenon occurs for the superpotentials $\mathcal{W}_i$. In general, the data $(\mathcal{F}, \mathcal{W}_i)$ can be chosen *locally*, but there is no global description of the theory valid everywhere on the Coulomb branch. Rather, we need to glue together different local descriptions. In this section we describe the global structure obtained from this gluing.

Let us first consider $\mathbb{S}$ with a single vacuum ($n = 1$). Call this vacuum i. Recall that globally the charge lattice Γ is a local system over $\mathcal{B}^*$, with monodromies

$$\gamma \to M(\gamma) \tag{3.24}$$

as we go around nontrivial loops in $\mathcal{B}^*$. Similarly, Γ_i is a local system of Γ-torsors over $\mathcal{B}^*$. The monodromy around nontrivial loops in $\mathcal{B}^*$ of Γ_i is similar to that of Γ, but can involve an additional shift by an element of Γ. To write this concretely we could choose an origin in Γ_i and thus identify Γ_i with Γ; then the monodromies would be represented by *affine-linear* maps of the form

$$\gamma_i \to M(\gamma_i) + \gamma. \tag{3.25}$$

We could also ask about the torsor Γ_{ii}; as we have noted, this is canonically isomorphic to Γ, so its monodromies are exactly those of Γ.

For surface defects with $n > 1$ vacua, there is a further complication. Globally, in addition to the kinds of monodromy we discussed above, the vacua i might be permuted among themselves. Locally we have n distinct Γ-torsors Γ_i. The Γ_i do not really exist globally over $\mathcal{B}^*$, but we can continue to use them locally, or we can talk about their direct sum $\Gamma_{\mathbb{S}}$, which does exist globally over $\mathcal{B}^*$. Upon choosing an origin in each Γ_i, we could represent the monodromy of $\Gamma_{\mathbb{S}}$ in the form

$$\gamma_i \to M(\gamma_{P(i)}) + \gamma^{(i)} \tag{3.26}$$

where P is a permutation of the vacua $i = 1, \dots, n$. Similarly, locally we have n^2 distinct Γ-torsors Γ_{ij}, which also do not exist globally over $\mathcal{B}^*$. Their monodromies are of the form

$$\gamma_{ij} \to M(\gamma_{P(i)P(j)}) + \gamma^{(ij)} \tag{3.27}$$

where $\gamma^{(ij)} = \gamma^{(i)} - \gamma^{(j)}$, and in particular $\gamma^{(ii)} = 0$, reflecting the fact that $\Gamma_{ii} \simeq \Gamma$ canonically.

We will refer to the above structure as a "local system of torsors," although this is a slight abuse of terminology. More precisely, we can combine the Γ_i into a single local system of torsors $\Gamma_{\mathbb{S}}$ defined over the ramified cover $\mathcal{B}_{\mathbb{S}} \to \mathcal{B}$ defined in §1, while Γ_{ij} combine into a single local system of torsors $\Gamma_{\mathbb{S}\mathbb{S}}$ over the fiber product $\mathcal{B}_{\mathbb{S}} \times_{\mathcal{B}} \mathcal{B}_{\mathbb{S}} \to \mathcal{B}$.

By now we have introduced almost all of the structure that appeared in the 2d-4d WCF we formulated in §2.3. It only remains to comment on one important — but unfortunately rather obscure — ingredient, the sign $\sigma(a, b)$. Already in the 4d WCF the analogous sign was rather subtle. Based on the halo picture of wall-crossing, we believe that the physical interpretation of the sign has to do with the fact that the electro-magnetic fields sourced by a pair of well-separated bulk particles of charges γ and γ' carry an extra contribution $\langle \gamma, \gamma' \rangle$ to the fermion number.

Locally in moduli space, $\sigma(a, b)$ does not really have more information than the bulk $\sigma(\gamma, \gamma') = (-1)^{\langle \gamma, \gamma' \rangle}$. Indeed, we can always pick an origin γ_i^0 in each Γ_i, thus giving "trivializations" $\Gamma_i \simeq \Gamma$ and $\Gamma_{ij} \simeq \Gamma$. Having done so, we may define $\langle a, b \rangle$ for any a, b using these trivializations and the $\langle \cdot, \cdot \rangle$ on Γ. Then, given any choice of $\sigma(a, b)$, it is possible to make a sign redefinition of the X_a to bring $\sigma(a, b)$ to the form $(-1)^{\langle a, b \rangle}$. To find the correct sign redefinition we just require that $X_{\gamma_i^0 + \gamma} = X_\gamma X_{\gamma_i^0}$ and $X_{\gamma_i^0 - \gamma_j^0 + \gamma} X_{\gamma_j^0} = X_{\gamma_i^0 + \gamma}$. To preserve the wall-crossing identities, such changes of sign should be accompanied by sign changes for $\mu(\gamma_{ij})$, as discussed in §2.3.1. We cannot exclude the possibility that this local redefinition might be impossible globally. We leave such global issues for later investigation in general, but we will give a sign prescription for the A_1 examples in §7.2.

3.3.6 Algebraic integrable systems, mirror symmetry, and surface defects

There is another, rather elegant way of describing the IR data associated to a surface defect in terms of a certain Lagrangian subvariety. This will give a concise summary of the considerations of the previous section.

Recall from §1 that we defined

$$\mathcal{M} = \Gamma_g^* \otimes_{\mathbb{Z}} \mathbb{R}/(2\pi\mathbb{Z}) \to \mathcal{B}. \tag{3.28}$$

As discussed in [1], this is a hyperkähler manifold. Over $\mathcal{B}^*$ it is a torus fibration, and with the special complex structure denoted $\zeta = 0$ in [1] the fibration (3.28) is a holomorphic fibration making it a completely integrable system. Closely allied to this is the system

$$\widetilde{\mathcal{M}} := \Gamma_g \otimes_{\mathbb{Z}} \mathbb{R}/(2\pi\mathbb{Z}) \to \mathcal{B}. \tag{3.29}$$

In fact, $\mathcal{M}$ and $\widetilde{\mathcal{M}}$ are mirror dual manifolds and define mirror conformal field theories upon compactification of the 4d theory to 2d along a two-dimensional torus. In particular $\widetilde{\mathcal{M}}$ is hyperkähler. Moreover, since Γ_g is self-dual, the manifold is self-mirror, i.e. $\mathcal{M}$ and $\widetilde{\mathcal{M}}$ are actually isomorphic. This is very natural since the sigma model with target space $\mathcal{M}$ arises from compactification of the 4d theory on a torus, and exchange of the circles induces mirror symmetry.

Now, the discussion of the previous subsection §3.3.5 implies that, given a surface defect $\mathbb{S}$, the corresponding GW parameters $\tilde{\nu}$ define local sections of $\widetilde{\mathcal{M}}$. Since the vacua form a ramified cover over $\mathcal{B}$ we cannot in general extend $\tilde{\nu}$ to a global section, although it will define a multisection of $\widetilde{\mathcal{M}}$ with a finite number of values in any fiber.

Next, we claim that this multisection is a holomorphic Lagrangian submanifold of $\widetilde{\mathcal{M}}$ in complex structure $\zeta = 0$.

To demonstrate this note that any local section γ_g of Γ_g canonically defines a local $U(1)$-valued function on $\mathcal{M}$. We denote this function as $e^{\mathrm{i}\theta_{\gamma_g}}$. Upon choosing a local framing $\{\gamma_g^j\}$ for Γ_g we obtain a set of coordinate functions $\theta_{\gamma_g^j}$ on the torus fibers of $\mathcal{M}$. The total space of the fibration carries a natural symplectic form

$$\langle \mathrm{d}Z, \mathrm{d}\theta\rangle := C_{ij}\mathrm{d}Z_{\gamma_g^i} \wedge \mathrm{d}\theta_{\gamma_g^j} \tag{3.30}$$

which at $\zeta = 0$ is holomorphic symplectic. Here C_{ij} is inverse to $C^{ij} = \langle \gamma_g^i, \gamma_g^j\rangle$.

Next, given a local section $\tilde{\nu}$ of $\widetilde{\mathcal{M}}$ we can define a corresponding local section of $\mathcal{M}$ whose coordinates at $u \in \mathcal{B}^*$ are given by

$$e^{\mathrm{i}\theta_{\gamma_g}(u)} = e^{2\pi\mathrm{i}\langle\tilde{\nu}(u),\gamma_g\rangle}. \tag{3.31}$$

This subvariety is clearly Lagrangian with respect to $\langle \mathrm{d}Z, \mathrm{d}\theta\rangle$. Moreover it is holomorphic at $\zeta = 0$. We conclude that the same properties hold for the global multisection $\tilde{\nu}$ associated to $\mathbb{S}$. Moreover, by (3.15) its generating function is $\mathcal{W}$ for an appropriate choice of superpotential.

Thus, we have arrived at the most parsimonious description of the IR data of the surface defect $\mathbb{S}$: It is a holomorphic Lagrangian multisection of $\mathcal{M}$ with generating function $\mathcal{W}$.

Remarks

1. In the presence of a continuous flavor symmetry group it is natural to consider a slight twisting of this story, as follows. Since $\Gamma_f^* \otimes_{\mathbb{Z}} \mathbb{R}/(2\pi\mathbb{Z})$ is trivially fibered over $\mathcal{B}$ we can define the continuous flavor symmetry group $\mathcal{F}$ to be the fiber. We can then weakly gauge this global symmetry, and consequently define surface defects with flat connection in the plane transverse to the defect whose holonomy is given by $\theta_f \in \mathcal{F}$. The Aharonov-Bohm phases of test particles transported around such a surface defect involve the flavor charges of the test particles, which can mix with the gauge charges under monodromy around paths in $\mathcal{B}$; in consequence, such a surface defect does not quite correspond to a section of $\widetilde{\mathcal{M}}$. Instead we should consider a torus fibration modeled on the full charge lattice Γ, i.e., $\Gamma^* \otimes_{\mathbb{Z}} \mathbb{R}/(2\pi\mathbb{Z})$. This space sits naturally in an exact sequence of torus fibrations, fiberwise Pontryagin dual to (3.6),

   $$0 \to \Gamma_g^* \otimes_{\mathbb{Z}} \mathbb{R}/(2\pi\mathbb{Z}) \to \Gamma^* \otimes_{\mathbb{Z}} \mathbb{R}/(2\pi\mathbb{Z}) \to \Gamma_f^* \otimes_{\mathbb{Z}} \mathbb{R}/(2\pi\mathbb{Z}) \to 0. \tag{3.32}$$

   The inverse image of θ_f is a twisted version $\mathcal{M}_{\theta_f}$ of $\mathcal{M}$. On $\mathcal{M}_{\theta_f}$ the coordinate transformations of the angular coordinates on the fiber are affine-linear rather than linear. A surface defect with flavor monodromy θ_f gives a multi-section of this twisted integrable system.

2. In §5 we will associate a hyperholomorphic vector bundle $V_{\mathbb{S}} \to \mathcal{M}$ to a surface defect $\mathbb{S}$. Roughly speaking, this vector bundle is the mirror dual to $\tilde{\nu}$. We discuss this further in §5.2.3 and Appendix G.

3.4 Surface defect (twisted) chiral rings: periods as superpotentials

In [17] a close relation was conjectured between surface defects with a non-trivial parameter space C of marginal deformations and Seiberg-Witten-like descriptions of the IR bulk physics. (Here C is a complex variety, possibly of dimension larger than one.) The basic idea is that the marginal deformations of the parameters of the surface defect, $z \to z + \delta z$, are associated to twisted chiral operators $\hat{x}$, and the vevs x of $\hat{x}$ define a 1-form $\lambda = x\, \mathrm{d}z$ on an n-fold cover Σ of C. Let us stress that the symbol z used here is part of the UV definition of the surface defect!

The points (x_i, z) lying over $z \in C$ correspond to the IR vacua of the surface defect with parameter z. Moreover, the charge lattice Γ should be a subquotient of the homology lattice $H_1(\Sigma, \mathbb{Z})$ of closed paths in Σ, and the central charge function Z_γ should be just

$$Z_\gamma = \frac{1}{\pi} \oint_\gamma \lambda. \tag{3.33}$$

So Σ is a (possibly higher-dimensional) analog of the Seiberg-Witten curve, and λ of the Seiberg-Witten differential. The Coulomb branch $\mathcal{B}$ should be a subset of the space of possible complex structure deformations of the covering Σ of the fixed complex manifold C. The lattice Γ should be the quotient by the kernel of the map (3.33) of some subset of the homology lattice closed under all the monodromies around the singular locus of $\mathcal{B}$.

The language developed in this paper is well suited to understand the origin of these statements. Indeed, in this formalism, we have a lot of information available about the effective twisted superpotential; we have [17]

$$x = \pi \partial_z \mathcal{W}. \tag{3.34}$$

We can use this to compute the *difference* $(\Delta\mathcal{W})_{ij}$ between two IR vacua of the surface defect, $i = (x_i, z)$ and $j = (x_j, z)$. According to (3.34), we should just integrate the SW differential $\lambda = x\,\mathrm{d}z$ along an open path γ_{ij} on the SW curve from (x_i, z) to (x_j, z):

$$(\Delta\mathcal{W})_{ij} = \frac{1}{\pi}\int_{\gamma_{ij}} \lambda. \tag{3.35}$$

It is immediately clear that this formula has an ambiguity, namely, different choices of the path γ_{ij} give different $(\Delta\mathcal{W})_{ij}$. The difference is a period of λ along some closed path γ in Σ. Thus we should really use the notation $(\Delta\mathcal{W})_{\gamma_{ij}}$.

We propose to identify the superpotential differences $(\Delta\mathcal{W})_{\gamma_{ij}}$ with the $Z_{\gamma_{ij}}$ controlling BPS solitons between vacua i and j. In particular we provisionally identify the set Γ_{ij} of soliton charges with the set of possible paths γ_{ij} (considered modulo boundaries of 2-chains on Σ). The fact that we can shift an open path by adding a closed path means that Γ_{ij} has a natural action of Γ. This action corresponds to adding a gauge charge to the soliton charge; the annoying ambiguity of $\Delta\mathcal{W}$ is thus identified with the important fact that the solitons carry gauge charges!

More precisely, to be consistent with our general discussion we require that Γ_{ij} should be a torsor for the 4d gauge charge lattice Γ, i.e. any two paths in Γ_{ij} should be related by adding an element of Γ. Since Γ is not quite the same as the full $H_1(\Sigma, \mathbb{Z})$, this implies that Γ_{ij} should not consist of *all* paths from x_i to x_j, but only some subset of paths closed under all the monodromies around the singular locus of $\mathcal{B}$.

Our discussion has a natural extension: we can consider two different surface defects $\mathbb{S}$ and $\mathbb{S}'$, labeled by parameters z and z'. If they are joined (in physical space) by an *interface* then it makes sense to discuss the difference of their superpotentials. This will be given by $(\Delta\mathcal{W})_{\gamma_{ij'}} = \frac{1}{\pi}\int_{\gamma_{ij'}} \lambda$ where $\gamma_{ij'}$ is a path connecting (x_i, z) to $(x_{j'}, z')$. The integral only depends on the relative homology class (with fixed endpoints) defined by $\gamma_{ij'}$ and, as in the above discussion the allowed set of homology classes will form a Γ-torsor $\Gamma_{ij'}$. For further discussion see §4.

Example 3.4. Consider the case where C and Σ are Riemann surfaces. A typical example of a singularity in $\mathcal{B}^*$ arises from the presence of a single light hypermultiplet, whose charge is represented by a vanishing cycle $\tilde{\gamma}$ in Σ. The monodromy of the local system Γ around this singularity is of the Lefschetz type, $\gamma \to \gamma + \langle\tilde{\gamma}, \gamma\rangle\tilde{\gamma}$. Let us now derive the monodromy of Γ_{ij}. The torsor Γ_{ij} is represented by elements of a relative homology group, paths on Σ which end on the preimages of z. These paths have a well defined intersection pairing with elements of another homology group $H_1(\Sigma - \pi^{-1}\{z\}, \mathbb{Z})$. In the region of $\mathcal{B}^*$ sufficiently close to the singularity, we can lift uniquely $\tilde{\gamma}$ to a well defined element $\tilde{\gamma}'$ of this enlarged homology group: $\tilde{\gamma}'$ is the representative of the vanishing cycle which can shrink at the singular locus without crossing the preimages of z. Then the monodromy will be given by the same formula

$$\gamma_{ij} \to \gamma_{ij} + \langle\tilde{\gamma}', \gamma_{ij}\rangle\tilde{\gamma}. \tag{3.36}$$

Another generic type of singularity arises when z hits a branch point of the covering $\Sigma \to C$. Monodromy around a branch point which exchanges two sheets i and j will result in a monodromy exchanging the torsors Γ_{ik} and Γ_{jk}. More precisely, if we denote as γ_{ij}^0 the path in Γ_{ij} which goes from (x_i, z) to (x_j, z) through the branch point (this path is unique in a neighborhood of the singular locus), then we have monodromies $\gamma_{jk} \to \gamma_{ij}^0 + \gamma_{jk}$ and $\gamma_{ik} \to -\gamma_{ij}^0 + \gamma_{ik}$, and of course $\gamma_{ji} \to \gamma_{ij}^0 + \gamma_{ji} + \gamma_{ij}^0$ and $\gamma_{ij} \to -\gamma_{ij}^0 + \gamma_{ij} - \gamma_{ij}^0$.[21] This is the monodromy expected around a locus where a single 2d particle of charge γ_{ij}^0 becomes massless: it is the unique reflection which sends $\gamma_{ij}^0 \to -\gamma_{ij}^0$.

3.4.1 A potential anomaly

We would like to define the superpotential $\mathcal{W}_i$ in a *given* vacuum, rather than the differences $(\Delta\mathcal{W})_{ij}$. In general it might not be possible to find suitable superpotentials $\mathcal{W}_i$ for both mathematical and physical reasons. In this section we indicate some of the possible reasons for such obstructions.

The first potential problem is that it might be impossible to find an appropriate local system of torsors Γ_i over $\mathcal{B}C^*$. (Here $\mathcal{B}C^*$ is $\mathcal{B} \times C$ with certain divisors removed, namely $\mathcal{B}^{\text{sing}} \times C$ and the branch locus of the covering $\Sigma \to C$.) We are trying to solve a cohomology problem by splitting $\Gamma_{ij} = \Gamma_i - \Gamma_j$ where Γ_i are Γ-torsors. (We consider all three terms concretely as subsets of $\Gamma \otimes \mathbb{R}$ rather than just abstract torsors.) In general, there can be an obstruction to this splitting.[22]

Even when torsors Γ_i can be found, we must then also define a suitable extension of Z from $\mathrm{Hom}(\Gamma_{ij}, \mathbb{C})$ to $\mathrm{Hom}(\Gamma_i, \mathbb{C})$. Concretely, we must find Z_{γ_i} so that $Z_{\gamma_{ij}} = Z_{\gamma_i} - Z_{\gamma_j}$, when $\gamma_{ij} = \gamma_i - \gamma_j$, Z_{γ_i} have monodromy shifts only around the singularities in $\mathcal{B}C^*$ (and in no other places), and these shifts are by Z_γ, $\gamma \in \Gamma$.

Even if such candidate superpotentials Z_{γ_i} can be found, we should then worry about possible non-uniqueness. For example, given one solution one could shift $Z_{\gamma_i} \to Z_{\gamma_i} + \mathcal{W}$ where $\mathcal{W}$ has suitable monodromies. (For example, this might be possible if the theory has a surface defect with a single vacuum.) Indeed, in a given physical situation we might wish to put further conditions on the Z_{γ_i}, including properties under the deck transformations permuting the vacua or on the asymptotic behavior in weak coupling limits. We expect that in a physical problem the Z_{γ_i}, if they exist, should be unique, up to trivial redefinitions such as shifts by constants.

We will see explicit examples in §3.4.3 and §7 below where suitable collections of superpotentials Z_{γ_i} in fact do *not* exist.

In order to illustrate what can go wrong, still working in rather general terms, let us consider fibrations Σ with the special property that the sum of λ over the sheets above any point z is zero. (This makes sense for a finite covering and in particular it turns out to be true of the examples in §7, obtained from six-dimensional engineering.) Now, choose some set of local sections $\gamma_{ij'} \in \Gamma_{ij'}$ and $\gamma_{ij} \in \Gamma_{ij}$ such that $\gamma_{ij} + \gamma_{jk} = \gamma_{ik}$ and $\gamma_{ij} + \gamma_{jk'} = \gamma_{ik'}$. Then one could consider the sum

---

[21] Note that it would be wrong to "simplify" $\gamma_{ij}^0 + \gamma_{ji} + \gamma_{ij}^0$ to $\gamma_{ji} + 2\gamma_{ij}^0$. The latter makes no sense!

[22] The precise obstruction lies in $H^2(\mathcal{B}^*, \Gamma)$. To see this we begin with local splittings $\Gamma_{ij} = \Gamma_i - \Gamma_j$. Starting from splittings over each patch of $\mathcal{B}^*$, one can always glue together over the intersections of two patches $\mathcal{U}_{\alpha\beta}$. However at the intersection of three patches $\mathcal{U}_{\alpha\beta\gamma}$ the composition of the three gluing maps is in general an automorphism of a local splitting over the triple overlap. But the automorphisms of the splitting are just given by shifts $\gamma_i \to \gamma_i + \gamma$. This defines a Čech 2-cocycle valued in Γ.

$$\tilde{Z}_i = \frac{1}{n} \sum_{j'=1}^{n} Z_{\gamma_{ij'}}. \tag{3.37}$$

Notice that $\partial_{z'} \tilde{Z}_i = 0$, so $\tilde{Z}_i$ really depends only on z. Furthermore,

$$\tilde{Z}_i - \tilde{Z}_k = \frac{1}{n} \sum_{j'=1}^{n} \left(Z_{\gamma_{ij'}} - Z_{\gamma_{kj'}} \right) = Z_{\gamma_{ik}}. \tag{3.38}$$

It is tempting to use the $\tilde{Z}_i$ to define Z_{γ_i}, as follows. First we embed Γ_{ij} in $\Gamma \otimes \mathbb{R}$. Then, after choosing representatives $\gamma_{ij} \in \Gamma_{ij}$ such that $\gamma_{ij} + \gamma_{jk} = \gamma_{ik}$, we define

$$\Gamma_i = \left(\frac{1}{n} \sum_{j=1}^{n} \gamma_{ij} \right) + \Gamma. \tag{3.39}$$

This satisfies $\Gamma_{ik} = \Gamma_i - \Gamma_k$ as desired. Then we define Z_{γ_i} by linearity.

Unfortunately, there is a problem with this definition. Suppose we change $\gamma_{ij} \to \gamma_{ij} + \tilde{\gamma}^{(ij)}$ where $\tilde{\gamma}^{(ij)} \in \Gamma$. Then Γ_i shifts by

$$\frac{1}{n} \sum_{j} \tilde{\gamma}^{(ij)} \tag{3.40}$$

which is only guaranteed to live in $\frac{1}{n}\Gamma$. Unless a set of γ_{ij} exists for which the monodromies around loops in $\mathcal{B}^*$ only act by shifts in Γ, this naive proposal will run into trouble. As we indicate in §§3.4.3 and 7, there are indeed examples where the proposal fails: Shift happens.

While the above attempt failed to produce a suitable torsor Γ_i, the problem is "only" that there is a shift by a fractional charge in Γ. Locally we could embed Γ_i in $\Gamma \otimes \mathbb{R}$ and then take the smallest subset closed under monodromy to produce a discrete covering space of $\mathcal{BC}^*$. Thus, in some sense, the Γ_i exist, and we can call these Γ_i *twisted torsors*. We will indicate in §3.6 how surface defects with such twisted torsors of charges can be made nonanomalous by gauging finite flavor symmetry groups. This is reassuring since the choice (3.37) is very natural in the A_n theories of class $\mathcal{S}$, considered in §7.

3.4.2 Example: The GW surface defect in pure $SU(2)$ gauge theory

In this section we consider in detail an example of the general type discussed above. Further discussion of this example can be found in §§7 and 8.

The pure $SU(2)$ gauge theory coupled to a GW surface defect was studied in [17]. The infrared theory is controlled by two order parameters: a bulk vev $u = \langle \frac{1}{2} \mathrm{Tr}\, \Phi^2 \rangle$ and the vev of a twisted chiral operator $x = \langle \frac{1}{2\mathrm{i}} \Phi^{U(1)} \rangle$. Classically, in the vacuum we will have $x^2 = -2u$. The quantum-corrected chiral ring relation is conjectured [17] to become

$$x^2 = \Lambda^2 e^t - 2u + \Lambda^2 e^{-t}. \tag{3.41}$$

We caution the reader about some potential notational confusions: We stress that t is a parameter defined in the UV, and should not be confused with the IR GW parameter of (3.15), which will be denoted t_{IR} in this subsection. In certain regimes, explained below, t can be identified with a UV GW parameter normalized such that $t = 2\pi \mathrm{i} t_{\mathrm{uv}}$, when compared with (3.2). Moreover, t in this subsection is denoted z above, but because of its dual interpretation as a 2d instanton amplitude we prefer to use t.

We take C to be a cylinder, identified as the quotient of the complex t-plane by translation by $2\pi\mathrm{i}\mathbb{Z}$. Physically, the cylinder C is the one used in Witten's M-theory construction of the SW model of [43], where $t = x^6 + \mathrm{i}x^{10}$. We will use the fundamental domain $-\pi < \operatorname{Im} t \leq \pi$ to illustrate C, as in Figure 4.

Equation (3.41) should be identified with the defining equation of the Seiberg-Witten curve of the 4d theory, thus identifying Σ in the above discussion with that ramified covering of C. Likewise, λ as described above becomes identified with the Seiberg-Witten differential $\lambda = x\,\mathrm{d}t$. At fixed u there are four branch points of the covering $\lambda \to t$. Two are at $\operatorname{Re} t \to \pm\infty$. There are also branch points where $x = 0$, i.e. for t such that

$$\Lambda^2 \cosh t = u. \tag{3.42}$$

There will only be two distinct branch points in C which we denote $t_\pm(u)$. Representing C by its fundamental domain we take $t_+(u)$ to be the solution with positive real part, when that real part is nonzero. Note that $t_-(u) = -t_+(u)$. For generic u the Seiberg-Witten curve Σ is a twice-punctured torus. The two punctures are the preimages of the branch points at $\operatorname{Re}(t) = \pm\infty$. As a function of u, the two branch points coincide when $u = \pm\Lambda^2$. These are the monopole and dyon points in the u-plane, $\mathcal{B}$, and we define $\mathcal{B}^*$ to be the complex u-plane with $\pm\Lambda^2$ removed.



Figure 4: The Riemann surface C, together with a basis of cycles generating Γ_u, for u in the weak coupling domain. We choose a square root $x \sim \sqrt{-2u}$ on the complement of the cuts. Paths on this sheet are drawn with solid lines while the paths on the sheet with $x \sim -\sqrt{-2u}$ are drawn with dashed lines. Wiggly orange lines denote branch cuts emanating from the branch points $t_\pm$.



Figure 5: The u-plane for $SU(2)$ Seiberg-Witten theory. The paths $\mathcal{P}_{\pm\Lambda^2}$ are based at u in the weak-coupling regime. Define $\mathcal{P}_\infty$ as the path $u \to e^{2\pi \mathrm{i}}u$. It is homotopic to the path given by traversing first $\mathcal{P}_{-\Lambda^2}$ and then $\mathcal{P}_{\Lambda^2}$.

Let us first review the (very well known) structure of the local system $\Gamma \to \mathcal{B}^*$ in this presentation. Recall that, physically, Γ is the set of IR charges $\Gamma_{\mathcal{L}}$ associated to a maximally local set of line defects in the theory. As described in [3] there are three choices for Γ depending on which set of mutually local line defects we choose to incorporate. We take the theory with gauge group $SU(2)$. In this case $\Gamma = H_1(\bar{\Sigma};\mathbb{Z})$, where $\bar{\Sigma}$ is the torus with no punctures. As we have noted there are two singular points $u_\pm = \pm\Lambda^2$ in $\mathcal{B}$ and we can describe Γ by giving the monodromy around the paths $\mathcal{P}_{\pm\Lambda^2}$ shown in Figure 5. It will be convenient to choose a basis of the fiber of Γ over a point u in the weak coupling region $|u| \gg |\Lambda^2|$. Therefore, we choose a system of cuts for the covering $\Sigma \to C$ as in Figure 4. We then choose a cycle γ_e to be given by taking a path in Σ whose image on C is $t(s) = (t_0 + 2\pi \mathrm{i} s) \bmod 2\pi\mathrm{i}\mathbb{Z}$, $0 \le s \le 1$, where t_0 is a point with $|\Lambda^2 \cosh t_0| \ll |u|$. Note that γ_e remains on one of the two sheets of the cover and is not (anti)invariant under the deck transformation of the cover. See Figure 4. However, the image of γ_e in $H_1(\bar{\Sigma};\mathbb{Z})$ *is* anti-invariant. We next choose a cycle γ_m on Σ whose image on C is a path circling around the two branch points, as in Figure 4. Note that $\langle\gamma_e,\gamma_m\rangle = 1$. These cycles generate the lattice Γ at u. Physically the cycle γ_m corresponds to a magnetic monopole and $2\gamma_e = \gamma_W$ is the charge of the W-boson in the weak-coupling region.

The periods of the cycles γ_e, γ_m are:

$$\begin{aligned} a &:= \frac{1}{\pi}\oint_{\gamma_e} \lambda, \\ a_D &:= \frac{1}{\pi}\oint_{\gamma_m} \lambda. \end{aligned} \tag{3.43}$$

Extending by linearity we obtain the central charge function $Z : H_1(\Sigma;\mathbb{Z}) \to \mathbb{Z}$. The kernel of Z coincides with the kernel of the map $H_1(\Sigma;\mathbb{Z}) \to H_1(\bar{\Sigma};\mathbb{Z})$ (this is generated by the small circles around the two punctures of the torus) so Z descends to an element of $\mathrm{Hom}(\Gamma,\mathbb{C})$.

It is well known that the monodromy of the periods a, a_D around the paths $\mathcal{P}_{\pm\Lambda^2}$ in Figure 5 is:

$$\mathcal{P}_{-\Lambda^2}: \qquad \begin{cases} a \to & a - a_D, \\ a_D \to & a_D. \end{cases} \tag{3.44}$$

$$\mathcal{P}_{\Lambda^2}: \qquad \begin{cases} a \to & 3a - a_D, \\ a_D \to & 4a - a_D. \end{cases} \tag{3.45}$$

It will be useful for what follows to have weak coupling expressions for these periods. Since $|\Lambda^2 \cosh t_0| \ll |u|$, we easily estimate

$$Z_{\gamma_e} = \frac{1}{\pi} \oint_{\gamma_e} \lambda \approx 2\mathrm{i}\sqrt{-2u}. \tag{3.46}$$

For the other period note that at weak coupling the two branch points are approximately

$$t_\pm(u) \approx \pm \log \frac{2u}{\Lambda^2} \tag{3.47}$$

and hence

$$Z_{\gamma_m} = -\frac{2}{\pi} \int_{t_-}^{t_+} \sqrt{2\Lambda^2 \cosh t' - 2u}\, \mathrm{d}t' \approx -\frac{4}{\pi}\sqrt{-2u} \log \frac{2u}{\Lambda^2} \tag{3.48}$$

up to instanton corrections. From (3.46) and (3.48) we easily check that the monodromy under $\mathcal{P}_\infty$, given by $u \to e^{2\pi\mathrm{i}}u$ for $|u| \gg |\Lambda^2|$, is $a \to -a$ and $a_D \to 4a - a_D$. One can check that this is indeed the composition of the transformations around $\mathcal{P}_{-\Lambda^2}$ and then $\mathcal{P}_{\Lambda^2}$.

Now let us add a surface defect at a point $t_0 \in C$. For definiteness, we will choose $\mathrm{Re}(t_0)$ large and positive. We will focus on two physically distinct regimes for the physics in the u-plane. The first regime is defined by $|\Lambda^2 \cosh t_0| \ll |u|$ and $|\Lambda^2| \ll |u|$. We call this the *semiclassical GW regime*, because in this regime the 4d theory is weakly coupled and the surface defect is well-described as a Gukov-Witten defect with UV GW parameter t_0. The second regime is defined by $|\Lambda^2| \ll |u| \ll |\Lambda^2 \cosh t_0|$. In this regime the first term in the twisted chiral ring will become important, while the third term will remain subdominant. In this intermediate region, the twisted chiral ring is well approximated by

$$x^2 = \Lambda^2 e^t - 2u \tag{3.49}$$

which can be identified with the twisted chiral ring of a $(2,2)$ $\mathbb{CP}^1$ sigma model, with twisted mass parameter for the $SU(2)$ flavor symmetry $m^2 = 8u$. Indeed the surface defect is more usefully described in this second regime as the $(2,2)$ $\mathbb{CP}^1$ sigma model weakly coupled to the bulk fields. The 2d dynamics become strongly coupled when the first term in the twisted chiral ring relation dominates over the second. Therefore, we will call this second regime the *weakly coupled sigma model regime.* Of course there is also a corresponding weakly coupled sigma model regime when t_0 has large negative real part, since the chiral ring has $t \to -t$ symmetry.

In the semiclassical GW regime it follows from (3.41) that we can write x as a double expansion

$$x = \pm\sqrt{-2u} \cdot \sum_{n \geq 0} \sum_{p=-n}^{\infty} c_{n,p} \left(\frac{\Lambda^4}{u^2}\right)^n \left(\frac{\Lambda^2}{u} e^{t_0}\right)^p, \tag{3.50}$$

Physically, the two expansion factors are interpreted as 4d and 2d instanton effects. The expansion breaks down when t_0 is near a branch point, i.e. near a zero of x. As we will explain in §7 this is due to the existence of light BPS states, modeled on paths running from t_0 to a branch point. Therefore, there will be a singularity in the physics along the divisor (3.42). In particular, if we fix t_0 there is a singularity in $\mathcal{B}$ at $u = u_0$ defined by

$$u_0 = \Lambda^2 \cosh t_0 \tag{3.51}$$

where the branch point t_+ collides with t_0. As in our general discussion above, we will henceforth redefine $\mathcal{B}^*$ to be the complex u-plane with $\pm\Lambda^2$ *and* u_0 removed.



Figure 6: When $u = u_{\mathrm{gw}}$ is in the semiclassical GW regime we can define open paths γ_{12}^{+} and γ_{12}^{-} on Σ, which belong to the fiber of Γ_{12} above u_{gw}.



Figure 7: Paths in the u-plane defining the monodromy action on the fiber of Γ_{12} above a point $u = u_{\mathrm{gw}}$ in the semiclassical GW regime.

We now construct the torsors $\Gamma_{ij} \to \mathcal{BC}^*$ where $\mathcal{BC}^*$ is $\mathcal{B} \times C$ with the divisors $\{(\pm\Lambda^2, t)\}$ and (3.42) removed. Since $\Sigma \to C$ is two-sheeted there will only be Γ_{12} and its negative Γ_{21}. We will fix $t_0 \in C$ with large positive real part as above and just describe the torsor over the u-plane $\mathcal{B}^*$ at fixed t_0. As we just explained physically, there are then three singular points: $u_\pm = \pm\Lambda^2$ and u_0 defined in (3.51). The vacua over t_0, defined by $x_\pm = \pm\sqrt{2(u_0 - u)}$, get permuted if u follows the path $\mathcal{P}_{u_0}$ around u_0 shown in Figure 7, and hence the covering $\mathcal{B}_{\mathbb{S}} \to \mathcal{B}$ is ramified at $u = u_0$ and $u = \infty$.

Now, to construct Γ_{12} explicitly we first consider the fiber above $u = u_{\mathrm{gw}}$ in the semiclassical GW regime, where we can define two natural elements γ_{12}^+ and γ_{12}^- of Γ_{12} which correspond to open paths on Σ whose projection to C goes from t_0 to the respective branch point $t_\pm$ and back as shown in Figure 6. The difference is clearly $\gamma_{12}^+ - \gamma_{12}^- = \gamma_m$. We then have, up to instanton corrections,

$$
\begin{aligned}
Z_{\gamma_{12}^+} &= \frac{2}{\pi}\int_{t_+}^{t_0} x\mathrm{d}t' \approx \frac{2\sqrt{-2u}}{\pi}(t_0 - t_+) = \frac{at_0}{\mathrm{i}\pi} + \frac{1}{2}a_D, \\
Z_{\gamma_{12}^-} &= \frac{2}{\pi}\int_{t_-}^{t_0} x\mathrm{d}t' \approx \frac{2\sqrt{-2u}}{\pi}(t_0 - t_-) = \frac{at_0}{\mathrm{i}\pi} - \frac{1}{2}a_D.
\end{aligned} \tag{3.52}
$$

From these equations it is easy to compute that the monodromy around the path $\mathcal{P}_\infty$ shown in Figure 7 shifts $Z_{\gamma_{12}^\pm} \to \pm 2a - Z_{\gamma_{12}^\pm}$.

Now let us consider the monodromy around the path $\mathcal{P}_0$ in Figure 7. In order to compute the monodromy of $Z_{\gamma_{12}^+}$ it is actually easier to fix $u = u_0$ and displace $t \to t_0 + \delta t$ and let the phase of δt increase from 0 to 2π. Since we are computing monodromy around a single divisor (3.42) in $\mathcal{B} \times C$ the result will be the same as holding $t = t_0$ and letting u run around $\mathcal{P}_0$. When $u = u_0$ and δt is small, we have

$$
x^2 = 2(\Lambda^2\cosh(t_0 + \delta t) - u_0) \approx (2\Lambda^2 \sinh t_0)\delta t \tag{3.53}
$$

and hence $Z_{\gamma_{12}^+} \sim (\delta t)^{3/2}$. It follows that the monodromy around this locus sends $Z_{\gamma_{12}^+} \to -Z_{\gamma_{12}^+}$.

We are now ready to discuss the full local system above the u plane. We have given a basis for the fiber of Γ_{12} over u_{gw} together with the monodromy around $\mathcal{P}_0$ and $\mathcal{P}_\infty$, shown in Figure 7. Around $\mathcal{P}_\infty$ we have $a \to -a$, $a_D \to 4a - a_D$, $Z_{\gamma_{12}^\pm} \to \pm 2a - Z_{\gamma_{12}^\pm}$ and around $\mathcal{P}_{u_0}$ we have the $Z_{\gamma_{12}^+} \to -Z_{\gamma_{12}^+}$ while a, a_D are unchanged. Note that it follows that around the composition $\mathcal{P}_{\Lambda^2}\mathcal{P}_{-\Lambda^2}$ we have

$$
Z_{\gamma_{12}^+} \to 2a + Z_{\gamma_{12}^+}. \tag{3.54}
$$

In order to compute the monodromy around $\mathcal{P}_{\pm\Lambda^2}$ separately we instead move our fiber of Γ_{12} to sit over $u = u_\sigma$ in the sigma-model regime, where $\Lambda^2 \ll |u_\sigma| \ll |u_0|$. (To be precise, we transport along the path $\mathcal{P}_{gw,\sigma}$ shown in Figure 9.) The cuts for the cover $\Sigma \to C$ are now shown in Figure 8. It is useful to consider a third element γ_{12} in the fiber of Γ_{12} over u_σ. It corresponds to the open path on Σ whose projection to C is $t(s) = (t_0 + 2\pi \mathrm{i} s) \bmod 2\pi\mathrm{i}\mathbb{Z}$, $0 \le s \le 1$. The crucial point is that this element has no monodromy in the region $|u_\sigma| \ll |u_0|$, because the branch points always stay away from this path. Thus γ_{12} is invariant under transport around either of $\mathcal{P}_{\pm\Lambda^2}$. (As a simple check, let us show that it is invariant around $\mathcal{P}_{\Lambda^2}\mathcal{P}_{-\Lambda^2}$. To do this first note that $\gamma_{12} - \gamma_{12}^+$ is



Figure 8: When $u = u_\sigma$ is in the semiclassical sigma model regime, we define an open path γ_{12} on Σ, which gives a useful element in the fiber of Γ_{12} above u_σ.



Figure 9: Paths in the u-plane defining the monodromy for Γ_{12}, acting on the fiber over $u = u_\sigma$ in the semiclassical sigma model regime.

simply γ_e; see Figure 8. Next, recall that we have (3.54) around the composition $\mathcal{P}_{\Lambda^2}\mathcal{P}_{-\Lambda^2}$ and $a \to -a$ around this path, so $Z_{\gamma_{12}} = Z_{\gamma_e} + Z_{\gamma_{12}^+} = a + Z_{\gamma_{12}^+}$ remains invariant.) On the other hand, around $\mathcal{P}_{u_0}$ we have $\gamma_{12} \to 2\gamma_e - \gamma_{12}$. This completes our explicit description of the local system Γ_{12}.

We are now ready to construct the local systems Γ_1, Γ_2 of Γ-torsors. Note that if we define

$$Z_{\gamma_1} = -Z_{\gamma_2} = \frac{1}{2} Z_{\gamma_{12}} = \frac{1}{2} Z_{\gamma_{12}^+} + \frac{a}{2} \tag{3.55}$$

then Z_{γ_i} really do shift by elements Z_γ for $\gamma \in \Gamma$ for all closed curves in $\mathcal{B}^*$, thanks to the factor of 2 in the monodromy $\gamma_{12} \to 2\gamma_e - \gamma_{12}$ around $\mathcal{P}_{u_0}$. Thus, in this example, the potential anomaly raised in §3.4.1 does not occur.

Now, with proper superpotentials in hand, we can fill in a gap in the above discussion and explain the phrase "semi-classical GW regime." The reason is that we can use the weak coupling expressions to show that the IR GW parameter is

$$t_{IR} = \frac{\partial Z_{\gamma_1}}{\partial a} \approx \frac{t}{2\pi \mathrm{i}} + \frac{1}{4}\tau + \frac{1}{2}. \tag{3.56}$$

Since the theory is weakly coupled we can identify t itself with a GW parameter.

3.4.3 An example of the anomaly

One class of theories, discussed further in §§7 and 8, are the so-called Argyres-Douglas ("AD") theories of type N. Here $C = \mathbb{C}$ and the chiral ring/SW equation is of the form:

$$\lambda^2 = P_N(z)(\mathrm{d}z)^2. \tag{3.57}$$

The parameter t of §3.4.2 is here restored to $z \in C$, and $P_N(z)$ is a polynomial of degree N. See [2] for background material. Crucial points for our discussion here are that when N is odd there are branch points at the N roots of P_N together with a branch point at $z = \infty$, but when N is even, $z = \infty$ is not a branch point. Related to this, the local system Γ has a one-dimensional flavor lattice when N is even, but no flavor lattice when N is odd.

We claim that for N odd it is possible to construct Γ-torsors Γ_i compatible with the $\mathbb{Z}_2$ deck transformation so that $\Gamma_{12} = \Gamma_1 - \Gamma_2$, but for N even this is not possible. Thus, the potential anomaly of §3.4.1 is realized when N is even. By "compatible" we mean the following. Since the chiral ring $x^2 = P_N(z)$ has a $\mathbb{Z}_2$ symmetry exchanging the sheets it is natural to invoke the physical principle that there should be symmetry under permutation of the two vacua so that $Z_{-\gamma_1}$ should have the same monodromy as Z_{γ_2} in the splitting $Z_{\gamma_{12}} = Z_{\gamma_1} - Z_{\gamma_2}$. But $Z_{\gamma_1} - Z_{-\gamma_1}$ shifts by *even* elements of Γ, and hence so must $Z_{\gamma_{12}}$.

In order to justify this claim let us briefly discuss the case $N = 3$ in a way which will generalize to all odd N. The construction is very closely analogous to the example of §3.4.2 above. In this case we may parameterize

$$P_3(z) = z^3 - 3\Lambda^2 z + u. \tag{3.58}$$

Λ is a non-normalizable parameter and $\mathcal{B}$ is the complex u-plane. We can choose cuts in the z-plane going from each of the three roots of P_3 to $z = \infty$. The local system $\Gamma \to \mathcal{B}^*$ has monodromy around $u = \pm 2\Lambda^2$ (where a pair of roots collide) and is of rank 2.

Now we add a surface defect at a point $z_0 \in C$. We assume that $|z_0|$ is very large. There is then one new singular point in the u-plane, at u_0 such that $z_0^3 - 3\Lambda^2 z_0 + u_0 = 0$. The analog here of the weakly coupled σ-model regime is

$$|z_0^3| \gg |u| \gg |\Lambda^2|. \tag{3.59}$$

We can construct a special element $\gamma_{12} \in \Gamma_{12}|_{u_\sigma}$, where u_σ is in the weakly coupled σ-model regime, by choosing a path in C beginning and ending at z_0 and circling around all three branch points. Since it has crossed an odd number of cuts it lifts to an open path in Σ joining the vacuum λ_0 to $-\lambda_0$. As in the $SU(2)$ example, this element has no monodromy around all the paths in $\mathcal{B}_*$ based at u_σ which wind around zeros of the discriminant of P_3 (namely $u = \pm 2\Lambda^2$). It *does* have monodromy around the path $\mathcal{P}_{u_0}$ based at u_σ which winds once around u_0. To compute this monodromy, note that as $u_\sigma \to u_0$ (thus leaving the σ-model regime) some branch point, say $z_*(u_\sigma)$, approaches z_0. We can then define an open path $\gamma_{12}^* \in \Gamma_{12}|_{u_\sigma}$ (the analog of γ_{12}^+ above) whose image in C begins at z_0, runs to z_* passing through a cut and comes back to z_0. Note that $Z_{\gamma_{12}} = Z_{\gamma_{12}^*} + Z_\gamma$ for some $\gamma \in \Gamma$. On the other hand, by an argument similar to that of the $SU(2)$ example, $Z_{\gamma_{12}^*} \to -Z_{\gamma_{12}^*}$ around $\mathcal{P}_{u_0}$ and hence the only nontrivial monodromy of $Z_{\gamma_{12}}$ in $\mathcal{B}^*$ is around $\mathcal{P}_{u_0}$ and given by $Z_{\gamma_{12}} \to 2Z_\gamma - Z_{\gamma_{12}}$. We can therefore define $Z_{\gamma_1} = -Z_{\gamma_2} = \frac{1}{2}Z_{\gamma_{12}}$, and these periods will have monodromy shifts by proper periods associated to Γ.

The above construction generalizes to AD theories with odd N. On the other hand, when N is even the situation is very different. Consider, for example, the first nontrivial case, $N = 4$, with

$$P_4(z) = z^4 + 4\Lambda^2 z^2 + 2mz + u. \tag{3.60}$$

Here Λ, m are fixed nonnormalizable parameters and $\mathcal{B}$ is again the complex u-plane. There are four branch points in C given by the roots of $P_4(z)$. The local system Γ has rank 3 with a one-dimensional flavor sublattice and $\Gamma \to \mathcal{B}^*$ has monodromy around the three roots of the disciminant of P_4. When we add a surface defect at $z_0 \in C$ there is a fourth singularity in the u-plane at u_0 such that $P_4(z_0) = 0$. Now, if $u = u_\sigma$ is in the weakly coupled sigma model regime

$$|z_0^4| \gg |u| \gg |\Lambda^2|, |m| \tag{3.61}$$

we *cannot* define the analog of the cycle $\gamma_{12} \in \Gamma_{12}|_{u_\sigma}$ since it will cross an even number of cuts and hence will not connect the two sheets above z_0. If, instead, we choose any other path $\tilde{\gamma}_{12} \in \Gamma_{12}|_{u_\sigma}$ surrounding an odd number of branch points, then that path will pass through a cut joining two branch points. Therefore, there will be a singularity in $\mathcal{B}$ where those two branch points collide. The monodromy of $\tilde{\gamma}_{12}$ around such a point will be $\tilde{\gamma}_{12} \to \tilde{\gamma}_{12} + \gamma_v$ where γ_v is the vanishing cycle associated with the colliding branch points. Therefore, if we attempt to introduce $Z_{\gamma_1} = -Z_{\gamma_2} = \frac{1}{2}Z_{\tilde{\gamma}_{12}}$ then γ_1, γ_2 will have half-integral monodromy $\gamma_1 \to \gamma_1 + \frac{1}{2}\gamma_v$. Thus our attempt has failed. Now, suppose there were some splitting $Z_{\gamma_{12}} = Z_{\gamma_1} - Z_{\gamma_2}$ where $Z_{\gamma_1}, Z_{\gamma_2}$ have proper integral monodromy. Then $-Z_{\gamma_1} = Z_{-\gamma_1}$ would have proper monodromy, and $Z_{\gamma_1} - Z_{-\gamma_1}$ would have monodromy of the type we have just proven cannot exist. Therefore, for $N = 4$ AD theories there is no splitting $\Gamma_{12} = \Gamma_1 - \Gamma_2$ which respects the symmetry between the sheets. The argument we have just given extends to all even values of N.

To use the language of §3.4.1 above, when N is even we can construct Γ_i as twisted torsors. The three examples we have just discussed are examples drawn from a class of theories described in §7 known as A_1 theories. As we discuss in §7 the A_1 theories with regular singular points all exhibit the anomaly we have discussed. Theories with irregular singular points are limits of those with regular singular points. In such limits particles can decouple and the nature of flavor symmetries can change. It can be shown that theories with odd irregular singular points (such as AD N odd and $SU(2)$, $N_f = 0$) have a natural candidate for the analog of the cycle γ_{12}^+ used above and do not exhibit the anomaly. Other theories with only even irregular singular points, such as AD theories with N even still exhibit the anomaly.

3.5 BPS degeneracies and wall-crossing

Now that we have given a physical description of the lattices Γ, torsors Γ_i, Γ_{ij} and central charges Z which appear in the wall-crossing formula, we can move to the main actors, Ω, μ and ω.

First, recall that in the 4d wall-crossing formula the integers $\Omega(\gamma)$ are interpreted as degeneracies of 4d BPS particles [1]. This is their interpretation in the 2d-4d wall-crossing formula as well.

The integers $\mu(\gamma_{ij})$ in the 2d-4d wall-crossing formula are a natural generalization of the index μ_{ij} in the 2d case [20]. $\mu(\gamma_{ij})$ is the degeneracy of 2d BPS solitons between vacuum i and vacuum j of the surface defect, carrying total charge γ_{ij}. Using the notation of Appendix A it can be written:

$$\mu(\gamma_{ij}) := \mathrm{Tr}_{\tilde{\mathcal{H}}^{\mathrm{BPS}}_{\gamma_{ij}}} e^{\mathrm{i}\pi F} e^{-\beta H}. \tag{3.62}$$

Here $\tilde{\mathcal{H}}^{\mathrm{BPS}}_{\gamma_{ij}}$ is the space of BPS states with center-of-mass degrees of freedom removed, while F is some $u(1)_V$ charge. An important point is that there is a *choice* of $u(1)_V$ generator F. One natural choice is $F = so(12)$, but there is some ambiguity in picking the overall fermion number when removing the center of mass degrees of freedom. We will take the $\mu(\gamma_{ij})$ to be integral although $e^{\mathrm{i}\pi F}$ acting on the full Hilbert space $\mathcal{H}_{\gamma_{ij}}$ need not be ± 1. It is standard in the 2d literature to choose F so that $\mu(\gamma_{ij}) = -\mu(-\gamma_{ij})$. The ambiguity in F will be relevant in §5.

The affine-linear functions $\omega(\gamma, \cdot)$ require more discussion. Roughly we want to interpret $\omega(\gamma, \gamma_i)$ as the number of 2d BPS particles in vacuum i, carrying gauge charge γ. If this were literally the case, then $\omega(\gamma, \gamma_i)$ would actually depend only on i, not on the choice of a particular γ_i. The reason for the dependence on γ_i is a rather subtle difficulty. In addition to 1-particle states with charge γ localized near the surface defect, the theory also contains 1-particle "4d" states with charge γ. When decomposed in terms of the unbroken 2d Poincare symmetry these states give a continuum of representations: The momentum in the transverse space to the surface defect is an *internal quantum number* and hence the representation of the little group has no gap from the BPS bound. The representation at the bottom of this continuum is the same one where a genuine 2d particle would be found. It is therefore difficult to disentangle the 2d and 4d spectra.

To get a bit more insight, let us focus on a region of parameter space containing a point where a specific $Z_{\gamma^e} \to 0$. At this point the Lagrangian we have been discussing until now — which includes only the abelian gauge fields and their superpartners — breaks down. One gets a better description near this point by including fields describing the light BPS particles with charge γ^e; the singularity of the original theory would then be reproduced by integrating those fields out. What kind of new fields do we need to add? Light 4d particles should be represented by hypermultiplets; light 2d particles should be represented by 2d chiral multiplets.

The effect of integrating out the light hypermultiplets is very well known: A simple 1-loop computation leads to a correction to the prepotential which creates logarithmic divergences in the central charges:

$$Z_\gamma = \frac{\mathrm{i}}{2\pi}\Omega(\gamma^e)\langle\gamma,\gamma^e\rangle Z_{\gamma^e}\log(Z_{\gamma^e}) + \mathrm{Regular} \tag{3.63}$$

This logarithmic behavior implies the well-known Picard-Lefschetz-like monodromy: as Z_{γ^e} goes around 0 we have $\gamma \to \gamma + \Omega(\gamma^e)\langle\gamma,\gamma^e\rangle\gamma^e$.

It turns out that the effect of integrating out a light 2d particle leads to a correction very similar to (3.63). Indeed, we can invoke the computation of Witten in the two-dimensional gauged linear sigma model (see [30], eq. (3.9) et. seq. and also [31]) to conclude that integrating out n two-dimensional chiral multiplets of four-dimensional charge γ^e induces a singular term in the two-dimensional *superpotential*:

$$\mathcal{W} = \frac{\mathrm{i}}{2\pi}nZ_{\gamma^e}\log Z_{\gamma^e} + \mathrm{Regular}. \tag{3.64}$$

For later reference (in §6.2 below) it will be useful to recall briefly the argument for (3.64). The twisted chiral superfield is (A.4) $\Upsilon = a + \cdots + \vartheta^+\vartheta^-(F_{03} + \mathrm{i}(D_{12} - F_{12}))$, where $Z_{\gamma_e} = qa$ vanishes at $a = 0$ and the n chiral multiplets X_i, $i = 1, \ldots, n$ have $U(1)$ charge q and hence have Lagrangian

$$\int \mathrm{d}x^0\mathrm{d}x^3 \int \mathrm{d}^4\vartheta \bar{X}_i e^{qV} X_i = -\int |\mathcal{D}X_i|^2 + (m^2 + q(D_{12} - F_{12}))|X_i|^2 + \cdots \tag{3.65}$$

where $m^2 = q^2|a|^2$ and we have only written the terms relevant to our computation. Integrating out the chiral superfields induces a one-loop determinant

$$\exp\left(-n\log\det[-\mathcal{D}^*\mathcal{D} + m^2 + q(D_{12} - F_{12})]\right)$$

which when expanded in the auxiliary field gives a coupling

$$\int \mathrm{d}x^0\mathrm{d}x^3(D_{12} - F_{12})nq\int\frac{\mathrm{d}^2k}{(2\pi)^2}\frac{1}{k^2 + m^2} + \cdots \tag{3.66}$$

Supersymmetry now determines the rest. Comparison with the last term in (3.17) reveals that we have induced a Gukov-Witten parameter

$$(\mathrm{Im}\tau)\alpha = -nq\int\frac{d^2k}{(2\pi)^2}\frac{1}{k^2 + q^2|a|^2} = \frac{nq}{4\pi}\log\frac{|a|^2}{|\Lambda|^2} \tag{3.67}$$

where Λ is a UV cutoff, necessary in this IR free gauge theory. On the other hand, a simple computation shows that such an α corresponds to

$$\mathcal{W} = nq\frac{\mathrm{i}}{2\pi}\left(a\log\frac{a}{\Lambda} - a\right) \tag{3.68}$$

thus establishing (3.64).

The above one-loop effect leads to the monodromy of the superpotential discussed in equation (3.21), and hence, as explained there, if we wish to extend the period function and define $\mathcal{W} = Z_{\gamma_i^0}$ then γ_i^0 must shift by $\gamma_i^0 \to \gamma_i^0 + n\gamma^e$ under one turn around $a = 0$. In other words, the presence of light 2d particles leads to affine-linear monodromies of the torsor Γ_i. In particular, we cannot hope to find a single γ_i^0 which makes sense globally: we will always have to make do with local descriptions, related to one another by shifts.

Moreover, more general paths in moduli space force us to consider more general monodromies, $\gamma_i^0 \to \gamma_i = \gamma_i^0 + \gamma$ for more general charges γ. In this case we have an effective superpotential $Z_{\gamma_i} = Z_{\gamma_i^0} + Z_\gamma$ which, by (3.63), is just:

$$Z_{\gamma_i} = \frac{\mathrm{i}}{2\pi}(n + \Omega(\gamma^e)\langle \gamma_i - \gamma_i^0, \gamma^e \rangle) Z_{\gamma^e} \log Z_{\gamma^e} + \text{Regular}. \tag{3.69}$$

Now, there is clearly something odd about this formula since it involves a the choice of a specific $\gamma_i^0 \in \Gamma_i$, which we used in writing our effective Lagrangian. What would have happened if we had made a different choice, say replacing γ_i^0 by $\gamma_i^0 + \tilde{\gamma}$? Perhaps surprisingly, from (3.69) we see that this change must be compensated by a change of n to $n + \Omega(\gamma^e)\langle \gamma^e, \tilde{\gamma} \rangle$ — so a rather innocuous-looking shift of the effective superpotential also requires a change in the number of 2d chiral multiplets! Thus, the number of chiral multiplets depends on the flux and we should write $n(\gamma_i^0)$. The most meaningful and invariant quantity we can extract from our considerations is the coefficient $n(\gamma_i^0) + \Omega(\gamma^e)\langle \gamma_i - \gamma_i^0, \gamma^e \rangle$. which appeared in (3.69), giving the physical effect of integrating out both 2d and 4d particles of charge γ^e. We propose to identify this with $\omega(\gamma^e, \gamma_i)$. In other words, if u is near a locus in $\mathcal{B}$ where some $Z_\gamma \to 0$, we define $\omega(\gamma, \gamma_i; u)$ in terms of the contribution from particles of charge γ to Z_{γ_i},

$$Z_{\gamma_i} = \frac{\mathrm{i}}{2\pi}\omega(\gamma, \gamma_i; u) Z_\gamma \log Z_\gamma + \text{Regular}. \tag{3.70}$$

Then the values of $\omega(\gamma, \gamma_i; u)$ at other vacua will be determined by wall-crossing. Note that $\omega(-\gamma, \gamma_i; u) = -\omega(\gamma, \gamma_i; u)$.

Example: Let us return to Example 3.4 above. We discussed there the example of a singularity in $\mathcal{B}^*$ which arises from the presence of a single light hypermultiplet whose charge is represented by a vanishing cycle $\tilde{\gamma}$ in Σ. From the monodromy of Γ_{ij}, equation (3.36) we recognize immediately $\omega(\tilde{\gamma}, \gamma_{ij}) = \langle \tilde{\gamma}', \gamma_{ij} \rangle$ in the neighbourhood of the singular locus. Similarly, a lot of simple wall-crossing examples can be derived from these statements, by simply considering a locus where z comes close to two branch points. See §7 for several examples along these lines.

Remark: The reader might well be distressed by the absence of a clear definition of $\omega(\gamma, \gamma_i)$. Fortunately, the only quantities that really enter the 2d-4d wall-crossing formula are the differences $\omega(\gamma, \gamma_{ij'})$. These do have a clear definition as a trace in a Hilbert space, which will be detailed in the next section 4. These will also be the quantities which we will have direct control on in our examples.

3.6 Cancellation of surface defect global anomalies by gauging flavor symmetries

In §3.4 we found that surface defects in 2d-4d systems potentially have global anomalies, and in §§3.4.1 and 3.4.2 we mentioned that there are simple examples which illustrate such anomalies. In this section we demonstrate in somewhat general terms how such anomalies can sometimes be cancelled by gauging an appropriate subgroup of the flavor group. (Such a subgroup might or might not exist in any given theory.)

First, let us put the problem we are trying to solve in the proper context:

We are used to thinking about the spectrum of zero-dimensional defects, or local operators, as an integral part of what a theory is, maybe as part of the very definition of the theory. We also think of the spectrum of one-dimensional defects, or line defects, this way, although this often involves a small refinement of our notion of "theory." For example, in four dimensions, there is typically a discrete choice of which class of line defects to allow: in, say, a gauge theory with Lie algebra $su(N)$, the fundamental Wilson loop and fundamental 't Hooft loop cannot appear simultaneously in correlation functions, because they would not be mutually local. The choice of which line defects to allow is part of the proper definition of the theory and involves, for example, a choice of the actual compact gauge group. For example, no theory with gauge algebra $su(N)$ can contain both the fundamental Wilson loop and the fundamental 't Hooft loop because they are mutually nonlocal. The fundamental Wilson loop is allowed if the gauge group is $SU(N)$ while the fundamental 't Hooft loop is allowed if the gauge group is $PSU(N)$. By contrast, we think of a surface defect not as a part of the theory, but rather as a modification of the theory. The spectrum of possible surface defects might be as intricate as the set of all possible 2d theories, or even more so.

In four dimensions, considering surface defects together with local operators might lead to problems with locality: transporting a local operator around a surface defect might induce some unwanted monodromy phases. As we usually consider the set of local operators as a given, we would normally just not allow such surface defects. In §7 we will encounter situations, though, where these surface defects (namely, the canonical surface defects $\mathbb{S}_z$) are too important simply to be thrown away, so it is the local operators which have to give. We will therefore change the theory to accomodate these surface defects. The idea is as follows. The effect of the monodromy around a surface defect must be some symmetry of the bulk theory. We will assume that the symmetry transformations associated to the surface defects we wish to retain lies in a finite abelian group $\mathcal{D}$ which is a subgroup of the group of flavor symmetries. This is indeed what happens to the surface defects $\mathbb{S}_z$ studied in §§7, 8 and 9. As with any finite global symmetry group, we can gauge it. In doing so we throw away the bothersome local operators, which are no longer gauge invariant. At the same time we introduce new surface defects corresponding to nontrivial $\mathcal{D}$-bundles in the space transverse to the surface defect.[23]

Of course, we do not want to throw away the vanilla BPS states. When the charge torsors Γ_i are twisted and have shifts by fractional elements of Γ, the AB phases of test particles of charges $\gamma_g^{\rm test}$ will not be well-defined. We will also see that by gauging a suitable finite flavor subgroup this anomaly can in principle be cancelled.

In general the new surface defects are labeled by conjugacy classes of the gauged finite flavor symmetry group, $\mathcal{D}$, and since we assume it is abelian they will be labeled by elements of that group. In what follows, one consequence of this gauging will be particularly important: surface defects which carry nontrivial flavor monodromy cannot have boundaries. They can, however, have interfaces to other surface defects carrying the same flavor monodromy. Throughout this paper, we will see that gauging the flavor symmetry has subtle and entertaining effects on the IR physics of the surface defect.

---

[23] In a recent paper the effects of gauging finite global symmetry groups in supergravity theories was studied [46, 47]. Some of the considerations of that paper are similar to ours. We thank N. Seiberg for discussions which considerably influenced our thinking on this topic.

3.6.1 Flavor holonomies

In this section we assume that there is a finite abelian flavor group $\mathcal{D}$, trivially fibered over $\mathcal{B}^*$, so that the Pontryagin dual group of characters $\hat{\mathcal{D}}$ fits in an exact sequence

$$0 \to \hat{\mathcal{D}} \to \Gamma_g^{\text{ext}} \to \Gamma_g \to 0. \tag{3.71}$$

As usual, this extension is trivial locally but not globally.

A useful example to keep in mind is given by choosing a finite subgroup of the fiber of $\Gamma_f^* \otimes \mathbb{R}/(2\pi\mathbb{Z})$. Then there is a projection map from Γ_f to $\hat{\mathcal{D}}$. Applying this map to (3.6) we find that the action of the discrete flavor group on charged particles is captured by an extension of local systems of abelian groups (3.71).

What does the exact sequence (3.71) mean? A gauge-invariant state (one whose charge projects to zero in Γ_g) could carry some element of $\hat{\mathcal{D}}$. There is an ambiguity when we try to lift this action to non-gauge-invariant states such as BPS particles: in other words, the discrete flavor charge of a BPS particle is ambiguous. To fix the flavor charges we have to choose a splitting of (3.71). (This is very similar to the ambiguity we have in assigning the usual flavor charges to BPS particles, i.e. in splitting the sequence (3.6).) Any two different splittings differ by an element of $\mathrm{Hom}(\Gamma_g, \hat{\mathcal{D}})$. If we gauge $\mathcal{D}$, then Γ_g^{ext} will be the new extended group of all gauge charges.

The fiberwise Pontryagin dual of (3.71) is an extension of the algebraic integrable system $\mathcal{M}$:

$$0 \to \mathcal{M} \to \mathcal{M}^{\text{ext}} \to \mathcal{D} \to 0. \tag{3.72}$$

The fiber of $\mathcal{M}^{\text{ext}}$ has one connected component $\mathcal{M}_d$ for each element $d \in \mathcal{D}$; the identity component $\mathcal{M}_1 = \mathcal{M}$, while the other $\mathcal{M}_d$ are twisted versions of $\mathcal{M}$.

Now we are ready to consider the surface defects which become available after gauging $\mathcal{D}$. These surface defects carry a flavor monodromy $d \in \mathcal{D}$. The Aharonov-Bohm phases of test particles around such a surface defect give a section of $\mathcal{M}^{\text{ext}}$, or more precisely, of its connected component $\mathcal{M}_d$ along the lines of (3.31). So globally the IR data of the surface defect is determined by a complex Lagrangian multi-section of $\mathcal{M}_d$.

We can describe the monodromy of $\mathcal{M}_d$ more explicitly as follows. Pick a splitting of (3.71) into discrete flavor charges and gauge charges, i.e. decompose charges γ_g^{ext} into $\xi \oplus \gamma_g$, where $\xi \in \hat{\mathcal{D}}$ and $\gamma_g \in \Gamma_g$. Under monodromy around a loop on the Coulomb branch $\mathcal{B}^*$, these charges are transformed by

$$\gamma_g \to M(\gamma_g), \tag{3.73}$$

$$\xi \to \xi + \delta\xi(\gamma_g) \tag{3.74}$$

where $\delta\xi$ is some homomorphism $\Gamma_g \to \hat{\mathcal{D}}$. Notice that we are using an additive notation for the characters ξ and $\delta\xi$. The multiplicative characters are $\exp 2\pi i \xi$, etc.

Then, we can attempt to define functions θ_{γ_g} on $\mathcal{M}_d$ analogous to those on $\mathcal{M}$: Choosing a splitting of (3.71) and a corresponding local trivialization of $\mathcal{M}^{\rm ext}$, $\xi \oplus \gamma_g$ defines a $U(1)$-valued function $\xi(d)e^{{\rm i}\theta_{\gamma_g}}$ on $\mathcal{M}^{\rm ext}$. On the other hand, monodromy takes $(0 \oplus \gamma_g)$ to $\delta\xi(\gamma_g) \oplus M(\gamma_g)$ and hence $\delta\xi$ deforms the monodromy of the fiber coordinates θ_{γ_g} on $\mathcal{M}_d$ from linear to affine-linear:

$$\theta_{\gamma_g} \to \theta_{M(\gamma_g)} + 2\pi[\delta\xi(\gamma_g)](d). \tag{3.75}$$

Since $[\delta\xi(\gamma_g)](d)$ is linear in both γ_g and d and since Γ_g is self-dual with respect to $\langle\cdot,\cdot\rangle$, we can define a useful quantity $\delta\xi(d) \in \Gamma_g \otimes \mathbb{R}/\mathbb{Z}$ by

$$[\delta\xi(\gamma_g)](d) = \exp[2\pi{\rm i}\langle\delta\xi(d), \gamma_g\rangle]. \tag{3.76}$$

Now let us explore the consequences for the Gukov-Witten parameters ν associated to a surface defect with flavor monodromy. If we define a multisection by (3.31) then, after monodromy around a loop in $\mathcal{B}^*$ we will find

$$\exp[i\theta_{M(\gamma_g)} + 2\pi{\rm i}\langle\delta\xi(d), \gamma_g\rangle] = \exp[2\pi{\rm i}\langle\nu', \gamma_g\rangle]. \tag{3.77}$$

It follows that, if we choose some lift $\widehat{\delta\xi(d)} \in \Gamma_g \otimes \mathbb{R}$ then the monodromy of the Gukov-Witten parameters is of the form

$$\nu' = M^{tr} \cdot \nu + \widehat{\delta\xi(d)} + \gamma \tag{3.78}$$

for some $\gamma \in \Gamma_g$. Therefore, by (3.15) the superpotential has a shift $\mathcal{W} \to \mathcal{W} + Z_{\gamma'}$ (where we lift γ' from $\Gamma_g \otimes \mathbb{R}$ to $\Gamma \otimes \mathbb{R}$).

Now, we should stress that the shift by $\widehat{\delta\xi(d)}$ is *not* by an element of Γ_g but rather by a fractional element of Γ_g. But such shifts were precisely the sort we found in potential anomalies in defining the torsors Γ_i in §3.4.1! This will be the key to the anomaly cancellation mechanism described in §3.6.2 below.

It is useful to give a concrete example here. Consider a typical example of a singularity in $\mathcal{B}^*$, which arises from the presence of a single light hypermultiplet, whose charge is $\tilde{\gamma}$. The monodromy of the local system Γ is of the Lefschetz type, $\gamma \to \gamma + \langle\tilde{\gamma}, \gamma\rangle\tilde{\gamma}$. This is easily generalized to the case with discrete flavor symmetries. The flavor monodromy will be $\xi \to \xi + \langle\tilde{\gamma}_g, \gamma_g\rangle\tilde{\xi}$ if the discrete charge of the hypermultiplet is $\tilde{\xi}$. But then $\delta\xi(\gamma_g) = \langle\tilde{\gamma}_g, \gamma_g\rangle\tilde{\xi}$ and thus $\delta\xi(d) = \tilde{\xi}(d)\tilde{\gamma}_g$. In general $\tilde{\xi}(d)$ will be a nontrivial element of $\mathbb{R}/\mathbb{Z}$.

3.6.2 Anomaly cancellation

In the previous section we have learned that by gauging a finite abelian flavor symmetry we can induce shifts of the GW parameters analogous to those forced on us by the cohomological obstruction discussed in §3.4. Therefore, under suitable conditions, gauging a finite flavor symmetry group can cancel the global anomalies obstructing the existence of a surface defect associated with a twisted torsor.

For example, consider the situation discussed in equations (3.37) to (3.40). The shift (3.40) means that AB phases are ill-defined because a monodromy around a loop in $\mathcal{BC}^*$

changes the AB phase of a test particle by $\exp[2\pi \mathrm{i}\langle \frac{1}{n}\sum_{j=1}^{n}\gamma^{(ij)},\gamma_g^{\mathrm{test}}\rangle]$. On the other hand, if we can find a suitable finite flavor subgroup $\mathcal{D}$ and an element d such that under the same monodromy transformation

$$\widehat{\delta\xi(d)} = -\frac{1}{n}\sum_{j=1}^{n}\gamma^{(ij)} \bmod \Gamma \tag{3.79}$$

then the AB phases of BPS test particles in the theory where the flavor group is gauged will be well-defined. In this case we can form a good theory of the surface defect, even though the torsor Γ_i is twisted.

Example 3.6.2. For an example of how we might find a suitable finite flavor group $\mathcal{D}$ in the IR, we return to the theories discussed in §3.4. We will show how to construct an extension like (3.71) with $\mathcal{D}\simeq\mathbb{Z}_n$, which will serve for anomaly cancellation. We begin with the exact sequence

$$0\to\mathcal{P}(z)\to H_1(\bar\Sigma-\pi^{-1}\{z\},\mathbb{Z})\to H_1(\bar\Sigma,\mathbb{Z})\to 0. \tag{3.80}$$

This sequence extends the charge lattice by a free abelian group of rank n,

$$\mathcal{P}(z)=\bigoplus_{i=1}^{n}P_i(z)\cdot\mathbb{Z}\simeq\mathbb{Z}^n, \tag{3.81}$$

generated by the homology classes of small circles $P_i(z)$ around the n preimages of z. Now consider the sublattice $\Lambda_n\subset\mathcal{P}(z)$ given by the kernel of the homomorphism $\mathcal{P}(z)\to\mathbb{Z}_n$ defined by $\sum a_iP_i(z)\mapsto\sum a_i \bmod n$. We can take a quotient of (3.80) by this lattice and its image in $H_1(\bar\Sigma-\pi^{-1}\{z\},\mathbb{Z})$ to produce

$$0\to\hat{\mathcal{D}}\to H_1(\bar\Sigma-\pi^{-1}\{z\},\mathbb{Z})/\Lambda_n\to H_1(\bar\Sigma,\mathbb{Z})\to 0. \tag{3.82}$$

where $\hat{\mathcal{D}}:=\mathcal{P}(z)/\Lambda_n\simeq\mathbb{Z}_n$. The actual charge lattice Γ_g is not exactly $H_1(\bar\Sigma,\mathbb{Z})$ but is closely related to it as a subquotient. The subgroup $\hat{\mathcal{D}}$ survives taking this subquotient, yielding a candidate for (3.71).

Now suppose we choose a system of charges $\gamma_{ij}\in\Gamma_{ij}$ as in §3.4.1. Using this data we can split gauge charges $\gamma_g^{\mathrm{ext}}\in H_1(\bar\Sigma-\pi^{-1}\{z\},\mathbb{Z})/\Lambda_n$ as $\xi\oplus\gamma_g$ where $\xi=[\sum a_iP_i]$ with

$$\sum_{i=1}^{n}a_i=\sum_{j=1}^{n}\langle\gamma_g^{\mathrm{ext}},\gamma_{ij}\rangle \quad \bmod n. \tag{3.83}$$

(One can check that the right hand side is indeed independent of i.)

Now let d be a primitive generator of $\mathcal{D}$. Let $\mathbb{S}$ be a surface defect with flavor holonomy d. Then, the flavor contribution to the AB phase of a particle of charge $\gamma_g^{\mathrm{test}}\in H_1(\bar\Sigma-\pi^{-1}\{z\},\mathbb{Z})/\Lambda_n$ transported around such a defect $\mathbb{S}$ will be

$$\xi(d)=\exp\left(-\frac{2\pi\mathrm{i}}{n}\left\langle\sum_j\gamma_{ij},\gamma_g^{\mathrm{test}}\right\rangle\right). \tag{3.84}$$

Monodromy of the local system Γ_{ij} around cycles in $\mathcal{B}C^*$ takes $\gamma_{ij} \to \gamma_{ij} + \gamma^{(ij)}$, and hence (3.84) satisfies (trivially) the anomaly cancellation condition (3.79).

Remarks:

1. We must stress that we have only given a potential mechanism for anomaly cancellation and we have not given a general proof that the problems of (3.39) can always be cured by conjecturing an extension of the lattice of gauge charges by characters for a $\mathbb{Z}_n$ flavor symmetry. Moreover, it is far from obvious that such tentative finite flavor charge groups $\mathcal{D}$ would always arise from actual symmetries of the bulk UV 4d theory. In §7.4 we explain that for the A_1 theories with regular singular points a suitable UV $\mathbb{Z}_2$ flavor symmetry does indeed exist. In addition, one can define a suitable flavor symmetry for the AD theories with N even.
2. Notice that if the surface defect carries a nontrivial flavor group label the non-integral monodromies of the Γ_i imply that the $\omega(\gamma, \gamma_i)$ are nonintegral. This is not so strange, considering that we have twisted boundary conditions around the surface defect for bulk particles. It is perfectly possible for bulk hypermultiplets with twisted boundary conditions to contribute to the 2d effective superpotential as a fraction of a 2d chiral multiplet. It would be nice to compute such contributions explicitly.

4. Line defects, interfaces, framed BPS states, and a spin 2d-4d wall-crossing formula

4.1 Interfaces

Past experience with the pure 4d wall-crossing formula suggests that a useful way of understanding it is to study supersymmetric *line defects* L [3]. In the 2d-4d context, in addition to the usual 4d line defects, we want to consider supersymmetric line defects which sit on a surface defect, dividing it into two pieces — or more generally, supersymmetric *interfaces* which have one surface defect $\mathbb{S}$ on the left and another surface defect $\mathbb{S}'$ on the right.

The supersymmetric interfaces we consider preserve 2 out of the 4 supercharges preserved by the surface defect. In the conventions of [3], they are

$$Q^1_1 - \zeta^{-1}\bar{Q}^1_{\dot{2}}, \qquad Q^2_2 + \zeta^{-1}\bar{Q}^2_{\dot{1}} \tag{4.1}$$

where the lower index is a spin index and the upper index is an $su(2)_R$ symmetry index. As in [3], ζ is a phase, but later will be analytically continued to $\mathbb{C}^\times$, and the interfaces in general depend on a lift to the universal cover $\widetilde{\mathbb{C}^\times}$.

Some simple examples of supersymmetric interfaces on surface defects can be constructed directly. For example, suppose we consider a supersymmetric gauge theory with gauge group G, with a Gukov-Witten-type surface defect $\mathbb{S}$ which breaks G to a Levi subgroup H. Then we can consider a supersymmetric Wilson line in an irreducible representation of H (which is *not* a representation of G). This gives a line defect which is restricted to lie on $\mathbb{S}$. Another useful machine for producing such interfaces is the "Janus" construction: here we allow a twisted F-term coupling on the surface defect to jump at some point x^3, and include an appropriate coupling at the interface to preserve supersymmetry [17].

In the presence of the interface, the IR physics depends on the choice of a vacuum i for $\mathbb{S}$ and j' for $\mathbb{S}'$. Having fixed that choice, the Hilbert space is graded by charges lying in $\Gamma_{ij'} = \Gamma_i - \Gamma_{j'}$.

4.2 Review of the 4d case: Framed BPS states and halos

Let us recall briefly the case studied in [3]. This is the case where $\mathbb{S}$ and $\mathbb{S}'$ are both the null surface defect. In this case we denoted a line defect of type ζ by L_ζ. Often ζ will be understood and we simply write L. The presence of the line defect modifies the Hilbert space of (one particle) states which is denoted $\mathcal{H}_{L,u}$, where $u \in \mathcal{B}$. Often the u-dependence will be suppressed in the notation. The Hilbert space $\mathcal{H}_L$ is graded by a Γ-torsor Γ_L and we write $\mathcal{H}_L = \oplus_{\gamma \in \Gamma_L} \mathcal{H}_{L,\gamma}$. The preserved supersymmetries lead to a BPS bound $E \geq -\mathrm{Re}(Z_\gamma/\zeta)$ in $\mathcal{H}_{L,\gamma}$ and states satisfying this bound are called *framed BPS states*. We define a protected spin character for these

$$\overline{\underline{\Omega}}(u, L_\zeta, \gamma; y) := \mathrm{Tr}_{\mathcal{H}^{\mathrm{BPS}}_{u,L,\zeta,\gamma}} (-1)^{2J_3} (-y)^{2\mathcal{J}_3} \tag{4.2}$$

where

$$\mathcal{J}_3 := J_3 + I_3, \tag{4.3}$$

and I_3 is an R-symmetry generator. We will often lighten the notation and just write $\overline{\underline{\Omega}}(L, \gamma; y)$. Specializing to $y = -1$ defines the framed BPS index $\overline{\underline{\Omega}}(L, \gamma)$.

When considered as functions of (u, ζ) the framed indices are piecewise constant but undergo wall-crossing, just like ordinary BPS degeneracies. The main physical justification for this is the "halo picture" described in [34, 3, 35, 48, 49]. Near BPS walls, defined for populated charges $\gamma \in \Gamma$ by,

$$W_\gamma := \{(u, \zeta) : Z_\gamma/\zeta < 0\} \tag{4.4}$$

some of the states in $\mathcal{H}^{\mathrm{BPS}}_{u,L,\zeta,\gamma}$ can be described as "halo configurations." They look like a "core" BPS state of the line defect very weakly bound to some "halo" particles. The distance between the line defect and the halo particles has a universal form depending only on the IR data: if γ^c, γ^h are the core and halo charges respectively, it is

$$r_h = \frac{\langle \gamma^c, \gamma^h \rangle}{2\,\mathrm{Im}\,(Z_{\gamma^h}/\zeta)}. \tag{4.5}$$

The bound state exists only when $r_h > 0$, i.e. only on one side of the wall; as we approach the wall from that side $r_h \to \infty$, so this bound state disappears from the framed BPS spectrum.

The framed BPS states which appear/disappear upon crossing the wall W_{γ^h} can be thought of as states in a Fock space consisting of states containing a core charge and some number of halo particles of charge γ^h orbiting around it. Since the halo particles are mutually BPS there are many such states, and they can be enumerated with Fock space combinatorics. In order to describe this more precisely we first introduce a generating function

$$F(u, L_\zeta, \{x_\gamma\}; y) := \sum_\gamma \overline{\underline{\Omega}}(u, L, \zeta, \gamma; y) x_\gamma. \tag{4.6}$$

where $x_\gamma x_{\gamma'} = x_{\gamma+\gamma'}$. In order to lighten the notation we will often drop various arguments, and sometimes simply write this as $F(L)$. Now, one isolates the contributions from a core charge γ^c surrounded by halo configurations of particles with charge γ^h. The $\mathbb{Z}_2$-graded Fock space is built on $(J_{\gamma^c,\gamma^h}) \otimes \tilde{\mathcal{H}}^{\mathrm{BPS}}(\gamma^h; u)$, where $\tilde{\mathcal{H}}^{\mathrm{BPS}}$ is the space of BPS states with the center-of-mass degree of freedom factored out. Here (J_{γ^c,γ^h}) is the spin representation with maximal spin $2J_3 = |\langle\gamma^c, \gamma^h\rangle|$. If we decompose

$$\Omega(u, \gamma^h; -z) = \mathrm{Tr}_{\tilde{\mathcal{H}}(\gamma^h;u)}(-z)^{2J_3} z^{2I_3} = \sum_{m\in\mathbb{Z}} a_{m,\gamma^h} z^m \tag{4.7}$$

then $a_m \geq 0$ for m even while $a_m \leq 0$ for m odd. There are creation operators $A^\dagger_{m',m,\alpha}$ with

$$-|\langle\gamma^c, \gamma^h\rangle| \leq m' \leq |\langle\gamma^c, \gamma^h\rangle| \tag{4.8}$$

while α runs over $|a_{m,\gamma^h}|$ values for each m. Somewhat counterintuitively, the oscillators with m even correspond to fermionic oscillators while those with m odd correspond to bosonic oscillators. The corresponding Fock space factor when a halo is created is thus

$$(1 + y^{m+m'} x_{\gamma^h})^{a_{m,\gamma^h}} \tag{4.9}$$

when m is even, and

$$\frac{1}{(1 - y^{m+m'} x_{\gamma^h})^{|a_{m,\gamma^h}|}} \tag{4.10}$$

when m is odd. These factors can be summarized as

$$(1 + (-1)^m y^{m+m'} x_{\gamma^h})^{a_{m,\gamma^h}}. \tag{4.11}$$

Reference [3] shows that the framed BPS indices (at $y = -1$) can be described by replacing x_γ by X_γ satisfying the twisted group law $X_\gamma X_{\gamma'} = (-1)^{\langle\gamma,\gamma'\rangle} X_{\gamma+\gamma'}$, and making an L-independent "coordinate" transformation of the X_γ,

$$X_{\gamma'} \to (1 - X_\gamma)^{\langle\gamma,\gamma'\rangle\Omega(\gamma)} X_{\gamma'}. \tag{4.12}$$

This transformation is nothing but (the inverse of) (2.11)! This is the key observation in one proof of the 4d wall-crossing formula for the $\Omega(\gamma)$ [3].

If one wants a wall-crossing formula for the full y-dependent framed BPS indices, then the above generating function must be split up into its contribution from various "core" charges. The reason for this is that the halo Fock spaces (in particular the range m' in (4.8)) depend on the core charge. This leads to somewhat awkward wall-crossing formulae.

However, as explained at length in [3] (see also [35, 49]) one can write an elegant formula by replacing F by the same expression evaluated on noncommuting variables,

$$\widehat{F}(u, L_\zeta, \{\hat{X}_\gamma\}; y) := \sum_\gamma \underline{\overline{\Omega}}(u, L, \zeta, \gamma; y)\hat{X}_\gamma, \tag{4.13}$$

where

$$\hat{X}_\gamma \hat{X}_{\gamma'} = y^{\langle\gamma,\gamma'\rangle}\hat{X}_{\gamma+\gamma'}, \tag{4.14}$$

Again, we will abbreviate this by $\widehat{F}(L)$.

The transformation of $\widehat{F}$ across walls W_γ for populated charges γ can be expressed in terms of conjugation by products of quantum dilogs evaluated on $\hat{X}_\gamma$, where the product is determined by the data $a_{m,\gamma}$. See [50, 3] for details.

4.3 Analog in two dimensions

Let us now consider the special case where the four-dimensional theory is null but there are nontrivial surface defects separated by an interface $\mathbb{S}L_\zeta\mathbb{S}'$. It is tempting to apply the approach of §4.2 to try to understand the purely 2d wall-crossing formula as well. We do this in the present section, although our arguments in this case will be somewhat less rigorous. Just as in the 4d situation, we can define "framed BPS states" in the Hilbert space of the theory with a supersymmetric interface L_ζ inserted between two surface defects. Then we describe the wall-crossing of the framed BPS degeneracies.

At least in specific models such as supersymmetric Landau-Ginzburg models, there is a picture of this wall-crossing which is much like that in the 4d case: there are "halo" states consisting of a single 2d particle loosely bound to the line defect. Although there is not a precise formula for this halo radius it can be shown that any reasonable measure of such a radius has the crucial property that it diverges at walls W_{ij} where $Z_{ij}/\zeta \in \mathbb{R}_-$. See Appendix C. At such walls, halo states appear or disappear from the framed BPS spectrum.

As in the 4d case, after introducing appropriate generating functions $F(L)$ for the framed BPS states, this picture leads to a prediction for the transformation of $F(L)$ across the walls. The transformation is

$$F(L) \to (1 - \mu_{ij}X_{ij})\, F(L). \tag{4.15}$$

For 2d particles on the right of the line defect, similarly

$$F(L) \to F(L)\,(1 + \mu_{ij}X_{ij})\,. \tag{4.16}$$

Note the change of sign between (4.15) and (4.16). We will see later that this is necessary for the wall-crossing invariance of the algebra of OPEs of line defects.

The transformation (4.15) is precisely (2.2). Starting from this observation, and following again the approach of [3], we get a new way of understanding the 2d wall-crossing formula.

Our discussion of the "halo states" above invoked facts specific to the Landau-Ginzburg model. It would be very natural to believe that these states in fact exist in any 2d theory and have the same behavior we have described; if so, this would give a real derivation of the 2d wall-crossing formula using only the halo picture. Unfortunately (and in contrast to the 4d case) it seems difficult to argue directly that this behavior is a universal feature depending only on the infrared physics. The difficulty is that we are considering infrared theories which are massive: the forces between the halo particle and the line defect are generated by irrelevant operators which are, after all, irrelevant at low energies. In [20] a full proof of the 2d wall-crossing formula was given, independent of the details of the theory, but it required the heavier machinery of tt^* geometry.

4.4 Framed indices in the 2d-4d case

Let us now consider the general 2d-4d case. There is a Hilbert space of one-particle states in the presence of surface defects. If we have vacua $i \in \mathcal{V}(\mathbb{S})$ and $j' \in \mathcal{V}(\mathbb{S}')$ then the Hilbert space is graded by the Γ-torsor $\Gamma_{ij'}$,

$$\mathcal{H}_{u,\mathbb{S}L_\zeta\mathbb{S}'} = \bigoplus_{i\in\mathcal{V}(\mathbb{S}),j'\in\mathcal{V}(\mathbb{S}')}\ \bigoplus_{a\in\Gamma_{ij'}} \mathcal{H}_{u,\mathbb{S}L_\zeta\mathbb{S}',a}. \tag{4.17}$$

As usual we will lighten the notation to $\mathcal{H}_L$ and $\mathcal{H}_{L,a}$.

The two supercharges (4.1) preserved by the line defect imply a BPS bound: states of charge $\gamma_{ij'}$ have

$$E \geq -\mathrm{Re}\ (Z_{\gamma_{ij'}}/\zeta). \tag{4.18}$$

So just as reviewed above, we can consider *framed BPS states* saturating this bound. Moreover, we can introduce a framed BPS degeneracy:

$$\underline{\overline{\Omega}}(u, \mathbb{S}L_\zeta\mathbb{S}', a; y) := \mathrm{Tr}_{\mathcal{H}^{\mathrm{BPS}}_{L,a}} e^{\mathrm{i}\pi F}(-y)^{\mathcal{J}} \tag{4.19}$$

As usual we abbreviate this by $\underline{\overline{\Omega}}(L, a; y)$ and the value at $y = -1$ by $\underline{\overline{\Omega}}(L, a)$. Here we have made a choice of $u(1)_V$ charge generator F. This generates a one-dimensional abelian Lie algebra and hence its spectrum is some set of real numbers, not necessarily integer-spaced. Moreover we have introduced the operator $\mathcal{J} := 2J_{12} + 2I_{12}$ which commutes with the preserved supercharges. Again the spectrum is some set of real numbers, not necessarily integer-spaced.

The framed BPS degeneracies will undergo wall-crossing across two kinds of walls. First, halo states will be gained/lost when crossing walls of type W_γ for populated charges $\gamma \in \Gamma$. Such states are shown in Figure 10. Second, there will also be wall-crossing across *soliton walls* (again for populated charges):

$$W_{\gamma_{ik}} := \{(u,\zeta) : Z_{\gamma_{ik}}/\zeta < 0\} \qquad i, k \in \mathcal{V}(\mathbb{S}), \tag{4.20}$$

$$W_{\gamma_{k'j'}} := \{(u,\zeta) : Z_{\gamma_{k'j'}}/\zeta < 0\} \qquad k', j' \in \mathcal{V}(\mathbb{S}'). \tag{4.21}$$

Across the walls (4.20), a bound state involving a 2d "halo" particle with charge γ_{ik} and degeneracy $\mu(\gamma_{ik})$, sitting to the left of a core state of the line defect, appears or disappears. Such states are illustrated in Figure 11. Similarly (4.21) involves bound states where the halo particle is on the right.



Figure 10: A 4d halo particle of charge γ^h is bound to an interface in charge sector $\gamma^c_{ij'}$ (the "core charge") to produce a framed BPS state of charge $\gamma_{ij'} = \gamma^h + \gamma^c_{ij'}$.



Figure 11: A soliton of type ik and 4d electromagnetic charge γ^h_{ik} is bound to an interface in charge sector $\gamma^c_{kj'}$ to produce a framed BPS state of charge $\gamma_{ij'} = \gamma^h_{ik} + \gamma^c_{kj'}$. A similar picture applies to solitons of charge $\gamma^h_{k'j'}$ bound on the right.

In §4.7 we will interpret the transformations (2.27) and (2.30) which enter the wall-crossing formula in terms of these "halo" states. The interpretation of (2.27) should be much like what we have just discussed in the pure 2d case. Similarly, we should interpret the wall-crossing associated with W_γ in terms of a halo of particles of charge γ, which surrounds a core state of the line defect with charge $\gamma_{ij'} = \gamma_i - \gamma_{j'}$. However, since there are a number of potentially subtle points in the halo discussion, we first make some general remarks on the structure of the wall-crossing formula.

4.5 Ring of line defects

In [3] an important role was played by an algebra of line defects whose coefficients are vector spaces. There is a natural generalization to the case of supersymmetric interfaces.

Indeed, consider two interfaces $\mathbb{S}L_\zeta\mathbb{S}'$ and $\mathbb{S}'L'_\zeta\mathbb{S}''$ preserving the same supersymmetry. We can define a new interface $\mathbb{S}\,(L \circ L')_\zeta\,\mathbb{S}''$ by a (nonsingular!) OPE, bringing them against each other along the surface defect. It should be possible to compute the framed BPS generating function of the composition of the two defects by counting the framed BPS states of the system of two well-separated line defects L and L': the index should be independent of the distance between the line defects. If the distance between the defects is sufficiently large, the Hilbert space of ground states of the system should be the tensor product of the Hilbert spaces of the two subsystems. We should allow the strip of surface defect between the two line defects to be in any of its possible vacua. The only IR subtlety we should take care of is the fermion number grading, which can be affected by the gauge charges carried by the ground states: a system of two well-separated dyons might carry fractional amounts of angular momentum hidden in the electro-magnetic fields.

The coefficients of this OPE are vector spaces which are graded by F and $\mathcal{J}$ [3]. Replacing these vector spaces by their characters defines a deformed product $L \circ_y L'$. As in [3] it is useful to describe this product by introducing the "quantum" generating function

$$\widehat{F}(L) := \sum_{i \in \mathcal{V}(\mathbb{S}), j' \in \mathcal{V}(\mathbb{S}'), \gamma_{ij'} \in \Gamma_{ij'}} \underline{\overline{\Omega}}(L_\zeta, \gamma_{ij'}; y) \hat{X}_{\gamma_{ij'}} \tag{4.22}$$

where, for the moment, $\hat{X}_{\gamma_{ij'}}$ are just placeholders.

We now claim that with a suitable multiplication law on $\hat{X}_a$ we have the rule

$$\widehat{F}(L \circ_y L') = \widehat{F}(L)\widehat{F}(L') \tag{4.23}$$

that is, $\widehat{F}$ is a homomorphism from the algebra of interfaces to the groupoid algebra[24] $\mathbb{C}[\mathbb{V}]$.

In order to find the requisite multiplication law on $\hat{X}_a$ we use the relation

$$\mathcal{H}_{L \circ_y L', c} = \oplus_{a+b=c} \mathcal{H}_{L,a} \otimes \mathcal{H}_{L',b} \otimes \mathcal{N}_{a,b} \tag{4.24}$$

where $\mathcal{N}_{a,b}$ is a one-dimensional representation of $so(2)_{12} \oplus u(1)_r$. Let us define $\sigma(a,b)$ and $n(a,b)$ via:

$$e^{\mathrm{i}\pi F}|_{\mathcal{N}_{a,b}} := \sigma(a,b) \tag{4.25}$$

$$(-y)^{\mathcal{J}}|_{\mathcal{N}_{a,b}} := (-y)^{n(a,b)} \tag{4.26}$$

Note that $\sigma(a,b)$ is in general a phase and $n(a,b)$ need not be integer because of angular momentum in massless degrees of freedom.

Equation (4.24) is reasonable because we are working with indices and therefore – by topological invariance – we can separate the line defects an arbitrary distance from each other. The states can factorize but there can be shifts in the action of various $u(1)$ generators.

Now, equation (4.23) will hold *provided*

$$\hat{X}_a \hat{X}_b = \sigma(a,b)(-y)^{n(a,b)} \hat{X}_{a+b} \tag{4.27}$$

for composable morphisms a, b. We continue to take $\hat{X}_a \hat{X}_b = 0$ if a, b are not composable. Associativity is guaranteed if $n(a,b)$ and $\sigma(a,b)$ are 2-cocycles on the groupoid. From the case with $\mathbb{S}$ and $\mathbb{S}'$ the null surface defects and the choice $F = 2J_{12}$ we know from [3] that $n(\gamma, \gamma') = \langle \gamma, \gamma' \rangle$ and $\sigma(\gamma, \gamma') = (-1)^{\langle \gamma, \gamma' \rangle}$. Now, these cocycles can be shifted by coboundaries. Physically, these correspond to shifts in the definitions of the generators F and $\mathcal{J}$. By an argument analogous to that at the end of §3.3.5 we may assume that $\sigma(a,b)$ are signs and $n(a,b)$ are integers, and we will henceforth make that assumption.

---

[24] The existence of supersymmetric interfaces extends the groupoid algebra to include that for several surface defects. We understand $\mathbb{V}$ to mean this extended groupoid.

4.6 A 2d-4d spin wall-crossing formula

Now we come to a crucial point. Wall-crossing must be compatible with the multiplication of line defects, since the latter is defined in the UV and hence independent of the choice of the vacuum. Therefore the wall-crossing transformation must be an automorphism of our groupoid algebra (4.27). This is a matrix algebra over the quantum torus, hence Morita equivalent to the coordinate algebra of the quantum torus, and hence a simple algebra. Therefore the automorphism must be inner. Therefore, across a wall of type W_γ, $\widehat{F}(L)$ should transform according to

$$\hat{T}_\gamma : \hat{X}_a \mapsto \Phi_{\mathbb{S},\gamma} \hat{X}_a \Phi^{-1}_{\mathbb{S}',\gamma}. \tag{4.28}$$

As we will see below, the halo picture of wall-crossing suggests that $\Phi_{\mathbb{S},\gamma}$ must have the form

$$\Phi_{\mathbb{S},\gamma} = \prod_{i \in \mathcal{V}(\mathbb{S})} \prod_{s \in \mathcal{S}_{i,\gamma}} (1 + \varphi_{i,s}(y) \hat{X}_{\gamma_{ii}})^{d(i,s)}, \tag{4.29}$$

with an analogous expression for $\Phi_{\mathbb{S}',\gamma}$. Here $\mathcal{S}_{i,\gamma}$ is some countably infinite set of real numbers depending on i, γ and $\varphi_{i,s}(y)$ is a "monomial in y," that is, it is a phase times a (possibly fractional) power of y. The important thing is that $\Phi_{\mathbb{S},\gamma}$ does not depend on the detailed core charge a being conjugated.[25]

Similarly, across walls of the kind $W_{\gamma_{ij}}$ we have again a conjugation (4.28), but now with

$$\Phi_{\mathbb{S},\gamma_{ij}} = (1 + \varphi_{\gamma_{ij}}(y) \hat{X}_{\gamma_{ij}}). \tag{4.30}$$

Here $\varphi_{\gamma_{ij}}(y)$ is a sum of monomials in y. (See (4.37) below.)

Given the transformation (4.28), we can use the same argument as in [3] to deduce a 2d-4d analog of the Kontsevich-Soibelman motivic wall-crossing formula. Given a path $\mathcal{P}$ between two points in $\mathcal{B} \times \mathbb{C}^\times$ the accumulated transformation of $\widehat{F}(L)$ is

$$\widehat{F}(L) \to T^{\text{left}}(\mathcal{P}) \widehat{F}(L) T^{\text{right}}(\mathcal{P}) \tag{4.31}$$

where

$$T^{\text{left}}(\mathcal{P}) =: \prod_b \Phi_{\mathbb{S},b} : \tag{4.32}$$

is the path-ordered product of the transformations taken across walls W_b of type (4.4) or (4.20) crossed by the path $\mathcal{P}$. Given two paths $\mathcal{P}$ and $\mathcal{P}'$ with common endpoints such that the local system does not have monodromy around the corresponding closed path, we can state the spin 2d-4d wall-crossing formula in the form:

$$T^{\text{left}}(\mathcal{P}) = T^{\text{left}}(\mathcal{P}'). \tag{4.33}$$

(An equivalent formula can be written with T^{right}.) A generalization can be stated which takes into account the monodromy of an arbitrary path $\mathcal{P}(\mathcal{P}')^{-1}$.

---

[25] In equation (4.29) we have assumed, for simplicity, that on W_γ the only populated charges with central charge Z parallel to Z_γ are particles that themselves have electromagnetic charge γ. We continue to make that assumption below.

4.7 Halo picture

We would now like to relate the quantities $\mathcal{S}_{i,\gamma}$, $\varphi_{i,s}(y)$, $d(i,s)$ and $\varphi_{\gamma_{ij}}(y)$ introduced in the wall-crossing formula above to physical quantities determined by Hilbert spaces of BPS states.

4.7.1 Crossing soliton walls

The first main statement of framed wall-crossing is that across $W_{\gamma_{ik}}$ we have

$$\Delta\mathcal{H}^{\mathrm{BPS}}_{L,\gamma_{ij'}} = \pm\tilde{\mathcal{H}}^{\mathrm{BPS}}_{\gamma_{ik}} \otimes \mathcal{H}^{\mathrm{BPS}}_{L,\gamma_{kj'}} \otimes \mathcal{N}_{\gamma_{ik},\gamma_{kj'},L} \tag{4.34}$$

as representations of the abelian Lie algebra $u(1) \oplus u(1)$ generated by F and $\mathcal{J}$. Here $\mathcal{N}_{\gamma_{ik},\gamma_{kj'},L}$ is a one-dimensional representation with character $e^{\mathrm{i}\pi f_{\mathcal{N}}}(-y)^{\mathcal{J}_{\mathcal{N}}}$. The overall sign of the right hand side depends on the direction the wall is crossed.

To express this in terms of generating functions, define

$$\mu(\gamma_{ij}; y) := \mathrm{Tr}_{\tilde{\mathcal{H}}^{\mathrm{BPS}}_{\gamma_{ik}}} e^{\mathrm{i}\pi F}(-y)^{\mathcal{J}}. \tag{4.35}$$

Comparing with the wall-crossing from (4.30) and taking traces we learn that

$$\varphi_{\gamma_{ik}}(y) = \pm\mu(\gamma_{ik}; y)\left(\frac{\sigma(\gamma_{ik},\gamma_{kj'})}{e^{\mathrm{i}\pi f_{\mathcal{N}}}}\right) y^{\mathcal{J}_{\mathcal{N}}-n(\gamma_{ik},\gamma_{kj'})}. \tag{4.36}$$

Putting $y = -1$ we find agreement with the formal structure described in equation (2.27) provided that $e^{\mathrm{i}\pi F}$ on the one-dimensional space $\mathcal{N}_{\gamma_{ik},\gamma_{kj'},L}$ has the value $\sigma(\gamma_{ik},\gamma_{kj'})$. Similarly, it is natural to guess that $n(a,b)$ defined in (4.26) satisfies $\mathcal{J}_{\mathcal{N}} - n(\gamma_{ik},\gamma_{kj'}) = 0$ and hence we conjecture that

$$\varphi_{\gamma_{ik}}(y) = \pm\mu(\gamma_{ik}; y). \tag{4.37}$$

4.7.2 Crossing 4d walls

Let us try to describe the halo particles relevant to wall-crossing associated with W_{γ}. We choose a specific IR Lagrangian description of the setup, i.e. a particular $\gamma_i^0 \in \Gamma_i$ and $\gamma_{j'}^0 \in \Gamma_{j'}$. As we have mentioned before, γ_i^0, $\gamma_{j'}^0$ measure the amount of electromagnetic flux which is threaded through the surface defects. The interface sources a total flux $\gamma_{ij'}$, of which $\gamma_i^0 - \gamma_{j'}^0$ is carried by the surface defects, while the remaining

$$\gamma^c = \gamma_{ij'} - \gamma_i^0 + \gamma_{j'}^0 \in \Gamma \tag{4.38}$$

emerges into space. Thus, the duality invariant gauge field is

$$\mathbb{F} = \gamma_{ij'}\omega_{S^2} + \gamma_{j'}^0\delta_N - \gamma_i^0\delta_S \tag{4.39}$$

See Figure 12. Here ω_{S^2} is the unit volume form of the sphere and δ_N, δ_S are unit weight delta functions at the north and south poles. The decomposition (4.39) is noncanonical, but $\mathbb{F}$ is canonically defined and satisfies Dirac quantization. (The projection of the charges to Γ_g is understood here.)



Figure 12: The configuration of electromagnetic flux around an interface between two surface defects. The total outgoing flux through the blue sphere is $\gamma_{ij'} + \gamma_{j'}^0 - \gamma_i^0 \in \Gamma$.

The "halo" then consists of 4d particles which couple to this flux $\gamma_{ij'}$ as in the usual 4d case, *plus* the contribution from 2d particles on the two surface defects. While these two contributions are not separately well-defined (they depend on our choice of Lagrangian), their sum must make good physical sense. As usual, the halo particles generate a Fock space, and in order to discuss the wall-crossing at the level of state spaces (i.e. at the "categorified level") we need to describe the oscillators which generate the Fock space. In Appendix D we analyze the Landau level problem for halo particles of charge γ in the presence of the field (4.39). In a description where there are no chiral multiplets on the surface defect, the halo particles may be described by fermionic and bosonic creation operators $A^\dagger_{m',m,\alpha}$ where α again runs over $|a_{m,\gamma^h}|$ values for every integer m, but now m' has a fixed fractional part and satisfies the inequality (D.12). The number of such m' values is the function N_{κ_1,κ_2} given in (D.13). The signed sum of the number of oscillators is given by (D.14). We identify this quantity with $\omega(\gamma, \gamma_{ij'})$.

The framed wall-crossing formula for these walls says that across W_γ the Hilbert space of framed BPS states with core charge $\gamma_{ij'} + N\gamma$ for some N gains or loses a Fock space factor generated by $A^\dagger_{m',m,\alpha}$. To express this formula more concisely we introduce a generating function F, analogous to (4.6), valued in the untwisted groupoid algebra:

$$F(u, L_\zeta, \{x_a\}; y) := \sum_{i\in\mathcal{V}(\mathbb{S}), j'\in\mathcal{V}(\mathbb{S}')} \sum_{a\in\Gamma_{i,j'}} \overline{\underline{\Omega}}(L, a; y) x_a, \tag{4.40}$$

which will be abbreviated to $F(L)$. The x_a satisfy (2.25) with $\sigma(a, b) = 1$.

As in the pure 4d case, it is useful to define the "part of the generating function with core charge $\gamma_{ij'}$",

$$F_{\bar{\gamma}_{ij'}} := \sum_{n\in\mathbb{Z}} \overline{\underline{\Omega}}(L, \gamma_{ij'} + n\gamma; y) x_{\gamma_{ij'}+n\gamma}. \tag{4.41}$$

Now each halo particle of type $A^\dagger_{m',m,\alpha}$ contributes a Fock space factor

$$(1 + (-1)^m y^{m'+m} x_\gamma), \tag{4.42}$$

and hence the transformation law across W_γ is:

$$T_\gamma : F_{\bar\gamma_{ij'}} \to \prod_{m,m'} (1+(-1)^m y^{m'+m} x_\gamma)^{a_{m,\gamma}} F_{\bar\gamma_{ij'}}. \tag{4.43}$$

As before, we are assuming that the only 4d particles with central charge parallel to Z_γ along W_γ in fact are those with charge γ.

Similarly, there is a transformation law for crossing soliton walls $W_{\gamma_{ik}}$:

$$T_{\gamma_{ik}} : F(L) \to (1 \pm \mu(\gamma_{ik}) x_{\gamma_{ik}}) F(L) (1 \mp \mu(\gamma_{ik}) x_{\gamma_{ik}}) \tag{4.44}$$

where the sign is determined by the direction in which the wall is crossed.

We now face an issue analogous to that in the pure 4d story, discussed just above (4.13). The transformation (4.43) is written directly in terms of the physical halo particles, but has the awkward feature that the prefactor depends on the core charge $\gamma_{ij'}$, through the range of m'. On the other hand, in §4.6 we found that there should be a nicer way of expressing the framed wall-crossing: indeed, it should be possible to write it as conjugation by an operator $\Phi_{\mathbb{S},\gamma}$ depending only on the wall which is crossed, not on the core charge. What we would really like is to determine $\Phi_{\mathbb{S},\gamma}$ physically, in terms of the halo picture we have been discussing in this section. We indicate briefly how to do this in §4.7.3.

While we will not carry out that analysis to the end, there is one specialization that is easier to deal with. Beginning with (4.43) and (4.44), taking $y \to -1$, converting to twisted (but commuting) variables, and making use of (D.14), we recover the transformations (2.30) and (2.27) which appeared in the 2d-4d wall-crossing formula of §2. In particular, after this specialization the spin 2d-4d wall-crossing formula (4.33) reduces to the ordinary (non-spin) 2d-4d wall-crossing formula we use in the rest of this paper.

4.7.3 Determining the generalized quantum dilogarithms $\Phi_{\mathbb{S},\gamma}$

In this section we finally explain how to relate the $\Phi_{\mathbb{S},\gamma}$ appearing in the framed wall-crossing across W_γ to physical data defined in terms of halo Fock spaces.

We define a linear map Ψ between the quantum groupoid algebra generated by $\hat X_a$ satisfying (4.27) and the (untwisted) groupoid algebra generated by x_a: $\Psi(\widehat{X}_a) = x_a$. Note that Ψ is *not* a ring homomorphism!

Compatibility of the framed wall-crossing formula with the physical halo picture is simply the statement that the following diagram commutes:

$$\begin{tikzcd} \widehat{F} \arrow[r, "\hat T_\gamma"] \arrow[d, "\Psi"] & \widehat{F} \arrow[d, "\Psi"] \\ F \arrow[r, "T_\gamma"] & F \end{tikzcd} \quad . \tag{4.45}$$

This condition can be used to write recursion relations for the exponents $d(i,s)$ and $d(j',t)$ in terms of $a_{m,\gamma}$, γ and $\gamma_{ij'}$. It can also be used to obtain the monomials $\varphi_{i,s}(y)$.

As an illustration of how this works, let us consider the formula

$$\hat X_a \hat X_b = \frac{\sigma(a,b)}{\sigma(b,a)} (-y)^{n(a,b)-n(b,a)} \hat X_b \hat X_a, \tag{4.46}$$

valid when both $a+b$ and $b+a$ are composable morphisms. In particular this holds for $a=\gamma$ and $b=\gamma_{ij'}$, and we can use this to rewrite (4.28) as

$$\hat{X}_a \to \prod_{i,s}(1+\varphi_{i,s}(y)\hat{X}_\gamma)^{d(i,s)} \prod_{j',t}(1+\tilde{\varphi}_{j',t}(y)\hat{X}_\gamma)^{-d(j',t)}\hat{X}_a, \tag{4.47}$$

where $\tilde{\varphi}_{j',t}(y) = \frac{\sigma(a,\gamma)}{\sigma(\gamma,a)}(-y)^{n(a,\gamma)-n(\gamma,a)}\varphi_{j',t}(y)$.

The products on the left only depend on the $\hat{X}_a$ through the single variable $\hat{X}_\gamma$ and hence can be treated as functions of a commutative variable. In particular they can be expanded as a series $(1+\sum_{N\geq 0}\Phi_N \hat{X}_{N\gamma})$, which can be combined to rewrite the right hand side of (4.47) in the form

$$\hat{X}_a + \sum_{N\geq 1}\Phi_N \sigma(N\gamma,a)(-y)^{n(N\gamma,a)}\hat{X}_{N\gamma+a}. \tag{4.48}$$

Now we can easily apply Ψ since that map simply takes monomials to monomials. Comparing the result with the transformation (4.43) of F, we would find recursion relations implying that $\Phi_{\mathbb{S},\gamma}$ is an infinite product, generalizing the quantum dilogarithm which appeared in the 4d case.

It would be worthwhile working out further details and examples of the spin version of the 2d-4d wall-crossing formula, but we leave this to another occasion.

5. Compactification to 3d and hyperholomorphic bundles

So far we have written the 2d-4d wall-crossing formula, explained in detail a physical context to which it is relevant, and given a physical derivation of it based on the halo phenomenon. We now turn to the 2d-4d analog of tt^* geometry.

5.1 The dimensionally reduced action

We consider the compactification of the combined 2d-4d theory on a circle of radius R. A surface defect which wraps the compactification circle will give rise to a line defect in three dimensions. As such, at energies much lower than $1/R$ we will obtain an effective 1d-3d theory, whose effective Lagrangian we wish to study. If $1/R$ is much smaller than the scale where the 4d IR description breaks down, then a good approximation to the 1d-3d Lagrangian will be obtained by naive dimensional reduction of the 2d-4d IR theory. This gives the exact 1d-3d Lagrangian in the limit $R\to\infty$. For finite R there are quantum corrections which can be interpreted as coming from BPS particles of the 2d-4d theory going around the compactification circle; these corrections are crucial to our story, but for now we are only discussing the limit of infinite R. As for the metric on the target space, this was discussed in detail in [1]. After patching, the dimensional reduction of the bulk 4d term in (3.12) gives a sigma model into a manifold $\mathcal{M}$, a torus bundle over the 4d Coulomb branch $\mathcal{B}$, with a simple explicit metric g^{sf}.

To find the corresponding 1d action we turn to the reduction of the term integrated over the surface defect, namely, the second line of (3.17). Before reduction this term can be written as

$$\int \mathrm{d}x^0 \mathrm{d}x^3 \, \left(\eta_I F^I_{03} - \alpha^I G_{03,I}\right) \tag{5.1}$$

where the dual gauge field is given by (3.18). The auxiliary field has been integrated out and the resulting infinite expression is cancelled by taking into account the last term in the definition of $G_{03,I}$. We can now dualize all of the gauge fields to scalar fields in 3d according to

$$\partial_i \theta_\gamma = \oint \mathrm{d}x^3 \langle \gamma, \mathbb{F}_{i3} \rangle. \tag{5.2}$$

This defines a scalar field θ valued in $\Gamma^* \otimes \mathbb{R}/(2\pi\mathbb{Z})$. [26] Note that for flavor charges $\gamma_f \in \Gamma_f$ we learn that θ_{γ_f} are constant. After choosing a duality frame we define $\theta^I_e := \theta_{e^I}$ and $\theta_{m,I} := \theta_{e_I}$. Note that for a charge $\gamma = p^I e_I + q_I e^I$ we have $\mathrm{d}\theta_\gamma = \mathrm{d}\theta \cdot \gamma = q_I \mathrm{d}\theta^I_e + \mathrm{d}\theta_{m,I} p^I$.

Thus, (5.1) leads to a very natural supersymmetric line defect in the sigma model. Its bosonic part is the path-ordered exponential of a simple $U(1)$ connection

$$\exp \int \mathrm{d}x^0 A^{\mathrm{sf}} \tag{5.3}$$

where in this expression we mean the integral of the pullback of the locally-defined one-form on $\mathcal{M}$:

$$A^{\mathrm{sf}} = \mathrm{i}\nu \cdot \mathrm{d}\theta = \mathrm{i}(\eta_I \mathrm{d}\theta^I_e + \alpha^I \mathrm{d}\theta_{m,I}). \tag{5.4}$$

When we wish to emphasize the choice of the vacuum i of $\mathbb{S}$ we write A^{sf}_i.

5.2 Globalizing

5.2.1 Line bundles

As emphasized in §3.3.5, the 2d-4d action is only locally defined on $\mathcal{B}$: there is no single Lagrangian description which encompasses the whole IR theory. So strictly speaking, we must perform the dimensional reduction in each patch and then glue the dimensionally reduced theories together, taking account of the necessary electromagnetic duality transformations and shifts of $\mathcal{W}_i$. The expression A^{sf}_i is duality invariant and will glue nicely as a one-form across patches requiring an electric-magnetic duality transformation. On the other hand, the superpotential $\mathcal{W}_i$ is not globally defined. It determines, locally, a choice of basepoint $\gamma^0_i \in \Gamma_i$. A change of superpotential is equivalent to $\gamma^0_i \to \gamma^0_i + \tilde{\gamma}$, with $\tilde{\gamma} \in \Gamma$. This leads to a shift of ν_i by $\nu_i \to \nu_i + \tilde{\gamma}$ which in turn induces a gauge transformation

$$A^{\mathrm{sf}}_i \to A^{\mathrm{sf}}_i + \mathrm{i}\mathrm{d}\theta_{\tilde{\gamma}}. \tag{5.5}$$

Therefore, we can cover $\mathcal{B}$ with patches $\mathcal{U}_\alpha$ so that on patch overlaps $\mathcal{U}_{\alpha\beta}$ the semiflat connections differ by gauge transformations of the form (5.5) for some $\tilde{\gamma}_{i,\alpha\beta} \in \Gamma$ which satisfy the cocycle condition on triple overlaps. We can therefore interpret A^{sf}_i as the connection on a line bundle V_i over $\mathcal{M}$, whose transition functions are $e^{\mathrm{i}\theta_{\tilde{\gamma}_{i,\alpha\beta}}}$. (This statement will need to be slightly amended. See the next section below.) Changing the local description by taking $\gamma_i^0 \to \gamma_i^0 + \tilde{\gamma}_\alpha$ in patch $\mathcal{U}_\alpha$ (with $\tilde{\gamma}_\alpha \in \Gamma$) changes the local framing of V_i and modifies the transition functions by a coboundary. Below we will identify V_i as the vacuum line bundle (for vacuum i) of the reduced one-dimensional quantum mechanical problem.

---

[26] To be precise, there are some subtle shifts of π when adding the scalars θ_γ, due to the supersymmetry-preserving boundary conditions. The correct additivity statement is that $e^{\mathrm{i}\theta_\gamma} e^{\mathrm{i}\theta_{\gamma'}} = (-1)^{\langle \gamma, \gamma' \rangle} e^{\mathrm{i}\theta_{\gamma+\gamma'}}$

It is worth remarking that if we interpret ν as a section of the torus fibration for the algebraic integrable system $\widetilde{\mathcal{M}}$ of the theory as in §3.3.6 then the connection $A^{\rm sf}$ naturally corresponds to a flat connection $A^{\rm sf}$ on the dual torus. Therefore, when restricted to a torus fiber we can view $A^{\rm sf}$ as the restriction of the canonical connection on the Poincaré line bundle to $\Gamma_g^* \otimes \mathbb{R}/(2\pi\mathbb{Z})$.

5.2.2 Vector bundles

There is an important subtlety we have suppressed above: the index i labeling the vacuum is not globally defined — rather it labels a sheet of the ramified cover $\mathcal{B}_{\mathbb{S}} \to \mathcal{B}$. This implies that the line bundles V_i also do not exist globally over $\mathcal{M}$. What does exist globally over $\mathcal{M}$ is their direct sum, a rank n vector bundle which we denote as

$$V_{\mathbb{S}} := \bigoplus_i V_i. \tag{5.6}$$

If we use trivializations given by local superpotentials as above then the transition functions on patch overlaps $\mathcal{U}_{\alpha\beta}$ will be either diagonal matrices with i^{th} element $e^{\mathrm{i}\theta_{\tilde{\gamma}_{i,\alpha\beta}}}$, or (constant) permutation matrices. $V_{\mathbb{S}}$ carries a connection $A^{\rm sf}$ which is diagonal (in each patch) with respect to the decomposition into the V_i. Given the above transition functions we can define a Hermitian structure by declaring that the framings $s_{i,\alpha}$ on $\mathcal{U}_\alpha$ implicit in choosing the gauge (5.4) form an orthonormal basis for $V_{\mathbb{S}}$. We will refer to this as the *unitary framing.*

We hasten to add that equation (5.6) is only a statement about $\mathcal{C}^\infty$ vector bundles. In particular, the diagonal structure of $A^{\rm sf}$ is an artifact of the $R \to \infty$ limit. At finite values of R the quantum effects coming from 2d solitons (whose worldlines are reinterpreted as 1d instantons) imply that the connection A on the bundle $V = V_{\mathbb{S}}$ will be corrected: in particular, it is no longer diagonal with respect to the decomposition of V as a sum of line bundles given in (5.6). In the framing we have described above, the connection A will be exponentially close to a diagonal connection for large R. The quantum-corrected connection is a rather nontrivial and interesting object: it is the 2d analogue of the quantum-corrected hyperkähler metric on $\mathcal{M}$ which played an important role in [1]. We will see below how to construct it more explicitly using results of §5.6 and Appendix E.

5.2.3 Twisted vector bundles and mirror symmetry

In §3.4.1 we explained that the torsors Γ_i might not exist as Γ-torsors. In such cases there will be further complications in the geometrical construction of $V_{\mathbb{S}}$ since the fractional shifts of ν are not good gauge transformations. As further explained in §3.4.1, under some conditions one *can* construct the Γ_i as *twisted* torsors. In these cases the Γ_i will have have monodromy shifts by $\frac{1}{n}\Gamma$ for some integer n. In such cases the overlap transition functions $e^{\mathrm{i}\theta_{\tilde{\gamma}_{i,\alpha\beta}}}$ will be ambiguous by an n^{th} root of unity. Even if we choose a good cover, so that we can choose an unambiguous n^{th} root on each patch overlap $\mathcal{U}_{\alpha\beta}$, the cocycle condition can fail on triple overlaps by an n^{th} root of unity. Such an object defines what is known as a *twisted bundle*. In principle, from the monodromies of the torsors Γ_{ij} one can construct the gerbe of the twisting. It would be desirable to have a more streamlined version of this construction.

In string theory, twisted bundles are associated with background B fields [51]. The role of B-fields in three-dimensional sigma models is discussed further in Appendix G, where they are related to global symmetries. Recall from §3.6.2 that when there are obstructions to splitting $\Gamma_{ij} = \Gamma_i - \Gamma_j$ we can, in some cases, nevertheless modify the theory to incorporate the troublesome (but desirable) surface defects in a well-defined theory by gauging a suitable discrete flavor symmetry. As explained at the end of Appendix G the B-field associated to the gauging of discrete flavor symmetry allows the definition of well-defined amplitudes in the 3d sigma model when the target space has a twisted bundle with (twisted) connection.

Another useful viewpoint on the emergence of twisted bundles comes from further reduction to two-dimensional sigma models, where we can use mirror symmetry. It is well-known that reduction of four-dimensional gauge theory on two circles produces a sigma model which has a self-mirror target. In the 4d $\mathcal{N} = 2$ theories we are discussing, that target will be $\mathcal{M}$ or $\widetilde{\mathcal{M}}$. Mirror symmetry — which amounts to T-duality on the fibers over $\mathcal{B}$ — is induced by switching the compactification circles, and the theory is self-mirror because the lattice of charges Γ_g is self-dual. If we now consider 2d-4d systems we can wrap the surface defect around one compactification circle — as we have been doing — to produce a line defect in the three-dimensional sigma model, and then we can take the second reduction circle to link the defect. That is, we reduce with respect to a $U(1)$ isometry, with a fixed point on the defect. This is often described as compactification on a cigar geometry. As described in [52], the defect at the tip of the cigar requires boundary conditions which describe branes in the sigma model. As we have explained above, the resulting brane has support on all of $\mathcal{M}$ and carries (see §5.3 below) a hyperholomorphic connection. If, on the other hand, we first reduce along the linking circle and then along the transverse circle the resulting brane has fixed values of the scalar fields θ_γ, and hence is a Lagrangian brane, a section ν of $\widetilde{\mathcal{M}} \to \mathcal{B}$. (See §3.3.6.) We conclude that these branes are mirror dual to each other. [27]

Let us now consider this two-dimensional reduction in the case when the surface defect has a set of superpotentials leading to an anomaly in the splitting $\Gamma_{ij} = \Gamma_i - \Gamma_j$. Then, once again, in good cases, we can retain the surface defect by gauging a suitable discrete flavor symmetry. In this case, the Lagrangian brane is a multisection of $\mathcal{M}_d$ defined below equation (3.72), while $V_{\mathbb{S}}$ is a twisted bundle. This picture fits in very well with the work of Hausel and Thaddeus [53] who showed that for Hitchin fibrations the phenomenon of having a disconnected fiber is mirror dual to having a discrete B-field. (See also §7.2 of [5] for a discussion in the physics literature.)

---

[27] This description is incomplete. One should specify the bundle and connection on the Lagrangian brane.

5.3 Constraints from supersymmetry: hyperkähler and hyperholomorphic

The compactified theory has 8 supercharges ($\mathcal{N}=4$ supersymmetry in 3 dimensions). This imposes strong constraints both on the 3d and 1d parts of the effective action.

The constraints on the 3d part are well known: they imply that the metric g on $\mathcal{M}$ must be hyperkähler. The hyperkähler property means $(\mathcal{M}, g)$ is Kähler with respect to a family of complex structures $J^{(\zeta)}$, parameterized by $\zeta \in \mathbb{CP}^1$.

We now consider the analogous constraint on the 1d part of the action. Thompson [54] studied supersymmetric loop operators in (topologically twisted) $\mathcal{N}=4$ sigma models, with bosonic part of the form

$$\operatorname{Tr} \operatorname{Pexp} \oint \mathrm{d}x^0\, A \tag{5.7}$$

where A is a connection pulled back from some bundle V over the target $\mathcal{M}$. (The full supersymmetric expression corrects A by a fermion bilinear times the curvature.) He found a strong constraint on the connection A: the field strength F must be of type $(1,1)$ in all the complex structures $J^{(\zeta)}$ of the hyperkähler manifold $\mathcal{M}$. Such connections are said to be *hyperholomorphic.* [28]

The reason for this moniker is that we can use the $(0,1)$ part of A to define a notion of holomorphic section of V, in any complex structure $J^{(\zeta)}$: namely a holomorphic section s is one for which $(\mathrm{d}+A)s$ is of type $(1,0)$. So A equips V with the structure of a holomorphic vector bundle $V^{(\zeta)}$ over $\mathcal{M}^{(\zeta)}$, for all $\zeta \in \mathbb{CP}^1$.

5.4 The semiflat connection is hyperholomorphic in the semiflat geometry

In this section we verify that the field strength of the connection we have found in the semiflat approximation,

$$F_i^{\mathrm{sf}} = \mathrm{d}A_i^{\mathrm{sf}} = \mathrm{i}\mathrm{d}\nu_i \cdot \mathrm{d}\theta, \tag{5.8}$$

is indeed of type $(1,1)$ in all complex structures on $(\mathcal{M}, g^{\mathrm{sf}})$.

To prove this, we first quickly review the hyperkähler structure of g^{sf}. The complex structures $J^{(\zeta=0)}$ and $J^{(\zeta=\infty)}$ are rather different from the rest: in these structures the torus fibers of $\mathcal{M}^{(\zeta)} = (\mathcal{M}, J^{(\zeta)})$ are compact complex submanifolds (in fact abelian varieties). In particular, the fibers admit no non-constant holomorphic functions. In contrast, at other ζ there are plenty of holomorphic functions, which we can even write explicitly:

$$\mathcal{Y}_\gamma^{\mathrm{sf}} := \exp\left[\pi R Z_\gamma/\zeta + \mathrm{i}\theta_\gamma + \pi R \bar{Z}_\gamma \zeta\right]. \tag{5.9}$$

(Notice that $\mathcal{Y}_\gamma^{\mathrm{sf}}$ is only a *locally* defined function on $\mathcal{M}$. If we try to analytically continue $\mathcal{Y}_\gamma^{\mathrm{sf}}$ to a global function it will typically come out multivalued, because of the monodromies of Γ around the singularities of $\mathcal{B}$.) It will be useful to introduce the symbol $\mathrm{d}\log \mathcal{Y}^{\mathrm{sf}}$ for a 1-form valued in Γ^*, $\mathrm{d}\log\mathcal{Y}^{\mathrm{sf}}_\gamma = \mathrm{d}\log\mathcal{Y}^{\mathrm{sf}}\cdot\gamma$. Then the semiflat geometry is summarized by the assertion that this form is of type $(1,0)$. [29]

---

[28] Using the notation and conventions of Appendix B of [3] a simple explanation of the result can be given as follows. The integral written out more fully is of the form $\int \mathrm{d}x^0 \left(A_{ai}\dot{\varphi}^{ai} + \kappa F_{aibj}\psi_\alpha^{Ai}\psi_\beta^{Bj}\delta_A^\alpha \delta_B^\beta \epsilon^{ab}\right)$ for some constant κ. If u^a is a vector in S determining the complex structure then $F_{aibj}u^a u^b$ determines the $(2,0)$ part of the curvature in that complex structure. But this symmetric combination does not appear in the fermion bilinear term, and hence for the supersymmetric variation of $A_{ai}\dot{\varphi}^{ai}$ to cancel the fermion bilinear term the $(2,0)$ part of the curvature must vanish in all complex structures.

The relation (3.15) between ν_i and the twisted superpotential $\mathcal{W}_i$ may be rewritten in the elegant form

$$\mathrm{d}\mathcal{W}_i = \nu_i \cdot \mathrm{d}Z. \tag{5.11}$$

It follows that $\mathrm{d}\nu_i \cdot \mathrm{d}Z = 0$ and also (by complex conjugation) $\mathrm{d}\nu_i \cdot \mathrm{d}\bar{Z} = 0$. Using these facts, and linearly extending $\mathrm{d}\log\mathcal{Y}^{\mathrm{sf}}$ to be valued in $\Gamma^*\otimes\mathbb{R}$, we can write

$$F^{\mathrm{sf}}_i = \mathrm{i}\,\mathrm{d}\nu_i \cdot \mathrm{d}\log\mathcal{Y}^{\mathrm{sf}}. \tag{5.12}$$

Since $\mathrm{d}\log\mathcal{Y}^{\mathrm{sf}}$ is of type $(1,0)$ this proves that F^{sf}_i has no $(0,2)$ component in any complex structure, and, since it is pure imaginary, this implies it is $(1,1)$ in all complex structures. We conclude that A^{sf}_i is a hyperholomorphic connection on V_i.

We can actually write down holomorphic sections explicitly: in the gauge (5.4) we take

$$\mathcal{Y}^{\mathrm{sf}}_i := \exp\left[\pi R\mathcal{W}_i/\zeta + \mathrm{i}\theta_i + \pi R\bar{\mathcal{W}}_i\zeta\right]. \tag{5.13}$$

Here θ_i is a generic constant introduced for future convenience. To check that $\mathcal{Y}^{\mathrm{sf}}_i$ is a *holomorphic* section, we calculate

$$\begin{aligned}(\mathcal{Y}^{\mathrm{sf}}_i)^{-1}(\mathrm{d}+A^{\mathrm{sf}}_i)\mathcal{Y}^{\mathrm{sf}}_i &= \nu_i\cdot\left(\frac{\pi R}{\zeta}\mathrm{d}Z + \pi R\zeta\mathrm{d}\bar{Z}\right) + \mathrm{i}\nu_i\cdot\mathrm{d}\theta \\ &= \nu_i\cdot\mathrm{d}\log\mathcal{Y}^{\mathrm{sf}}\end{aligned} \tag{5.14}$$

which is manifestly of type $(1,0)$. If we take $Z_{\gamma_i^0} := \mathcal{W}_i$, then for $\gamma_i = \gamma_i^0 + \gamma$ we can define

$$\mathcal{Y}^{\mathrm{sf}}_{\gamma_i} := \mathcal{Y}^{\mathrm{sf}}_i\mathcal{Y}^{\mathrm{sf}}_\gamma = \exp\left[\frac{\pi R Z_{\gamma_i}}{\zeta} + \mathrm{i}\theta_{\gamma_i} + \pi R\bar{Z}_{\gamma_i}\zeta\right] \tag{5.15}$$

where for future convenience we set $e^{\mathrm{i}\theta_{\gamma_i}} = \sigma(\gamma_i^0,\gamma)e^{\mathrm{i}\theta_i}e^{\mathrm{i}\theta_\gamma}$. The $\mathcal{Y}^{\mathrm{sf}}_{\gamma_i}$ provide a basis of holomorphic sections.

From now on, we will define angles θ_a and sections

$$\mathcal{Y}^{\mathrm{sf}}_a = \exp\left[\pi R Z_a/\zeta + \mathrm{i}\theta_a + \pi R\bar{Z}_a\zeta\right] \tag{5.16}$$

in such a way that $\mathcal{Y}^{\mathrm{sf}}_a$ satisfy the same twisted multiplication rules as the X_a in (2.25) of §2. (Note that when a,b are not composable it does not make geometric sense to multiply $\mathcal{Y}^{\mathrm{sf}}_a$ by $\mathcal{Y}^{\mathrm{sf}}_b$, so their "product" is zero in the sense that we will never meet it.) This choice is motivated in §5.8.

---

[29] It is sometimes useful to choose a duality frame and work with an explicit basis of $(0,1)$ vector fields. One can check that the vector fields

$$\begin{gathered}\frac{\partial}{\partial a^I} + \frac{\mathrm{i}\pi R}{\zeta}\left(\frac{\partial}{\partial\theta^I_e} + \tau_{IJ}\frac{\partial}{\partial\theta_{m,J}}\right) \\ \frac{\partial}{\partial\bar{a}^I} + \mathrm{i}\pi R\zeta\left(\frac{\partial}{\partial\theta^I_e} + \bar{\tau}_{IJ}\frac{\partial}{\partial\theta_{m,J}}\right)\end{gathered} \tag{5.10}$$

indeed annihilate $\mathrm{d}\log\mathcal{Y}^{\mathrm{sf}}$.

Remark: As we have said, in the canonical framing implied by (5.4) we must make a gauge transformations on patch overlaps $\mathcal{U}_{\alpha\beta}$ so that $A_i^{\rm sf} \to A_i^{\rm sf} + {\rm i}{\rm d}\theta_{\tilde{\gamma}_{i,\alpha\beta}}$ because $\mathcal{W}_i \to \mathcal{W}_i + Z_{\tilde{\gamma}_{i,\alpha\beta}}$. From the description of the holomorphic sections in complex structure ζ which we have just given we can see that we could take the $\mathcal{Y}_i^{\rm sf}$ to be a local framing of V_i and then across patches the transition functions are holomorphic: $\mathcal{Y}_i^{\rm sf} \to \sigma(\gamma_i^0, \tilde{\gamma}_{i,\alpha\beta})\mathcal{Y}^{\rm sf}_{\tilde{\gamma}_{i,\alpha\beta}}\mathcal{Y}_i^{\rm sf}$, thus endowing V_i with the structure of a holomorphic line bundle in complex structure ζ. Now, in (5.6) we have defined V as a $\mathcal{C}^\infty$ bundle as a direct sum of line bundles. However, the hyperholomorphic connection we will construct is not diagonal with respect to this decomposition. Thus, although both V and the V_i can be endowed with the structure of holomorphic bundles in complex structure ζ, (5.6) is not an isomorphism of holomorphic bundles.

5.5 Physical interpretation of V: Expectation values

Now let us consider the expectation values of supersymmetric interfaces.

In the uncompactified IR theory, such interfaces are easy to describe. Indeed, to describe an interface in the IR between two surface defects with twisted superpotentials $\mathcal{W}_i$ and $\mathcal{W}_{j'}$, we simply divide $\mathbb{R}^{1,1}$ into two half-spaces along some line, say $x_0 = 0$, and integrate $\mathcal{W}_i$ on one half-space and $\mathcal{W}_{j'}$ on the other. By itself this breaks supersymmetry, but we can make the interface half-BPS by adding a simple boundary term to the action [17]. In other words, we insert at $x_0 = x_1 = x_2 = 0$ the line defect

$$L = L_{\gamma^0_{ij'}} = \exp \frac{1}{2} \int {\rm d}x_3 \, [(\mathcal{W}_i - \mathcal{W}_{j'})/\zeta + (\bar{\mathcal{W}}_i - \bar{\mathcal{W}}_{j'})\zeta]. \tag{5.17}$$

Here ζ is an arbitrary phase, so this is really a family of interfaces $L = L_\zeta$. (See Appendix A for further explanation of how (5.17) restores holomorphy.)

When we wrap the defect on a circle of radius R we can speak of expectation values $\langle L_{\gamma^0_{ij'}} \rangle$. In the $R \to \infty$ limit where naive dimensional reduction works, we immediately see using (5.17) that the expectation value of this IR interface is simply

$$\langle L_{\gamma^0_{ij'}} \rangle := \exp \left[\pi R \left(\mathcal{W}_i - \mathcal{W}_{j'} \right) / \zeta + \pi R \left(\bar{\mathcal{W}}_i - \bar{\mathcal{W}}_{j'} \right) \zeta \right]. \tag{5.18}$$

The expectation value of UV-defined interfaces will be a sum over sectors of vevs of such IR interfaces. Combining (5.43) with (4.19) leads then to a consistency condition on the choice of framings $e^{{\rm i}\theta_i} = e^{{\rm i}\theta_{\gamma_i^0}}$,

$$e^{{\rm i}\pi f_{ij'}} = \sigma(\gamma^0_{ij'}, \gamma^0_{j'}) e^{{\rm i}(\theta_i - \theta_{j'})}. \tag{5.19}$$

$L_{\gamma^0_{ij'}}$ is not the only possible supersymmetric interface between these two surface defects. We can get a whole family of such interfaces by bringing IR line defects with charge $\gamma \in \Gamma$ close to $L_{\gamma^0_{ij'}}$. The analog of (5.17) for the IR line defect is

$$L_\gamma := \exp \frac{1}{2} \int {\rm d}x_3 \left[\frac{Z_\gamma}{\zeta} + {\rm i}\theta \cdot \mathbb{A} + \zeta \overline{Z}_\gamma \right]. \tag{5.20}$$

Thus, the vacuum expectation values of these other interfaces are $\sigma(\gamma, \gamma_i^0 - \gamma_{j'}^0)\mathcal{Y}^{\rm sf}_{\gamma_i^0 - \gamma_{j'}^0}\mathcal{Y}^{\rm sf}_{\gamma}$. What distinguishes $L_{\gamma^0_{ij'}}$ from the other members of this class is only that $L_{\gamma^0_{ij'}}$ looks particularly simple with respect to the Lagrangian description we have chosen.

Now suppose we change our choice of Lagrangian description by shifting the superpotential $\mathcal{W}_i$ by $Z_{\tilde{\gamma}}$. In the new picture the vacuum expectation values of interfaces look slightly different. There is again a "simplest" interface L', whose vacuum expectation value is given by (5.18) with the new $\mathcal{W}_i$. The interface L' is not quite L, however; it differs by the addition of an IR line operator with charge $\tilde{\gamma}$. (One could see this, for example, by comparing the central charges of the two interfaces.) So $\langle L \rangle$ in the new picture is obtained by starting with (5.18), shifting $\mathcal{W}_i$ by $Z_{\tilde{\gamma}}$ and then dividing by $\mathcal{Y}^{\rm sf}_{\tilde{\gamma}}$. These two effects cancel each other out as far as the ζ-dependent terms go, but leave behind a ζ-independent shift: we find that if we change our Lagrangian in this way, $\langle L \rangle$ is multiplied by $e^{{\rm i}\theta_{\tilde{\gamma}}}$. Similarly, if we shift $\mathcal{W}_j$ by $Z_{\tilde{\gamma}}$ then $\langle L \rangle$ is multiplied by $e^{-{\rm i}\theta_{\tilde{\gamma}}}$. This fact can be summarized by saying that $\langle L \rangle$ is a section of the bundle $\mathrm{Hom}(V_{j'}, V_i)$.

Moreover, since each V_i has the connection $A_i^{\rm sf}$ making it into a hyperholomorphic bundle, we also get a hyperholomorphic structure on $\mathrm{Hom}(V_{j'}, V_i)$, with respect to which $\langle L \rangle$ is a holomorphic section. This holomorphy reflects the fact that L preserves the 2 supercharges (4.1).

This interpretation of $\langle L \rangle$ allows us to give a more physical understanding of the line bundles V_i and $V_{j'}$. Indeed, on general grounds $\langle L \rangle$ depends on the choice of quantum vacua for the surface defects $\mathbb{S}$ and $\mathbb{S}'$; it should really be thought of as a linear map from the vacuum ray in vacuum i to that associated to vacuum j'. This matches very well with the fact that $\langle L \rangle$ is a section of $\mathrm{Hom}(V_{j'}, V_i)$, if we identify V_i as the vacuum ray associated to the classical vacuum i on $\mathbb{S}$, and $V_{j'}$ as that associated to classical vacuum j' on $\mathbb{S}'$.

As we have mentioned above, at finite R we expect that the vector bundles of vacua $V_{\mathbb{S}}$ and $V_{\mathbb{S}'}$ still make sense, but their decomposition into individual line bundles V_i and $V_{j'}$ is not very useful. Nevertheless, the constraints from supersymmetry still operate, and imply in particular that the vev $\langle L \rangle$ of any supersymmetric line defect L should be a holomorphic section of the hyperholomorphic bundle $\mathrm{Hom}(V_{\mathbb{S}'}, V_{\mathbb{S}})$.

There is a useful observation concerning the twisted vector bundles associated to surface defects carrying flavor monodromies. Physically, interfaces are expected to exist only between surface defects which carry the same flavor monodromy. This is consistent with the idea that the vector bundles associated to these surface defects are twisted by a B-field as in §5.2.3. The bundle $\mathrm{Hom}(V_{\mathbb{S}'}, V_{\mathbb{S}})$ is a true vector bundle, with well-defined sections, only if the two surface defects have the same twisting, which in turn means they must carry the same flavor monodromy.

Finally, we remark that another physical interpretation of the connection A is that it is a Berry phase connection. This can be deduced from the considerations of Appendix A of [17].

5.6 Integral equations

In this section we introduce the main integral equations used to construct the hyperholomorphic connection A on the bundle $V_{\mathbb{S}}$. For brevity we simply denote the bundle by V. The examples of §6 are meant to demonstrate that this is a physically reasonable connection, and is indeed the exact result of the sum of the quantum corrections from 4d and 2d BPS states to the semiflat connection.

As we have mentioned, the construction of A is closely analogous to the construction in [1] of a hyperkähler metric on $\mathcal{M}$. A key step in reference [1] was the construction of holomorphic functions $\mathcal{Y}_\gamma$, corrected versions of the $\mathcal{Y}_\gamma^{\mathrm{sf}}$ which appeared above. The $\mathcal{Y}_\gamma$ were obtained as solutions of an appropriate integral equation, whose building blocks were $\mathcal{Y}_\gamma^{\mathrm{sf}}$ and the BPS data $\Omega(\gamma)$:

$$\mathcal{Y}_\gamma(\zeta) = \mathcal{Y}_\gamma^{\mathrm{sf}}(\zeta) \exp\left[\sum_{\gamma'} \Omega(\gamma')\langle\gamma',\gamma\rangle \frac{1}{4\pi \mathrm{i}} \int_{\ell_{\gamma'}} \frac{\mathrm{d}\zeta'}{\zeta'} \frac{\zeta'+\zeta}{\zeta'-\zeta} \log(1-\mathcal{Y}_{\gamma'}(\zeta'))\right]. \tag{5.21}$$

In this section we present a similar system of integral equations which determines certain sections $\mathcal{Y}_{\gamma_i}(\zeta)$ of V, likewise built from $\mathcal{Y}_{\gamma_i}^{\mathrm{sf}}$ together with the BPS data $\Omega(\gamma)$, $\mu(\gamma_{ij})$, and $\omega(\gamma,\gamma_a)$. For any fixed $\zeta \in \mathbb{C}^\times$, the $\mathcal{Y}_{\gamma_i}(\zeta)$ induce a holomorphic structure on V, in a tautological way: we take the holomorphic structure such that all $\mathcal{Y}_{\gamma_i}(\zeta)$ are holomorphic sections. Moreover, all of these holomorphic structures come from a single hyperholomorphic connection A on V. We explain the construction of A from the $\mathcal{Y}_{\gamma_i}$ in Appendix E.

The sections $\mathcal{Y}_{\gamma_i}$ are piecewise holomorphic in ζ and will undergo morphisms corresponding to those of §2.3 when ζ crosses $\mathcal{K}$-rays and $\mathcal{S}$-rays. The key to our system is to separate the Riemann-Hilbert problems associated with $\mathcal{K}$-rays and $\mathcal{S}$-rays by writing $\mathcal{Y}_{\gamma_i}$ as a product of two pieces:

$$\mathcal{Y}_{\gamma_i} = g_i x_{\gamma_i}, \tag{5.22}$$

where $x_{\gamma_i}(\zeta)$ is a section of V_i and $g_i(\zeta)$ a linear map $V_i \to V$.

The $x_{\gamma_i}(\zeta)$ are defined by the equation

$$x_{\gamma_i}(\zeta) := \mathcal{Y}_{\gamma_i}^{\mathrm{sf}}(\zeta) \exp\left[\sum_{\gamma'} \omega(\gamma',\gamma_i) \frac{1}{4\pi \mathrm{i}} \int_{\ell_{\gamma'}} \frac{\mathrm{d}\zeta'}{\zeta'} \frac{\zeta'+\zeta}{\zeta'-\zeta} \log(1-\mathcal{Y}_{\gamma'}(\zeta'))\right]. \tag{5.23}$$

For $i \neq j$, we also define sections $x_{\gamma_{ij}}(\zeta)$ of $V_j^* \otimes V_i$ by representing $\gamma_{ij} = \gamma_i - \gamma_j$ and taking

$$x_{\gamma_{ij}} := \sigma(\gamma_{ij},\gamma_j)^{-1} x_{\gamma_i} x_{\gamma_j}^{-1}. \tag{5.24}$$

Now the key integral equation for $g_k : V_k \to V$ is

$$g_k(\zeta) = g_k^{\mathrm{sf}} + \sum_{\ell \neq k, \gamma_{\ell k}} \mu(\gamma_{\ell k}) \frac{1}{4\pi \mathrm{i}} \int_{\ell_{\gamma_{\ell k}}} \frac{\mathrm{d}\zeta'}{\zeta'} \frac{\zeta'+\zeta}{\zeta'-\zeta} g_\ell(\zeta') x_{\gamma_{\ell k}}(\zeta'), \tag{5.25}$$

where g_k^{sf} is a ζ-independent and nowhere vanishing linear map $V_k \to V$. (Recalling that as a $\mathcal{C}^\infty$ bundle $V = \oplus V_i$, the obvious choice for g_k^{sf} would be just the inclusion map, but other choices could be convenient.)

In [1, 2] it was argued that the integral equation (5.21) has solutions when R is taken large enough, and that in this case the solutions have no poles or zeroes, but the situation might become more complicated for smaller R. We expect a similar story for (5.25). In what follows we mostly restrict ourselves to large R and assume the solutions do exist. In particular we assume that R is sufficiently large that no zeros develop so that g_k has no kernel. In that case $\mathcal{Y}_{\gamma_i}$ have well behaved asymptotics: precisely, $\mathcal{Y}_{\gamma_i}(\zeta)(\mathcal{Y}^{\rm sf}_{\gamma_i}(\zeta))^{-1}$ is finite in the limit $\zeta \to 0$. They are also multiplicative in the expected way:

$$\mathcal{Y}_{\gamma}\mathcal{Y}_{\gamma_i} = \sigma(\gamma,\gamma_i)\mathcal{Y}_{\gamma+\gamma_i}. \tag{5.26}$$

It is also very convenient to consider $g_i(\zeta)$ to be column vectors in the unitary framing of V described in §5.2.2. Choosing some ordering of the vacua we can then assemble these column vectors into a matrix $g(\zeta) := \oplus g_i$. We consider g to be a linear transformation $g : \oplus V_i \to V$. When ζ crosses a BPS ray $\ell_{\gamma_{ji}}$ in the counterclockwise direction we find that

$$g(\zeta) \to g(\zeta)(1 - \mu(\gamma_{ji})e_{ji}x_{\gamma_{ji}}) \tag{5.27}$$

where $e_{ji} : \oplus V_k \to \oplus V_k$ is the matrix unit. (Thus, only $e_{ji}x_{\gamma_{ji}}$ makes good geometric sense.)

Now, an important application of the above integral equations is that they allow us to construct the expectation values of general supersymmetric interfaces between $\mathbb{S}$ and $\mathbb{S}'$. Denote the corresponding vector bundles by V, V' respectively. Then we aim to construct locally defined holomorphic sections $\mathcal{Y}_{\gamma_{ij'}}$ of $\mathrm{Hom}(V', V)$. Recall that γ_i is a morphism in $\hom(i, o)$ in the vacuum groupoid $\mathbb{V}[\mathbb{S}]$ so the $\mathcal{Y}_{\gamma_i}$ are geometrically sections of V and physically are expectation values of interfaces with the vacuum i on the *left* of the interface. Similarly we can define objects $\mathcal{Y}_{-\gamma_i}$ corresponding to having the vacuum i on the right. Geometrically these are sections of V^*. The analog of (5.22) would be $\mathcal{Y}_{-\gamma_i} = g_{-i}x_{-\gamma_i}$ where $g_{-i} : V_i^* \to V^*$. However, for our purposes it is more useful to invoke the canonical isomorphism $\mathrm{Hom}(V_i^*, V^*) \simeq \mathrm{Hom}(V, V_i)$ and regard $g_{-i} : V \to V_i$ and therefore

$$\mathcal{Y}_{-\gamma_i} = x_{-\gamma_i}g_{-i} \tag{5.28}$$

as a section of V^*. The $g_{-i}(\zeta)$ are constructed from the integral equation

$$g_{-k}(\zeta) = g^{\rm sf}_{-k} - \sum_{\ell\neq k,\gamma_{k\ell}} \mu(\gamma_{k\ell})\frac{1}{4\pi \mathrm{i}}\int_{\ell_{\gamma_{k\ell}}} \frac{\mathrm{d}\zeta'}{\zeta'}\frac{\zeta'+\zeta}{\zeta'-\zeta}x_{\gamma_{k\ell}}(\zeta')g_{-\ell}(\zeta'). \tag{5.29}$$

In analogy to $g = \oplus g_i$ we can define $g_- := \oplus_k g_{-k}$, and regard g_- as a linear transformation $V \to \oplus_k V_k$. Now, as ζ crosses a BPS ray $\ell_{\gamma_{ji}}$ in the counterclockwise direction we find that

$$g_-(\zeta) \to (1 + \mu(\gamma_{ji})e_{ji}x_{\gamma_{ji}})g_-(\zeta) \tag{5.30}$$

It now follows that $gg_- : V \to V$ is free of singularities in the ζ plane, and hence must be constant. Taking the $\zeta \to 0$ limit means that it is equal to its semiflat value, which may be taken to be the identity transformation. Similarly, $g_-g : \oplus V_k \to \oplus V_k$ transforms by conjugation, but the identity matrix will satisfy the Riemann-Hilbert problem and hence from either order we conclude that if we take $g_-^{\rm sf} = (g^{\rm sf})^{-1}$ then the quantum corrections respect $g_- = g^{-1}$. In particular, if we regard g_k as column vectors then g_{-k} is the k^{th} row of the inverse matrix g^{-1}.

Finally, with these remarks in hand we are ready to define, for $i \neq j$:

$$\mathcal{Y}_{\gamma_{ij}} := g_i e_{ij} x_{\gamma_{ij}} g_{-j} \tag{5.31}$$

and, more generally, with two different surface defects we have the obvious modification giving:

$$\mathcal{Y}_{\gamma_{ij'}}(\zeta) := g_i e_{ij'} x_{\gamma_{ij'}} g_{-j'} \tag{5.32}$$

as a locally defined section of $\mathrm{Hom}(V', V)$.

We conclude this section with a number of remarks:

1. The fact that g_{-i} is the $i^{\rm th}$ row of the inverse matrix to g is not obvious from iterating the integral equations (5.25) and (5.29).

2. In general we expect the linear transformations g_i and g_{-i} to become singular on the locus $\mathcal{B}^{\rm sing}$.

3. If we define $x_{\gamma_{ii}} = \mathcal{Y}_\gamma$ then we can extend our definition (5.31) to construct $\mathcal{Y}_{\gamma_{ii}} \in \mathrm{Hom}(V, V)$, and $\sum_i \mathcal{Y}_{\gamma_{ii}} = \mathcal{Y}_\gamma 1_V$.

4. Again using the unitary framing, we can also put the x_{γ_i} together into a *diagonal* matrix x. Then the fact that $[(\mathrm{d}+A)\mathcal{Y}_{\gamma_i}]^{0,1} = 0$ (as explained in Appendix E) implies that

$$(g^{-1}Ag + g^{-1}\mathrm{d}g)^{0,1} = -(x^{-1}\mathrm{d}x)^{0,1} \tag{5.33}$$

   and hence g can be regarded as the gauge transformation which diagonalizes A to a direct sum of connections on V_i.

5. In [1] a reality condition was imposed stating that $\overline{\mathcal{Y}_\gamma(\zeta)} = \mathcal{Y}_{-\gamma}(-1/\bar\zeta)$. This is compatible with the integral equation provided $\overline{\Omega(\gamma)} = \Omega(-\gamma)$. (Of course, $\Omega(\gamma)$ is also real.) An analogous reality constraint can be imposed on the quantities defined above. Recall these are expressed in the unitary frame, allowing us to define an Hermitian structure on V_k and V. First to γ_i^0 we associate a $-\gamma_i^0$ by requiring that $Z_{-\gamma_i^0} = -Z_{\gamma_i^0}$. Then

$$\overline{x_{\gamma_i}(\zeta)} = \frac{1}{\sigma(\gamma_i, -\gamma_i)} x_{-\gamma_i}(-1/\bar\zeta) = x_{\gamma_i}^{-1}(-1/\bar\zeta) \tag{5.34}$$

   provided

$$\overline{\omega(\gamma, \gamma_i)} = \omega(-\gamma, -\gamma_i) \tag{5.35}$$

   Once again, we can also take $\omega(\gamma, \gamma_i)$ to be real. Second, using the Hermitian structure on V_k and V (defined in in §5.2) we can demand that

$$(g_k(\zeta))^\dagger = g_{-k}(-1/\bar\zeta). \tag{5.36}$$

One can check using the integral equations that (5.36) will hold provided that it is true for the semiflat value and provided that

$$\overline{\mu(\gamma_{\ell k})x_{\gamma_{\ell k}}(\zeta)} = -\mu(\gamma_{k\ell})x_{\gamma_{k\ell}}(-1/\bar{\zeta}) \tag{5.37}$$

where $\gamma_{k\ell} = -\gamma_{\ell k}$. This can be written as

$$\overline{\mu(\gamma_{\ell k})} = -\mu(\gamma_{k\ell})\sigma(\gamma_{\ell k}, \gamma_{k\ell}). \tag{5.38}$$

(In deriving this one might find it useful to note that if $\gamma_{ij} = \gamma_i - \gamma_j = -\gamma_{ji}$ then

$$\sigma(\gamma_{ij}, \gamma_j)\sigma(\gamma_{ji}, \gamma_i) = \sigma(\gamma_{ij}, \gamma_{ji}), \tag{5.39}$$

a relation which will be useful again later.)

6. Finally, regarding (5.25) in its matrix form, a special case reduces to the important integral equation used by Cecotti and Vafa in their discussion of tt^* geometry [20].

5.7 Wall-crossing formula as a consistency condition

We have made a proposal for a construction of the exact A incorporating corrections both from 2d and 4d BPS particles. One of the main points of our story is that the 2d-4d WCF is the condition implying that this A does not jump at walls of marginal stability. Let us now describe how this works.

Suppose $\mathcal{Y}_{\gamma_i}$ is constructed as above. What is its analytic structure like? From the integral equations we see that $\mathcal{Y}_{\gamma_i}(\zeta)$ depends holomorphically on ζ, *except* when ζ hits one of the integration contours ℓ_γ or $\ell_{\gamma_{ij}}$. $\mathcal{Y}_{\gamma_i}(\zeta)$ jumps when ζ crosses one of these contours. The situation is quite parallel to that of [1]. As in that case, one can determine the jump of $\mathcal{Y}_{\gamma_i}(\zeta)$ by Cauchy's theorem:

- As ζ crosses ℓ_γ, we have the discontinuity $x_{\gamma_i}(\zeta^+) = x_{\gamma_i}(\zeta^-)(1-\mathcal{Y}_\gamma)^{\omega(\gamma,\gamma_i)}$, where ζ^+ is on the counterclockwise side of ℓ_γ, which gives

$$\mathcal{Y}_{\gamma_i}(\zeta^+) = \mathcal{Y}_{\gamma_i}(1-\mathcal{Y}_\gamma)^{\omega(\gamma,\gamma_i)}(\zeta^-). \tag{5.40}$$

  This is exactly the effect of the transformation $\mathcal{K}_\gamma^{-\omega}$ from (2.30).

- As ζ crosses $\ell_{\gamma_{ji}}$, we have $g_i(\zeta^+) = g_i(\zeta^-) - \mu(\gamma_{ji})g_j x_{\gamma_{ji}}(\zeta^-)$, which gives

$$\mathcal{Y}_{\gamma_i}(\zeta^+) = \mathcal{Y}_{\gamma_i}(\zeta^-) + \mu(\gamma_{ji})\sigma(\gamma_{ji}, \gamma_i)\mathcal{Y}_{\gamma_j}(\zeta^-). \tag{5.41}$$

  This is exactly the effect of $\mathcal{S}_{\gamma_{ji}}^{-\mu}$ as given in (2.27).

These discontinuities are, fortunately, rather benign: $\mathcal{Y}$ jumps, but (as explained in Appendix E) this jump does not lead to a discontinuity of A. However, in order for A to be continuous, it is important that there should be no *other* discontinuities of $\mathcal{Y}$. Imposing this requirement leads directly to the 2d-4d WCF. The argument is basically the same as one described in [1, 3] for the 4d WCF, and so we do not repeat it here.

5.8 $\mathcal{Y}_{\gamma_i}$ and framed BPS states

So far we have constructed some holomorphic sections $\mathcal{Y}_{\gamma_i}(\zeta)$ of the holomorphic bundle $V^{(\zeta)}$ over $\mathcal{M}^{(\zeta)}$. Let us now consider what they mean physically. It follows from the discussion in §5.5 that - heuristically - we can identify $\mathcal{Y}_{\gamma_i}$ as the expectation value of an "IR line defect" with charge γ_i, which sits at the boundary of the surface defect $\mathbb{S}$. This intuitive picture is not entirely sharp, because it is hard to give a precise definition of an "IR line defect."

We can give a sharper interpretation of the $\mathcal{Y}_{\gamma_i}$ following our earlier work [3], where we addressed a similar issue for the functions $\mathcal{Y}_\gamma$. There we argued that the vev of any UV line defect L wrapped on the compactification S^1 decomposes as a sum of the $\mathcal{Y}_\gamma$:

$$\langle L\rangle = \sum_\gamma \overline{\underline{\Omega}}(L,\gamma)\mathcal{Y}_\gamma(\zeta). \tag{5.42}$$

This expansion identifies $\mathcal{Y}_\gamma$ as the contribution from a single framed BPS state of charge γ going around the compactification circle. Repeating the arguments of [3] we have a very similar story in our present 2d-4d context: letting L_ζ be a UV line defect which is an interface between $\mathbb{S}$ and $\mathbb{S}'$, we expect that there is a universal expansion

$$\langle L\rangle = \sum_{i,\gamma_i,j',\gamma_{j'}} \overline{\underline{\Omega}}(L,\gamma_{ij'})\mathcal{Y}_{\gamma_{ij'}}(\zeta). \tag{5.43}$$

(It is universal in the sense that the expansion functions $\mathcal{Y}_{\gamma_{ij'}}(\zeta)$ do not depend on L.) So we identify $\mathcal{Y}_{\gamma_{ij'}}$ similarly as the contribution from a single framed BPS state of charge $\gamma_{ij'}$ going around the compactification circle. Of course in order for this expansion to be compatible with the OPE discussed in §4 it is crucial that the $\mathcal{Y}_a$ and $\hat{X}_a$ satisfy the same twisted multiplication rules. This was the motivation for our introduction of the constant angles θ_a in (5.9). By analogy to [1, 2], we expect that the $\mathcal{Y}_{\gamma_{ij'}}$ could be characterized by a pair of requirements: first, every $\langle L\rangle$ admits a finite expansion of the form (5.43); second, the analytic continuation of $\mathcal{Y}_{\gamma_{ij'}}$ from a generic ray $\zeta \in e^{\mathrm{i}\vartheta}\mathbb{R}_+$ to the half-plane $\mathbb{H}_\vartheta$ centered on that ray behaves asymptotically like $\mathcal{Y}^{\mathrm{sf}}_{\gamma_{ij'}}$ as $\zeta \to 0$.

Note that while $\mathcal{Y}_{\gamma_{ij'}}$ and $\overline{\underline{\Omega}}(L_\zeta,\gamma_{ij'})$ are both discontinuous as functions of (u,ζ) (the former as noted in §5.7, the latter because of the framed wall-crossing) these discontinuities precisely cancel in (5.43), leaving $\langle L_\zeta\rangle$ continuous, as it should be.

6. Local models and resolution of singularities

In this section we consider two examples where the quantum corrected connection A can be described more explicitly. These examples give a concrete check of our integral equations by studying models with various simplified assumptions on the spectrum Ω, ω, μ. These simplifications typically arise when describing the light spectrum near certain local singularities in $\mathcal{B}$. These examples, which are the analog of §4 of [1], also provide illustrations of one of the important features of the story: the real-codimension-2 singularities in the connection A^{sf} are smoothed out (or at least improved) by the quantum corrections, in the sense that the corrected A only has singularities in codimension greater than 2.

Recall that in the four-dimensional theory a typical singularity arises when a 4d particle of charge γ becomes massless, i.e. $Z_\gamma(u) \to 0$ as $u \to u_0$ but with $\Omega(\gamma; u_0) \neq 0$. If — as we assume — the only particles which become massless at u_0 have charges proportional to γ then near u_0, the space $\mathcal{M}$ locally factors as a product of a smooth hyperkähler space times a four-dimensional piece, the latter fibered over a one-complex-dimensional Coulomb branch. This four-dimensional geometry, which we call periodic Taub-NUT space,[30] can be described rather explicitly by summing up a series of quantum corrections around the semiflat geometry [55, 56, 1]. These corrections smooth out the singularity of the semiflat geometry.

When generalizing this discussion to the 2d-4d case we can have $Z_\gamma(u) \to 0$ or $Z_{\gamma_{ij}}(u) \to 0$ for some $u \to u_0$. Generically we do not expect these to happen at the same point u_0, so there are two cases to consider:

1. A 2d particle of 4d charge γ becomes massless, i.e. again $Z_\gamma(u) \to 0$ as $u \to u_0$, but now with $\omega(\gamma, \cdot; u_0) \neq 0$. Nevertheless, the 2d solitons measured by $\mu(\gamma_{ij}; u_0)$ remain heavy and can be ignored in a sufficiently small neighborhood of u_0. In general there can be several particles of different charges γ becoming massless, but we will make the key simplifying assumption that they are all collinear. The case where mutually nonlocal particles become massless is significantly more difficult.

2. A 2d soliton of 4d charge γ_{ij} becomes massless, i.e. $Z_{\gamma_{ij}}(u) \to 0$ for $u \to u_0$, with $\mu(\gamma_{ij}; u_0) \neq 0$. Meanwhile all 2d particles with $\omega \neq 0$ and 4d particles with $\Omega \neq 0$ remain massive and can be ignored.

In the following sections we will analyze the local 2d-4d geometry in these two situations, and we will see how the singularity of the semiflat connection becomes partially resolved.

As we have said, we do not expect to have $Z_\gamma \to 0$ and $Z_{\gamma_{ij}} \to 0$ simultaneously for populated charges. Nevertheless, this can of course happen, and our equations (5.23) and (5.25) could be used to describe hyperholomorphic vector bundles, i.e. instantons, over the PTN geometry. The geometry of these instantons might constitute an interesting generalization of the results of Kronheimer and Nakajima. The end of §8.1.2 describes a useful local model of this situation.

6.1 Massless 4d and 2d particles on a one-dimensional Coulomb branch

6.1.1 Local system and BPS degeneracies

We will describe the local system of (gauge) charges over a one-dimensional Coulomb branch

$$D^*_\Lambda := \{a \,|\, 0 < |a| < |\Lambda|\} \tag{6.1}$$

where Λ is a UV cutoff for the effective 4d IR free theory. We let D_Λ stand for the disk with $a = 0$ restored. We can trivialize the local system after pulling back to the universal cover:

---

[30] It is also known as the Ooguri-Vafa geometry in the math literature. We will use the abbreviation "PTN space."

$$\mathbb{C}_\Lambda := \{z \,|\, \mathrm{Re}(z) < \log |\Lambda|\}, \tag{6.2}$$

and we will use the projection $p(z) = e^z = a$, so we may think of $z \sim \log a$ as a branch of the logarithm. Our local system will be

$$\Gamma := (\mathbb{C}_\Lambda \times (\mathbb{Z}\gamma_e \oplus \mathbb{Z}\gamma_m))\,/\mathbb{Z} \tag{6.3}$$

where the generator $1 \in \mathbb{Z}$ acts by taking

$$T \cdot (z; q\gamma_e + p\gamma_m) := (z + 2\pi\mathrm{i}; (q - \Delta p)\gamma_e + p\gamma_m) \tag{6.4}$$

where q, p are integers, and Δ is an integer characterizing the monodromy of the local system. The symplectic structure is given by

$$\langle \gamma_e, \gamma_m \rangle = 1. \tag{6.5}$$

The central charge function $Z \in \mathrm{Hom}(\Gamma, \mathbb{C})$ is defined by

$$Z(\gamma_e; z) = e^z = a, \tag{6.6}$$

$$Z(\gamma_m; z) = \tau_0 a + \Delta a_d, \tag{6.7}$$

where τ_0 is a constant with positive imaginary part and

$$a_d := \frac{1}{2\pi\mathrm{i}} e^z (z - (\log \Lambda + 1)) = \frac{1}{2\pi\mathrm{i}}(a \log \frac{a}{\Lambda} - a), \tag{6.8}$$

and finally we extend Z by linearity. To extend Z to the torsor Γ_i we take the low energy effective superpotential as

$$\mathcal{W} = \delta a_d + \mathcal{W}^{\mathrm{analytic}} \tag{6.9}$$

where $\mathcal{W}^{\mathrm{analytic}} = w_0 + w_1 a + w_2 a^2 + \cdots$ is some analytic superpotential. The w_i are complex numbers. The constant w_0 does not affect the geometry, so we will take $w_0 = 0$.

The period matrix is

$$\begin{aligned} \tau(a) &= \tau_0 + \frac{\Delta}{2\pi\mathrm{i}} \log \frac{a}{\Lambda} \\ &= \frac{\Delta}{2\pi\mathrm{i}} \log \frac{a}{\Lambda_4}, \end{aligned} \tag{6.10}$$

where the second equation is only valid if $\Delta \neq 0$, in which case we have $\Lambda_4 = \Lambda e^{-2\pi\mathrm{i}\tau_0/\Delta}$. In this case we can also write

$$Z(\gamma_m) = \frac{\Delta}{2\pi\mathrm{i}}(a \log \frac{a}{\Lambda_4} - a). \tag{6.11}$$

If $\Delta = 0$ then we cannot absorb τ_0 into Λ. In the remainder of §6.1 we will assume that $\Delta \neq 0$, and thus, without loss of generality, we can assume $\Lambda = \Lambda_4$. In §6.2 we will consider the special case where $\Delta = 0$. Similarly, if we take $\Lambda = \Lambda_4$ and $\delta \neq 0$ then we can write

$$\mathcal{W} = \frac{\delta}{2\pi\mathrm{i}} \left(a \log \frac{a}{\Lambda_2} - a \right) + w_2 a^2 + \cdots \tag{6.12}$$

where

$$w_1 = \frac{\delta}{2\pi \mathrm{i}} \log \frac{\Lambda_4}{\Lambda_2} \tag{6.13}$$

so we can interpret w_1 in terms of Λ_2.

We will assume that the light BPS degeneracies are of the form

$$\Omega(q\gamma_e + p\gamma_m; z) = \delta_{p,0}\Omega_q, \tag{6.14}$$

$$\omega(q\gamma_e + p\gamma_m; \gamma_0; z) = \delta_{p,0}\omega_q. \tag{6.15}$$

Here $q, p \in \mathbb{Z}$ and Ω_q and ω_q are z-independent and satisfy the parity properties:

$$\Omega_{-q} = \Omega_q \qquad\qquad \omega_{-q} = -\omega_q \tag{6.16}$$

Self-consistency of the above spectrum with the central charge and superpotential require that

1. The integer Δ is
$$\Delta = \frac{1}{2}\sum_{q\in\mathbb{Z}} q^2\Omega_q = \sum_{q>0} q^2\Omega_q, \tag{6.17}$$
because integrating out light charge q hypermultiplets with BPS degeneracy Ω_q in a theory which originally has prepotential $\tau_0 a$ results in the effective central charge Z given above.

2. Similarly, as we discussed in equation (3.68) above, integrating out light 2d chiral multiplets results in a twisted chiral superpotential as above, with
$$\delta = \frac{1}{2}\sum_{q\in\mathbb{Z}} q\omega_q = \sum_{q>0} q\omega_q. \tag{6.18}$$

3. As we have emphasized, the description of the system depends on a choice of superpotential. These are the elements denoted γ_i in §3. If we change $\gamma_i \to \gamma_i' = \gamma_i + (q_1\gamma_e + p_1\gamma_m)$, then
$$\omega_q \to \omega_q + p_1 q\Omega_q \tag{6.19}$$
and $\mathcal{W}^{\text{analytic}} \to \mathcal{W}^{\text{analytic}} + q_1 a$, or equivalently $\Lambda_2 \to \Lambda_2 e^{-2\pi \mathrm{i} q_1/\delta}$.

6.1.2 Review of the PTN geometry

The dual of the local system over the cover $\mathbb{C}_\Lambda$ is

$$\mathcal{N}_\Lambda := \mathbb{C}_\Lambda \times ((\mathbb{R}/2\pi\mathbb{Z})\gamma_e^* \oplus (\mathbb{R}/2\pi\mathbb{Z})\gamma_m^*) \tag{6.20}$$

We would like to define $\mathcal{M}^{\text{sf}} = \mathcal{N}_\Lambda/\mathbb{Z}$ where the generator of $\mathbb{Z}$ acts by

$$T \cdot (z; \theta_e\gamma_e^* + \theta_m\gamma_m^*) = (z + 2\pi\mathrm{i}; \theta_e\gamma_e^* + (\theta_m + \Delta\theta_e)\gamma_m^*). \tag{6.21}$$

This gives the Pontryagin dual local system $\Gamma^* \otimes \mathbb{R}/(2\pi\mathbb{Z})$ to Γ. However, there is a subtle complication (related to self-duality) which forces us to consider instead a local system $\widehat{\mathcal{M}}^{\mathrm{sf}} = \mathcal{N}_\Lambda/\mathbb{Z}$ where the generator of $\mathbb{Z}$ instead acts as

$$\widehat{T} \cdot (z; \theta_e \gamma_e^* + \theta_m \gamma_m^*) = (z + 2\pi \mathrm{i}; \theta_e \gamma_e^* + (\theta_m + \Delta\theta_e + \pi\Delta)\gamma_m^*). \tag{6.22}$$

The extra shift in θ_m will affect the signs of various quantities under the monodromy transformation. Of course these two actions are related by $\theta_e \to \theta_e + \pi\Delta$ so the resulting spaces are isomorphic, but the quantum corrections are a bit more natural in the coordinates appropriate to the $\widehat{T}$ quotient.

The semiflat metric is:

$$g^{\mathrm{sf}} = R\,\mathrm{Im}\tau |\mathrm{d}a|^2 + \frac{1}{4\pi^2 R\,\mathrm{Im}\tau} |\mathrm{d}\theta_m - \tau \mathrm{d}\theta_e|^2 \tag{6.23}$$

This is invariant under $\widehat{T}$ and descends to a metric on $\widehat{\mathcal{M}}^{\mathrm{sf}}$. It is positive definite for $a \in D_\Lambda^*$ provided $\Delta > 0$.

Now recall the Gibbons-Hawking ansatz for a hyperkähler four-dimensional metric on a principal circle bundle over a region of $\mathbb{R}^3$:

$$U^{-1} \left(\frac{\mathrm{d}\chi}{2\pi} + A^{\mathrm{gh}} \right)^2 + U \mathrm{d}\vec{x}^2 \tag{6.24}$$

where

$$\mathrm{d}A^{\mathrm{gh}} = \star \mathrm{d}U \tag{6.25}$$

where $\star$ is computed using the metric $\mathrm{d}\vec{x}^2 = (\mathrm{d}x^i)^2$ and orientation $\mathrm{d}x^1 \mathrm{d}x^2 \mathrm{d}x^3$. Here $\Theta^{\mathrm{gh}} = \frac{\mathrm{d}\chi}{2\pi} + A^{\mathrm{gh}}$ is a 1-form on the total space of the circle bundle, normalized so that $\pi_* \Theta^{\mathrm{gh}} = 1$.

The semiflat metric g^{sf} can be put into the form (6.24) with

$$(x_1 + \mathrm{i}x_2, x_3) = \left(a, \frac{\theta_e}{2\pi R} \right), \qquad \chi = \theta_m, \tag{6.26}$$

and

$$U^{\mathrm{sf}} = R\,\mathrm{Im}\tau, \qquad A^{\mathrm{gh,sf}} = -R\,\mathrm{Re}\tau\, \mathrm{d}x^3, \qquad F^{\mathrm{gh,sf}} = -\frac{\Delta}{(2\pi)^2} \mathrm{d}\phi\, \mathrm{d}\theta_e, \tag{6.27}$$

where $a = |a| e^{\mathrm{i}\phi}$.

Note that since θ_e is invariant under the monodromy operator $\widehat{T}$ we have a projection $\widehat{\mathcal{M}}^{\mathrm{sf}} \to D_\Lambda^* \times S^1$ defining a principal $U(1)$ bundle. The base space contracts to T^2 and the first Chern class is $-\Delta$.

The semiflat metric is hyperkähler, and we define twistor coordinates

$$\mathcal{X}_e^{\mathrm{sf}} = \exp \left[\frac{\pi R}{\zeta} a + \mathrm{i}\theta_e + \pi R \zeta \bar{a} \right], \tag{6.28}$$

$$\mathcal{X}_m^{\mathrm{sf}} = \exp \left[\frac{\pi R}{\zeta} Z(\gamma_m; z) + \mathrm{i}\theta_m + \pi R \zeta \overline{Z(\gamma_m; z)} \right] \tag{6.29}$$

with the untwisted group law $\mathcal{X}^{\rm sf}_{\gamma}\mathcal{X}^{\rm sf}_{\gamma'} = \mathcal{X}^{\rm sf}_{\gamma+\gamma'}$. (Had we worked on $\mathcal{M}$ instead of $\widehat{\mathcal{M}}$ we would have used a twisted group law.) The coordinate $\mathcal{X}^{\rm sf}_m$ is only a function on the covering space $\mathcal{N}_\Lambda$ and under $\widehat{T}$ we have:

$$\widehat{T}^* \mathcal{X}^{\rm sf}_m = (-1)^{\Delta} \mathcal{X}_e^{\Delta} \mathcal{X}^{\rm sf}_m. \tag{6.30}$$

The quantum corrected metric is again of the form (6.24) but now with

$$U = U^{\Omega} := \sum_{q=1}^{\infty} q^2 \Omega_q U_q, \tag{6.31}$$

where

$$U_q := \frac{1}{4\pi} \sum_{n=-\infty}^{\infty} \left(\frac{1}{\sqrt{q^2|a|^2 + R^{-2}(q\frac{\theta_e}{2\pi} + n)^2}} - \kappa_{n,4} \right). \tag{6.32}$$

The regularization $\kappa_{n,4}$ is chosen so that in the instanton expansion the semiflat term is $U^{\rm sf}$ given above with Λ_4. There is a corresponding gauge potential such that $\mathrm{d}A^{\Omega} = \star \mathrm{d}U^{\Omega}$. Formulae for it, in one gauge, are in [1], eqs. (4.8)-(4.10).

For any integer n we define the points $s_{n,q} := (a = 0, \theta_e = 2\pi n/q) \in D_\Lambda \times S^1$, together with the sets:

$$S_q := \{ s_{n,q} | n \in \mathbb{Z} \}. \tag{6.33}$$

Then U^{Ω} is nonsingular away from $S_{\Omega} = \cup_{q:\Omega_q \neq 0} S_q$. The metric (6.24) with (6.31) defines a metric on a principal $U(1)$ bundle over $D_\Lambda \times S^1 - S_{\Omega}$. We will not be extremely careful about specifying the different patches and trivializations of this $U(1)$ bundle and will generally denote the globally well-defined connection one-form on the total space of the bundle by $\Theta^{\Omega} = \frac{\mathrm{d}\chi}{2\pi} + A^{\Omega}$. The total space with its hyperkähler metric will be denoted $\mathcal{M}(\Omega)$ in what follows and the projection is denoted

$$\pi_{\Omega} : \mathcal{M}(\Omega) \to D_\Lambda \times S^1 - S_{\Omega} \tag{6.34}$$

If we restrict this principal bundle to a small sphere linking the point $s_{n,q}$, where (n,q) are relatively prime and $0 \le n \le q-1$, then the function U behaves like

$$U \sim \frac{N_q}{4\pi |\vec{x}|} \tag{6.35}$$

and hence the first Chern class on the linking sphere is just $c_1 = -N_q$. Here N_q is given by

$$N_q = \sum_{j=1}^{\infty} |jq| \Omega_{jq}. \tag{6.36}$$

Note that $N_q > 0$ is required for a good metric.[31]

If we attempt to extend (6.34) over the points S_{Ω} then the fiber in the fibration collapses. We can complete $\mathcal{M}(\Omega)$ to $\overline{\mathcal{M}}(\Omega)$ by adding corresponding points $p_{n,q} \in \overline{\mathcal{M}}(\Omega)$.

---

[31] In particular, massless vectormultiplets lead to singularities at finite a.

The metric near the point $p_{n,q}$ is locally a $\mathbb{C}^2/\mathbb{Z}_{N_q}$ singularity. Note that if N_q is nonzero then there is such a singularity at all the points $s_{n,q}$ for $n = 0, 1, \dots, q-1$.

Over the subspace

$$D_\Lambda^* \times S^1 \subset D_\Lambda \times S^1 - S_\Omega \tag{6.37}$$

we can identify the $U(1)$ bundles and hence compare the metrics of the PTN and semiflat spaces. If we cut out a small tube around $a = 0$ then the principal bundle of the quantum corrected space has Chern class given by the sum over the Chern classes for the points $s_{n,q}$, and this sum is just

$$c_1|_{D_\Lambda^* \times S^1} = \sum_{n,q} (-N_q) = -\Delta,$$

where the sum is over n, q such that $q > 0$, $(n, q) = 1$, and $0 \le n \le q-1$. Since the bundles are isomorphic we can choose a "common fiber coordinate $\chi = \theta_m$" for both the semiflat bundle and the full PTN space. When this is done the PTN metric and the semiflat metric g^{sf} are exponentially close for $R \to \infty$ as can be seen by rewriting U as U^{sf} plus a series of instanton corrections.

To describe the quantum-corrected twistor coordinates we should introduce the functions

$$F_q(z, \theta_e, \zeta) := \exp\left[-\frac{1}{4\pi \mathrm{i}} \int_{\ell_{q\gamma_e}} \frac{d\zeta'}{\zeta'} \frac{\zeta' + \zeta}{\zeta' - \zeta} \log[1 - \mathcal{X}_e(\zeta')^q]\right] \tag{6.38}$$

defined for any $q \in \mathbb{Z}$. Recall that

$$\ell_{q\gamma_e} := \left\{ (z, \zeta) : \frac{qa}{\zeta} < 0 \right\}. \tag{6.39}$$

The functions F_q live on the cover $\mathcal{N}_\Lambda$ (and also on the universal cover of the punctured twistor sphere). There are in fact two possible definitions of F_q. We may define a piecewise analytic function away from the BPS rays. Suppose for fixed ζ that z_0 is a value at which the BPS ray contains ζ; then

$$F_q(z_0 + \mathrm{i}\epsilon, \theta_e, \zeta) = (1 - \mathcal{X}_e^q) F_q(z_0 - \mathrm{i}\epsilon, \theta_e, \zeta) \tag{6.40}$$

The resulting function is *periodic* in z but only piecewise holomorphic as a function of z, and in fact is not defined on the BPS rays. This is the definition generally adopted in this paper. However, for our present purposes it is more convenient to take (6.38) to be the definition of F_q only in the strip

$$\arg\zeta - \pi < \mathrm{Im} z < \arg\zeta + \pi \tag{6.41}$$

for $q > 0$, and for $q < 0$ we shift the strip down by π. Then we *analytically continue* in z to define a function without discontinuities. This function satisfies

$$F_q(z + 2\pi \mathrm{i}, \theta_e, \zeta) = (1 - \mathcal{X}_e^q)^{-1} F_q(z, \theta_e, \zeta). \tag{6.42}$$

It is the latter functions, entire in z but only quasi-periodic, which we will use in our discussion of the local PTN geometry.

The formula of [1] for the quantum-corrected magnetic twistor coordinate is

$$\mathcal{X}_m = \mathcal{X}_m^{\rm sf} \prod_{q\in\mathbb{Z},q\neq 0} F_q^{q\Omega_q}. \tag{6.43}$$

Now, under $z \to z + 2\pi{\rm i}$ we have,[32] according to (6.42)

$$\prod_{q\in\mathbb{Z},q\neq 0} F_q^{q\Omega_q}(z+2\pi{\rm i}) = (-1)^{\Delta}\mathcal{X}_e^{-\Delta} \prod_{q\in\mathbb{Z},q\neq 0} F_q^{q\Omega_q}(z) \tag{6.44}$$

Thus, the functions $\mathcal{X}_m$ are invariant under the deck transformation $\widehat{T}$, and descend to $U(1)$-equivariant functions for the fundamental representation on the principal $U(1)$ fibration $\widehat{\mathcal{M}}(\Omega) \to D_\Lambda \times S^1 - S_\Omega$.

6.1.3 The semiflat line bundle

The semiflat connection is

$$A^{\rm sf} = {\rm i}(\eta\, {\rm d}\theta_e + \alpha\, {\rm d}\theta_m). \tag{6.45}$$

If $t = \eta + \alpha\tau$ with η, α real,

$$\eta = \frac{{\rm Im}\ (\bar{t}\tau)}{{\rm Im}\ \tau}, \qquad\qquad \alpha = \frac{{\rm Im}\ t}{{\rm Im}\ \tau}. \tag{6.46}$$

Defining

$$\frac{\partial \mathcal{W}^{\rm analytic}}{\partial a} := t^{\rm analytic}, \tag{6.47}$$

then

$$t = \frac{\delta}{\Delta}\tau(a) + t^{\rm analytic} \tag{6.48}$$

and hence

$$\widehat{T}^*(\eta) = \eta - \Delta\alpha + \delta, \qquad\qquad \widehat{T}^*(\alpha) = \alpha. \tag{6.49}$$

$A^{\rm sf}$ is globally defined on $\mathcal{N}_\Lambda$ but does not descend to $\mathcal{M}^{\rm sf}$. Also note that the curvature of the semiflat connection comes entirely from $\mathcal{W}^{\rm analytic}$.

We should consider $A^{\rm sf}$ to define a connection on a principal $U(1)$ bundle over $\widehat{\mathcal{M}}^{\rm sf}$ defined by

$$\widehat{P}^{\rm sf} = (\mathcal{N}_\Lambda \times U(1))/\mathbb{Z}, \tag{6.50}$$

where the generator of $\mathbb{Z}$ acts by

$$\widehat{T}\cdot(z;\theta_e\gamma_e^* + \theta_m\gamma_m^*; e^{{\rm i}\psi}) = (z+2\pi{\rm i};\theta_e\gamma_e^* + (\theta_m + \Delta\theta_e + \pi\Delta)\gamma_m^*; e^{{\rm i}\psi}e^{-{\rm i}\delta\theta_e}(-1)^{n_\omega}), \tag{6.51}$$

so that $\Theta^{\rm sf} = {\rm i}({\rm d}\psi + \eta{\rm d}\theta_e + \alpha{\rm d}\theta_m)$ is well defined on the total space of $\widehat{P}^{\rm sf}$:

$$\widehat{T}^*\Theta^{\rm sf} = \Theta^{\rm sf} \tag{6.52}$$

---

[32] In deriving this the "extra sign" comes out more naturally in the form $(-1)^{n_\Omega}$ where $n_\Omega := \sum_{q>0} q\Omega_q$. However $n_\Omega = \Delta \bmod 2$.

Here $n_\omega = \sum_{q>0} \omega_q$ defines a shift of the fiber coordinate ψ, analogous to the shift of θ_m in (6.22). Now we form the semiflat section:

$$\mathcal{X}^{\mathrm{sf}}_{\mathcal{W}} = \exp\left(\frac{\pi R}{\zeta}\mathcal{W} - \mathrm{i}\psi + \pi R \zeta \overline{\mathcal{W}}\right), \tag{6.53}$$

so that $(\mathrm{d} + \Theta^{\mathrm{sf}})\mathcal{X}^{\mathrm{sf}}_{\mathcal{W}}$ is type $(1,0)$ in all complex structures. Note that $\mathcal{X}^{\mathrm{sf}}_{\mathcal{W}}$ has the equivariance property

$$\widehat{T}^*(\mathcal{X}^{\mathrm{sf}}_{\mathcal{W}}) = (-1)^{n_\omega}\mathcal{X}^{\mathrm{sf}}_{\mathcal{W}}\mathcal{X}^{\delta}_{e}, \tag{6.54}$$

so, like $\mathcal{X}^{\mathrm{sf}}_{m}$, it is well-defined on $\mathcal{N}_\Lambda$ but does not descend to $\widehat{\mathcal{M}}^{\mathrm{sf}}$.

6.1.4 The quantum-corrected bundle V

Since we are assuming $\mu(\gamma_{ij}; a) = 0$, it follows from (5.25) that there will be no quantum mixing of line bundles and the quantum corrected connection is on the semiflat line bundle V. Using the equation (5.23) we see that the holomorphic section is given by

$$\mathcal{X}_{\mathcal{W}} = \mathcal{X}^{\mathrm{sf}}_{\mathcal{W}} \prod_{q\in\mathbb{Z}, q\neq 0} F_q^{\omega_q}, \tag{6.55}$$

which transforms under $\widehat{T}$ as

$$\widehat{T}^*\mathcal{X}_{\mathcal{W}} = \mathcal{X}_{\mathcal{W}}. \tag{6.56}$$

It thus descends to a $U(1)$-equivariant (under shifts $\psi \to \psi + \psi_0$) function from $(\mathcal{N}_\Lambda \times U(1))/\mathbb{Z}$ to $\mathbb{C}$, and hence defines a section of the associated line bundle $V \to \widehat{\mathcal{M}}^{\mathrm{sf}}$.

There is a strong formal resemblance between the function $\mathcal{X}_{\mathcal{W}}$ and the magnetic twistor function $\mathcal{X}_m$ we used to describe the PTN geometry. To bring it out, define

$$\Omega^{\mathrm{eff}}_q := \omega_q/q. \tag{6.57}$$

Then if we also set $\theta^{\mathrm{eff}}_m := -\psi$ we can write:

$$\mathcal{X}_{\mathcal{W}} = \mathcal{X}^{\omega}_m \mathcal{X}^{\mathrm{analytic}} \tag{6.58}$$

where $\mathcal{X}^{\omega}_m$ is precisely the functions $\mathcal{X}_m$ computed with Ω^{eff} (and let us stress that it is independent of θ_m, but does depend on $\theta^{\mathrm{eff}}_m = -\psi$). Moreover

$$\mathcal{X}^{\mathrm{analytic}} = \exp\left(\frac{\pi R}{\zeta}\mathcal{W}^{\mathrm{analytic}} + \pi R\zeta\overline{\mathcal{W}}^{\mathrm{analytic}}\right). \tag{6.59}$$

It follows from (6.57) that we should view $\mathcal{X}_{\mathcal{W}}$ as an equivariant function on an *effective* PTN space $\widehat{\mathcal{M}}(\omega) := \widehat{\mathcal{M}}(\Omega^{\mathrm{eff}})$. It is equivariant under translation in the circle coordinate of the fibration by π_ω analogous to (6.34). Of course, we are supposed to be defining sections of a line bundle over the space $\widehat{\mathcal{M}}(\Omega)$. Since the sections $\mathcal{X}_{\mathcal{W}}$ are independent of θ_m we can say the following. We consider the diagram:

$$\begin{tikzcd}
 & \widehat{\mathcal{M}}(\omega) \arrow[d, "\pi_\omega"] \\
\widehat{\mathcal{M}}(\Omega) \arrow[r, "\pi_\Omega"] & D_\Lambda \times S^1 - (S_\Omega \cup S_\omega)
\end{tikzcd} \tag{6.60}$$

For the moment we excise all the singular points in the base, and for simplicity we use the same notation for the total space over this smaller base. Finally, we can consider the pullback

$$\pi_\Omega^*(\widehat{\mathcal{M}}(\omega)) \to \widehat{\mathcal{M}}(\Omega) \tag{6.61}$$

which is a $U(1)$ fibration over $\widehat{\mathcal{M}}(\Omega)$. The functions $\mathcal{X}_m^\omega$ pull back to equivariant functions on the $U(1)$ fibration (6.61). These are equivalent to sections of a line bundle over $\widehat{\mathcal{M}}(\Omega)$.

6.1.5 Deriving the connection

Now we use the differential equations (4.61)-(4.64) in [1] to give an explicit construction of the hyperholomorphic connection.

We first note that since eqs. (4.61)-(4.64) of [1] are supposed to be the Cauchy-Riemann equations we can read off a basis for $T^{(0,1)}\mathcal{M}(\Omega)$ in complex structure ζ:

$$\begin{aligned} V_1^{0,1} &= \frac{\partial}{\partial a} + \frac{\mathrm{i}\pi R}{\zeta}\frac{\partial}{\partial \theta_e} - \left(\frac{\pi}{\zeta}(U^\Omega + 2\pi\mathrm{i}RA^\Omega_{\theta_e}) + 2\pi A^\Omega_a\right)\frac{\partial}{\partial \theta_m} \\ V_2^{0,1} &= \frac{\partial}{\partial \bar{a}} + \mathrm{i}\pi R\zeta\frac{\partial}{\partial \theta_e} + \left(\pi\zeta(U^\Omega - 2\pi\mathrm{i}RA^\Omega_{\theta_e}) - 2\pi A^\Omega_{\bar{a}}\right)\frac{\partial}{\partial \theta_m} \end{aligned} \tag{6.62}$$

where U^Ω denotes (6.31), and so forth. For later reference, in the semiflat limit

$$U^\Omega \to R\frac{\Delta}{4\pi}\log\left|\frac{\Lambda}{a}\right|^2 = R\,\mathrm{Im}\,\tau, \tag{6.63}$$

$$A^\Omega \to \frac{\Delta}{4\pi^2}\mathrm{Re}\,\left(\mathrm{i}\log\frac{a}{\Lambda}\right)\mathrm{d}\theta_e = -\frac{1}{2\pi}\mathrm{Re}\,(\tau)\mathrm{d}\theta_e + \mathcal{O}(e^{-R|a|}). \tag{6.64}$$

Now let Θ denote the full quantum-corrected connection. We aim to extract Θ from the equations

$$\langle V_A^{0,1}, (\mathrm{d}+\Theta)\mathcal{X}_\mathcal{W}\rangle = 0, \qquad A = 1,2. \tag{6.65}$$

First consider the case where $\mathcal{W}^{\mathrm{analytic}} = 0$ and call the corresponding connection Θ^s. In the case $\mathcal{W}^{\mathrm{analytic}} = 0$ we have $\mathcal{X}_\mathcal{W} = \mathcal{X}_m^\omega$ and hence we can use the differential equations (4.61)-(4.64) of [1] and then subtract the terms corresponding to $\langle V_A^{0,1}, \mathrm{d}\mathcal{X}_\mathcal{W}\rangle$ to get four equations for the components of Θ. We find a somewhat elegant result:

$$\Theta^s_{\theta_m} = \mathrm{i}\frac{U^\omega}{U^\Omega}, \tag{6.66}$$

$$\Theta^s_i = 2\pi\mathrm{i}\frac{(U^\omega A_i^\Omega - U^\Omega A_i^\omega)}{U^\Omega}, \tag{6.67}$$

where $i = \theta_e, a, \bar{a}$. Here U^ω means the function (6.31) computed with (6.57), and A^ω is the corresponding one-form. It is nice to check that the semiflat limit works out perfectly and that there is then a series of instanton corrections to this limit.

Now when $\mathcal{W}^{\mathrm{analytic}}$ is nonzero we can write

$$0 = \langle V_A^{0,1}, \mathcal{X}_\mathcal{W}^{-1}(\mathrm{d}+\Theta)\mathcal{X}_\mathcal{W}\rangle \tag{6.68}$$

$$= \langle V_A^{0,1}, (\mathcal{X}_m^\omega)^{-1}(\mathrm{d}+\Theta)\mathcal{X}_m^\omega\rangle + \left\langle V_A^{0,1}, \mathrm{d}\left(\frac{\pi R}{\zeta}\mathcal{W}^{\mathrm{analytic}} + \pi R\zeta\overline{\mathcal{W}}^{\mathrm{analytic}}\right)\right\rangle. \tag{6.69}$$

Let us write the total connection as

$$\Theta = \Theta^s + \Theta^{\text{analytic}}. \tag{6.70}$$

Then from (6.68) we derive

$$\Theta^{\text{analytic}}_{\theta_m} = \mathrm{i}\frac{R(\operatorname{Im}\tau)}{U^\Omega}\alpha^{\text{analytic}}, \tag{6.71}$$

$$\Theta^{\text{analytic}}_{\theta_e} = \mathrm{i}(\eta^{\text{analytic}} + (\operatorname{Re}\tau)\alpha^{\text{analytic}}) + 2\pi\mathrm{i}\frac{R(\operatorname{Im}\tau)\alpha^{\text{analytic}}}{U^\Omega}A^\Omega_{\theta_e}, \tag{6.72}$$

$$\Theta^{\text{analytic}}_{i} = 2\pi\mathrm{i}\frac{R(\operatorname{Im}\tau)\alpha^{\text{analytic}}}{U^\Omega}A^\Omega_{i}, \tag{6.73}$$

where $i = a, \bar{a}$, and we have defined $t^{\text{analytic}} := \eta^{\text{analytic}} + \tau(a)\alpha^{\text{analytic}}$. Again, the semiflat limits reproduce the contribution of $\mathcal{W}^{\text{analytic}}$ to the semiflat connection.

6.1.6 Analyzing the singularities

We consider the extension of the pullback diagram (6.60) over the singular sets S_ω and S_Ω. There are three different cases to consider with a different story for each case. Denote the integer in equation (6.36) computed for Ω^{eff}_q of (6.57) by N^{eff}_q.

1. $s_{n,q} \in S_\Omega - S_\omega \cap S_\Omega$: Above these points the metric is singular but the line bundle is not. It pulls back to a locally trivial line bundle over the neighborhood $\mathbb{C}^2/\mathbb{Z}_{N_q}$. The $\mathbb{Z}_{N_q}$-action on the line over the origin is trivial.

2. $s_{n,q} \in S_\omega - S_\omega \cap S_\Omega$: Here the line bundle is singular but the metric is not. The fiber above a point $s_{n,q}$ has $c_1 = -N^{\text{eff}}_q$. Above the point $s_{n,q}$ (in the metric fibration π_Ω) there is a whole fiber in $\widehat{\mathcal{M}}(\Omega)$. The line bundle is singular all along that ring. Thus, in this case, the singularity is only reduced from codimension two to codimension three.

3. $s_{n,q} \in S_\omega \cap S_\Omega$. Here we consider the pullback of $\mathbb{C}^2/\mathbb{Z}_{N^{\text{eff}}_q} \to D^3_*$ to a $U(1)$ fibration over $\mathbb{C}^2/\mathbb{Z}_{N_q}$ via $\pi_\Omega : \mathbb{C}^2/\mathbb{Z}_{N_q} \to D^3_*$. This is easily determined by the following simple remark. Consider the quotient Hopf fibration $\pi_{N_1} : S^3/\mathbb{Z}_{N_1} \to S^2$. This is a principal $U(1)$ bundle over S^2 with first Chern class N_1 (measured relative to a unit volume form generating $H^2(S^2;\mathbb{Z})$). On the other hand, $\pi^*_{N_1} : H^2(S^2;\mathbb{Z}) \to H^2(S^3/\mathbb{Z}_{N_1};\mathbb{Z})$ is a homomorphism $\mathbb{Z} \to \mathbb{Z}_{N_1}$ and this homomorphism is simply reduction modulo N_1. Now therefore we can consider the principal $U(1)$ fibration over $S^3/\mathbb{Z}_{N_2}$ given by $\pi^*_{N_2}(S^3/\mathbb{Z}_{N_1})$. This principal $U(1)$ bundle over $S^3/\mathbb{Z}_{N_2}$ has Chern class N_1 mod N_2. Now, there are tautological line bundles $\mathcal{R}_\rho \to \mathbb{C}^2/\mathbb{Z}_N$ labeled by ρ in the Pontryagin dual $\widehat{\mathbb{Z}}_N$. They restrict to the linking $S^3/\mathbb{Z}_N$ to be the associated bundle to the $U(1)$ principal bundle with $c_1 = \rho$. Thus in our pullback diagram (6.60), if we restrict to the neighborhood of a singular point $s_{n,q} \in S_\Omega \cap S_\omega$ we get the tautological line bundle over $\mathbb{C}^2/\mathbb{Z}_{N_q}$ given by N^{eff}_q mod N_q. Note this is invariant under the transformations $\omega_q \to \omega_q + q\Omega_q$, as it must be.

The connections we have derived are only singular on codimension three or codimension four singularities, as opposed to the semiflat connections, where the line bundle is not defined on a codimension two singularity. In this sense, the quantum corrections have smoothed out the geometry.

6.1.7 Mirror manifolds and mirror branes

We can illustrate the remarks we made above about mirror branes in the present example. The PTN space is a torus fibration $\pi : \widehat{\mathcal{M}}(\Omega) \to D_\Lambda$ with a singular fiber over $a = 0$. The fiber can be viewed as a necklace of intersecting "spheres." More precisely, the circle fibers with coordinate θ_m collapse at the points $s_{n,q} \in S_\Omega$. These fibers together with the intervals on the θ_e circle constitute a sphere with $\mathbb{Z}_{N_q}$ orbifold singularities at the north and south poles.

The section $\tilde{\nu} = (\alpha\gamma_m + \eta\gamma_e)$ mod Γ is a section of the dual fibration, which should be interpreted as the mirror manifold. Indeed, the T-dual of the semiflat metric $R\, g^{\rm sf}$ is the metric

$$\widetilde{Rg^{\rm sf}} = R^2 \ {\rm Im}\ \tau |{\rm d}a|^2 + \frac{1}{4\pi^2 {\rm Im}\ \tau} |{\rm d}\tilde{\theta}_e - \tau {\rm d}\tilde{\theta}_m|^2 \tag{6.74}$$

(where $\tilde{\theta}_i$ is T-dual to θ_i) and hence isomorphic to the original metric $Rg^{\rm sf}$, with $\tilde{\theta}_e = \theta_m$ and $\tilde{\theta}_m = \theta_e$. Hence the mirror manifold to $\widehat{\mathcal{M}}(\Omega)$ should be diffeomorphic to $\widehat{\mathcal{M}}(\Omega)$ itself. Note that it is the $\tilde{\theta}_e$ circle which shrinks at the singular points $\tilde{s}_{n,q}$ defined by $\tilde{\theta}_m = 2\pi n/q$ with $\Omega_q \neq 0$.

The section $\tilde{\nu} = (\alpha\gamma_m + \eta\gamma_e){\rm mod}\Gamma$ has coordinates $\tilde{\theta}_e = \eta$ and $\tilde{\theta}_m = \alpha$. On the other hand, it follows from (6.48) that

$$\eta = {\rm Re}\ (w_1) - {\rm Im}\ (w_1) \frac{{\rm Re}\ \tau(a)}{{\rm Im}\ \tau(a)} + \mathcal{O}(\frac{a}{\log a}), \tag{6.75}$$

$$\alpha = \frac{\delta}{\Delta} + \frac{{\rm Im}\ w_1}{{\rm Im}\ \tau(a)} + \mathcal{O}(\frac{a}{\log a}) \tag{6.76}$$

(recall that w_1 is the linear term in $\mathcal{W}^{\rm analytic}$). Hence their reductions modulo 1 have good limits at $a \to 0$. In particular, note that $\alpha \to \frac{\delta}{\Delta}$.

The section $\tilde{\nu}$ defines a Lagrangian cycle which is the support of an A-brane. Sometimes this section goes through the singular points $\tilde{s}_{n,q}$. The A-brane should be mirror to the brane described by the line bundle with hyperholomorphic connection over $\widehat{\mathcal{M}}$ we have constructed above. However, as we have seen, the latter brane depends on the details of Ω_q and ω_q whereas the support of the A-brane is only sensitive to the data w_1, δ, Δ. It must be that there is further data needed to specify this A-brane (in particular its flat connection and possible binding to fractional branes). There is undoubtedly an interesting story here, related to [57], but it lies beyond the scope of this paper.

6.2 A massless 2d particle with 4d gauge charges only

Our second example is slightly artificial, but illustrates nicely an important physical point. We return to the analysis of §6.1 but with $\Delta = 0$, so that $\tau = \tau_0$ is constant. In particular, we will assume that at some point u_0 of $\mathcal{B}$ where is a charge $\gamma \in \Gamma$ with $Z_\gamma(u_0) = 0$ and

$\omega(\gamma, \cdot; u_0) \neq 0)$ but $\Omega(\gamma; u_0) = 0$. We will also assume that $\mu(\gamma_{ij}; u_0) = 0$ so the vacua don't mix and we can take the case of a single vacuum. Note that $\omega(\gamma, \gamma_i; u_0)$ is independent of which γ_i we choose in Γ_i.

We will take the simple effective superpotential

$$\mathcal{W} = \delta \frac{\mathrm{i}}{2\pi} \left(a \log \left(\frac{a}{\Lambda_2} \right) - a \right). \tag{6.77}$$

The singularity of $\mathcal{W}$ at $a = 0$ implies a singularity for $t = \partial_a \mathcal{W}$ and hence for A^{sf} there. Note that

$$\eta = -\frac{\delta}{2\pi \mathrm{Im}\tau_0} \mathrm{Re} \left(\bar{\tau}_0 \log \frac{a}{\Lambda_2} \right), \qquad \alpha = \frac{\delta}{4\pi \mathrm{Im}\tau_0} \log \left| \frac{a}{\Lambda_2} \right|^2. \tag{6.78}$$

Now, we would like to study the IR Lagrangian obtained after compactifying the whole system on S^1. The most straightforward way to proceed is to compactify the theory including the 2d chiral multiplet. The KK mode expansion of the 2d chiral multiplet then gives an infinite set of 1d fields, each charged under the 3d gauge field. The n-th KK mode has mass $m_n = \sqrt{|qa|^2 + (n + \frac{q\theta_e}{2\pi})^2/R^2}$, where $\theta_e = \theta_{\gamma_e}$ is the Wilson line of the $U(1)$ gauge field around S^1. Thus the compactified version of (3.67) gives

$$(\mathrm{Im}\ \tau_0)\alpha_{1d} = -\sum_{q>0} q\omega_q \frac{1}{4\pi^2 R} \sum_{n=-\infty}^{\infty} \int_{-\infty}^{\infty} \mathrm{d}k \frac{1}{m_n^2 + k^2}. \tag{6.79}$$

The integral on k is elementary, but the sum requires regularization. The regularized answer can be written

$$\begin{aligned} (\mathrm{Im}\ \tau_0)\alpha_{1d} &= -\sum_{q>0} \frac{q\omega_q}{4\pi R} \sum_{n=-\infty}^{\infty} \left(\frac{1}{m_n} - \kappa_{n,2} \right) \\ &= -\sum_{q>0} \frac{q\omega_q}{4\pi R} \sum_{n=-\infty}^{\infty} \left(\frac{1}{\sqrt{|qa|^2 + (n + q\frac{\theta_e}{2\pi})^2/R^2}} - \kappa_{n,2} \right) \end{aligned} \tag{6.80}$$

where $\kappa_{n,2}$ are some regularization constants, independent of a, θ_e, chosen so that the sum converges and the leading term in the large R expansion reproduces α. Note that $\alpha_{1d} = -U^{\omega}/(R\,\mathrm{Im}\ \tau_0)$, where U^{ω} is the function U^{Ω} of equation (6.31) computed with $\Omega_q^{\mathrm{eff}} = q^{-1}\omega_q$.

The semiflat geometry is uncorrected, so we will be discussing connections on a line bundle over $D_{\Lambda}^* \times T^2$. The hyperholomorphic connection Θ has a component along θ_m determined by equation (5.1)

$$\iota_{\partial_{\theta_m}} \Theta = -\mathrm{i}\alpha_{1d}. \tag{6.81}$$

In fact, this together with the condition that A is hyperholomorphic is almost enough to determine A completely. Hyperholomorphicity in the case of a 4-dimensional $\mathcal{M}$ just means that $\mathrm{d}A$ is an anti-self-dual 2-form on $\mathcal{M}$. [33] Writing

$$\Theta = \Theta^{[3]} - \mathrm{i}\alpha_{1d} \left(\mathrm{d}\theta_m - (\mathrm{Re}\ \tau_0)\mathrm{d}\theta_e \right) \tag{6.82}$$

---

[33] We choose an orientation $e^1 \wedge e^2 \wedge e^3 \wedge e^4$ where $e^1 + \mathrm{i}e^2 = \sqrt{R\mathrm{Im}\ \tau_0} da$ and $e^3 + \mathrm{i}e^4 = \frac{\mathrm{i}}{2\pi\sqrt{R}\mathrm{Im}\ \tau_0}(d\theta_m - \tau_0 d\theta_e)$.

where $A^{[3]}$ is a $U(1)$ connection over the $\mathbb{R}^2 \times S^1$ parameterized by $(a, \bar{a}, \theta_e)$, this amounts to

$$\mathrm{d}\Theta^{[3]} = (2\pi\mathrm{i}\sqrt{R\,\mathrm{Im}\;\tau_0}) \star_3 \mathrm{d}\alpha_{1d}. \tag{6.83}$$

Here $\star_3$ denotes the 3-dimensional Hodge star with respect to the metric $(e^1)^2+(e^2)^3+(e^3)^2$ with orientation $e^1e^2e^3$. In particular the existence of an $\Theta^{[3]}$ obeying (6.83) requires that α_{1d} is a harmonic function on $\mathbb{R}^2 \times S^1$, which is indeed the case. The general $\Theta^{[3]}$ obeying this equation was again written down in §4.1 of [1] (more precisely we have $\Theta^{[3]} = -\frac{2\pi\mathrm{i}}{q}A^{\mathrm{there}}$ when ω has support at one value of q and value 1 there.)

So we have determined the form of the hyperholomorphic connection Θ. The important new point here is that it agrees with our general expression (6.66) with the replacement $U^{\Omega} \to R\mathrm{Im}\;\tau_0$. Therefore, in the general expression (6.66) we should view the instanton series for the numerator as due to instantons from worldlines of 2d BPS particles and the instanton expansion of the denominator as due to instantons from worldlines of 4d BPS particles. We can therefore interpret the result (6.66) as a combined 2d-4d instanton expansion.

The rest of the discussion of the topology of the quantum corrected line bundle proceeds as in the previous section, with the replacement of $\widehat{\mathcal{M}}(\Omega)$ by $D^*_\Lambda \times T^2$.

6.3 A massless 2d soliton

Now let us turn to the other type of singularity, where $Z_{\gamma_{ij}}(u) \to 0$ for some 2d soliton with $\mu_{\gamma_{ij}}(u_0) \neq 0$ but all occupied 4d charges have nonvanishing Z_γ at u_0.

There is a neat toy model for this situation: take a free $U(1)$ gauge theory in 4 dimensions and a surface defect supporting a single twisted chiral field X, coupled together by a twisted superpotential $\mathcal{W} = \frac{\Lambda_{\mathrm{2d}}}{3}X^3 - aX$, where a is the twisted chiral multiplet coming from the 4d theory. (So this is essentially a 2d Landau-Ginzburg model whose superpotential depends on the Coulomb branch modulus of the 4d theory.)

The 2d system has two vacua labeled by $i \in \{+,-\}$. These two vacua correspond to the two points $X = \pm\sqrt{a/\Lambda_{\mathrm{2d}}}$, with

$$\mathcal{W}_\pm = \mp\frac{2}{3\Lambda_{\mathrm{2d}}^{1/2}}a^{3/2}, \quad t_\pm = \partial_a\mathcal{W}_\pm = \mp\left(\frac{a}{\Lambda_{\mathrm{2d}}}\right)^{1/2}. \tag{6.84}$$

(For simplicity we will take Λ_{2d} positive and $\Lambda_{\mathrm{2d}}^{1/2}$ positive.) There is a single BPS soliton interpolating between the two vacua, carrying charge γ^0_{+-}, and a soliton interpolating the other way, with charge γ^0_{-+}. We adopt the sign conventions for the A_1 theories in §7.2, so $\mu(\gamma^0_{+-}) = \mu(\gamma^0_{-+}) = 1$ while $\sigma(\gamma^0_{+-}, \gamma^0_{-+}) = -1$, in accord with (5.38). There is an obvious singularity of $t_\pm$ at $a = 0$ where the soliton becomes massless, which implies a singularity of A^{sf}.

In the exact compactified theory, A^{sf} should be corrected to some smooth self-dual connection A on a rank 2 bundle V over $\mathcal{M} = \mathbb{R}^2 \times T^2$. Unlike the previous example, we will not compute this A directly: fortunately we will be able to determine it more indirectly. We first describe A in a bit more detail and then check that it indeed matches what comes from our construction.

As in the previous example, the 4d theory is just the free $U(1)$ theory, so the metric on $\mathcal{M}$ is given by (6.23). Note that the physics of the compactified system is independent of the electric and magnetic Wilson lines around S^1, as all of the UV fields are uncharged. Hence the exact corrected A is invariant under translations on the T^2 factor of $\mathcal{M}$. (Strictly speaking, the notion of "translation invariant" only makes sense once we fix a trivialization of V; here we are working with the trivialization determined by the superpotentials $\mathcal{W}_\pm$.) Dividing such an A up into components as

$$A = A^{[2]} + \varphi_a \frac{\mathrm{d}\theta_m - \bar{\tau}\mathrm{d}\theta_e}{2\pi(\mathrm{Im}\ \tau)} + \bar{\varphi}_{\bar{a}} \frac{\mathrm{d}\theta_m - \tau\mathrm{d}\theta_e}{2\pi(\mathrm{Im}\ \tau)} \tag{6.85}$$

(where $A^{[2]}$ denotes a connection along the $\mathbb{R}^2$ factor and $\bar{\varphi}_{\bar{a}}$ is the Hermitian adjoint of φ_a) the self-dual Yang-Mills equations for A become

$$\partial_{\bar{a}}\varphi_a + [A^{[2]}_{\bar{a}}, \varphi_a] = 0, \tag{6.86}$$

$$F^{[2]} + R^2[\varphi_a, \bar{\varphi}_{\bar{a}}] = 0, \tag{6.87}$$

i.e. the Hitchin equations for the pair $(A^{[2]}, \varphi_a\,\mathrm{d}a)$ on $\mathbb{R}^2 = \mathbb{C}$. In the semiflat approximation (5.4), $A^{[2]} = 0$ and φ_a is diagonal, with eigenvalues $\pm\pi t$ with $\pi t = \pi\partial_a\mathcal{W} = -\pi(\frac{a}{\Lambda_{2\mathrm{d}}})^{1/2}$. This approximation is good at sufficiently large a, up to corrections of order $e^{-2\pi R|a|}$. In particular, using Liouville's theorem these asymptotics imply

$$\mathrm{Tr}\,\varphi_a^2 = \pi^2 a/\Lambda_{2\mathrm{d}}. \tag{6.88}$$

There is a unique smooth solution $(A^{[2]}, \varphi)$ of the Hitchin equations on $\mathbb{C}$ obeying (6.88). It is radially symmetric on $\mathbb{C}$, and determined by a solution of Painleve III — see for example [58], §8.1. Fortunately, we will not need the explicit form of $(A^{[2]}, \varphi)$. Rather, we will use an indirect characterization, essentially the tt^* technology of [58, 32] (see also §9.4.1 of [2].)

Let us briefly review how that technology works in this case. We consider the flat connections

$$\nabla(\zeta) = R\zeta^{-1}\varphi + A^{[2]} + R\zeta\bar{\varphi}. \tag{6.89}$$

This family of connections can be completed to a single connection over the $\mathbb{C} \times \mathbb{CP}^1$ parameterized by (a, ζ). This connection has irregular singularities at $\zeta = 0$ and $\zeta = \infty$, exhibiting Stokes phenomena. The formal asymptotics of solutions as $\zeta \to 0$ are

$$\mathcal{Y}_\pm \sim \exp\left[\mp\frac{\pi R}{\zeta}\frac{2}{3\Lambda_{2\mathrm{d}}^{1/2}}a^{3/2}\right]e_\pm \tag{6.90}$$

where $e_\pm$ denote some ζ-independent sections. There are two anti-Stokes rays emerging from either $\zeta = 0$ or $\zeta = \infty$, located where the exponent becomes real. In the basis $(\mathcal{Y}_+, \mathcal{Y}_-)$ the corresponding Stokes factors are known to be simply

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}. \tag{6.91}$$

See Appendix B for more details. Moreover, these formal asymptotics and Stokes data are enough to *determine* the full connection, hence the solution $(A^{[2]}, \varphi)$ of Hitchin's equations, and hence finally the hyperholomorphic connection A.

Now let us see how this description is reproduced by our construction. In our present setup, where $\omega(\cdot,\cdot) = 0$, (5.22), (5.23) just say

$$\mathcal{Y}_{\gamma_\pm} = g_\pm \mathcal{Y}^{\mathrm{sf}}_{\gamma_\pm}, \tag{6.92}$$

while $\mu(\gamma^0_{+-}) = 1$, $\mu(\gamma^0_{-+}) = 1$, with all other $\mu(\cdot) = 0$, so (5.25) says

$$g_+(\zeta) = g_+^{\mathrm{sf}} + \frac{1}{4\pi\mathrm{i}} \int_{\ell_{\gamma^0_{-+}}} \frac{\mathrm{d}\zeta'}{\zeta'} \frac{\zeta' + \zeta}{\zeta' - \zeta} g_-(\zeta') \mathcal{Y}^{\mathrm{sf}}_{\gamma^0_{-+}}(\zeta'), \tag{6.93}$$

$$g_-(\zeta) = g_-^{\mathrm{sf}} + \frac{1}{4\pi\mathrm{i}} \int_{\ell_{\gamma^0_{+-}}} \frac{\mathrm{d}\zeta'}{\zeta'} \frac{\zeta' + \zeta}{\zeta' - \zeta} g_+(\zeta') \mathcal{Y}^{\mathrm{sf}}_{\gamma^0_{+-}}(\zeta'). \tag{6.94}$$

(recall that $\mathcal{Y}^{\mathrm{sf}}_{\gamma^0_i}$ is defined in (5.13), (5.15) but $\mathcal{Y}^{\mathrm{sf}}_{\gamma^0_{ij}}$ is defined by (5.24)). In particular, $\mathcal{Y}_{\gamma^0_\pm}$ are independent of base coordinates θ_e, θ_m on T^2, i.e. they are invariant under translations along T^2. We may thus consider them as sections of a bundle over the base $\mathbb{C}$. As just explained, the translation invariance implies this bundle supports a solution $(A^{[2]}, \varphi)$ of the Hitchin equations. The Cauchy-Riemann equations for holomorphic sections of $V^{(\zeta)}$, when restricted to translation invariant sections, reduce to flatness under $\nabla(\zeta)$. So $\mathcal{Y}_{\gamma^0_\pm}$ are $\nabla(\zeta)$-flat. (6.92)-(6.94) show that $\mathcal{Y}_{\gamma^0_\pm}$ have the asymptotics (6.90). Finally, we can read off the Stokes factors from the discontinuities of $\mathcal{Y}_{\gamma^0_\pm}$ across the two rays $\ell_{\gamma^0_{+-}}$ and $\ell_{\gamma^0_{-+}}$. These discontinuities are determined by (6.93) and (6.94). The net result is the following:

1. Across the wall $a^{3/2}/\zeta \in \mathbb{R}_-$ we have

$$\begin{aligned} \mathcal{Y}_+(+) &= \mathcal{Y}_+(-) + \sigma(\gamma_{-+}, \gamma_+)\mathcal{Y}_- \\ \mathcal{Y}_-(+) &= \mathcal{Y}_-(-) = \mathcal{Y}_- \end{aligned} \tag{6.95}$$

   where $\mathcal{Y}_\gamma(\pm)$ refers to the side of the wall with $\mathrm{Im}(\pm a^{3/2}/\zeta) > 0$.

2. Across the wall $a^{3/2}/\zeta \in \mathbb{R}_+$ we have

$$\begin{aligned} \mathcal{Y}_+(+) &= \mathcal{Y}_+(-) = \mathcal{Y}_+ \\ \mathcal{Y}_-(+) &= \mathcal{Y}_-(-) + \sigma(\gamma_{+-}, \gamma_-)\mathcal{Y}_+ \end{aligned} \tag{6.96}$$

   where $\mathcal{Y}_\gamma(\pm)$ refers to the side of the wall with $\mathrm{Im}(\mp a^{3/2}/\zeta) > 0$.

Using (5.39) we know that $\sigma(\gamma_{+-}, \gamma_-)\sigma(\gamma_{-+}, \gamma_+) = -1$. With an appropriate choice of σ subject to this constraint, (6.95), (6.96) correspond to the desired Stokes factors (6.91). This is enough to show that our construction produces the A we described above.

In particular, our construction has produced a *smooth* hyperholomorphic connection: this is an example of the advertised resolution of singularities induced by 2d BPS states.

7. Example: A type theories and Hitchin systems

One important example of our construction is its application to the "canonical surface defects" in theories of class $\mathcal{S}$.

Recall that theories of class $\mathcal{S}$ are obtained from an ADE group and a curve C (with punctures) as described in [43, 2, 44]. For simplicity, throughout this section we will restrict our attention to the case where the ADE group is of type A_{n-1}. Then there are canonical surface defects $\mathbb{S}_z$ corresponding to points $z \in C$ [59, 44]. (If we think of the theory as obtained from a decoupling limit of M5-branes, then $\mathbb{S}_z$ can be constructed from an M2-brane ending on the M5-branes at the point z.) In this class of theories and surface defects, we can describe the 2d-4d wall-crossing data of §§2.3 and 3 explicitly in terms of the geometry of C and its spectral cover.

We will need a bit of care in defining the canonical surface defect, due to some subtleties about locality of surface defects in the 6d theory. It is not difficult to argue (looking for example at a KK reduction of the theory to 5d Yang-Mills or to the Abelian theory on the Coulomb branch of the 6d theory) that the basic surface defects in 6d have a potential mutual locality problem. If we keep a surface defect fixed, look at the S^3 which surrounds it, nucleate a second surface defect near the north pole of the sphere, and annihilate it near the south (in analogy with what is done in 4d with Wilson and 'tHooft operators) the partition function gains a factor of $e^{2\pi \mathrm{i}/n}$. [34] This factor plays an important role when one looks at mutual locality of line defects in 4d corresponding to paths on C. There is a second application, though, to mutual locality between surface defects sitting at a point in C and point operators in 4d engineered by wrapping surface defects on C: the point operator has $e^{2\pi \mathrm{i}/n}$ monodromy around the surface defect. The local operator which descends from a surface defect wrapping C was identified in [60] with a specific chiral operator which receives vevs on the Higgs branch. Hence in order to include the canonical surface defect $\mathbb{S}_z$ in the 4d theory, strictly speaking one needs to gauge the $\mathbb{Z}_n$ flavor symmetry corresponding to the monodromy of these specific chiral operators around the canonical surface defect. At least in the A_1 case with regular singularities we will, in §7.4, identify the $\mathbb{Z}_2$ flavor symmetry in the four dimensional Lagrangian description of the theories, and verify that the Γ_i fail to be Γ torsors in a way which is exactly controlled by the $\mathbb{Z}_2$ flavor monodromy, leading to the kind of anomaly cancellation discussed in §3.6.2. The generalization to higher rank should not be hard, though it might be hampered by the lack of a Lagrangian description.

The fact that $\mathbb{S}_z$ carries flavor monodromies nicely resolves some puzzles rased in [59] (in a way anticipated in [59]). Whenever C approaches a factorization limit and develops a long thin tube, a weakly coupled gauge group appears in the four dimensional field theory.

---

[34] One way to see this is to work on the Coulomb branch. Then the worldsheet of the nucleated surface defect has a topologically interesting term $\exp[2\pi \mathrm{i} \int_\Sigma (v_1, B)]$ where B is a self-dual two-form valued in the Cartan subalgebra of A_{n-1}. Here v_1 is a vector in the weight lattice of A_{n-1}. The other surface defect, which we locate at the center of $\mathbb{R}^4$ surrounded by the linking sphere S^3 is a magnetic source of B of strength $H = v_2 \omega_3$ where ω_3 is the unit normalized volume form on S^3 and v_2 is in the weight lattice of A_{n-1}. Here we have identitified the Cartan subalgebra with its dual using a metric so that roots have length-squared two. Then the topologically interesting phase for the process described above is $\exp[2\pi \mathrm{i}(v_1, v_2)]$, and since both v_1, v_2 are in the weight lattice of A_{n-1} this is an n^{th} root of unity.

If the point z is in the tube region, then, as was observed in [59], the surface defect $\mathbb{S}_z$ appears to be well described by a GW surface defect. However, a direct identification with a GW surface defect leads to two puzzles: First, the GW surface defect alone does not break any flavor symmetry, but the canonical surface defect often appears to break some flavor symmetry. The resolution of this puzzle is that the appropriate GW defect must carry some flavor monodromy and this monodromy does indeed break the flavor group down to the commutant in the flavor group of the gauged $\mathbb{Z}_n$ subgroup. The second problem is that several class $\mathcal{S}$ descriptions of the same 4d theory might exist, and in general these will have sharply different canonical surface defects. How can they all correspond to a single GW surface defect? The resolution is that there are several GW defects, distinguished by having different flavor monodromies.

7.1 Review of pure 4d data

First we recall from [2] some purely 4d aspects of the story. Although the theory exists for any simply laced Lie algebra we will for simplicity focus on the case of A_{n-1}. Later we will specialize to $n = 2$.

A point u of the Coulomb branch $\mathcal{B}$ corresponds to a set of meromorphic k-differentials ϕ_k $(k = 2, 3, \dots, n)$ on C. The ϕ_k have poles only at the punctures of C, and part of the data defining the theory is a set of linear conditions on the singular parts of the ϕ_k at the punctures; roughly these conditions identify the residues of the ϕ_k with various combinations of the mass parameters of the theory. $\mathcal{B}$ is the space of all tuples $(\phi_2, \phi_3, \dots, \phi_n)$ of meromorphic differentials obeying these linear conditions. $\mathcal{B}$ is thus a finite-dimensional affine space modeled on $\oplus_{d=2}^{n} H^0(C, K^{\otimes d})$.

For every $u \in \mathcal{B}$, there is a corresponding Seiberg-Witten curve,

$$\Sigma_u = \{\phi_n + \phi_{n-1}\lambda + \cdots + \phi_2\lambda^{n-2} + \lambda^n = 0\} \subset T^*C. \tag{7.1}$$

λ is a cotangent vector, so that each term in the above equation is an n-differential on C. At a generic point of C the equation has n distinct solutions, so $\Sigma_u \subset T^*C$ is an n-fold covering of C. Locally the n solutions give n 1-forms λ_i on C. These 1-forms pull back to a single globally defined 1-form on Σ_u which we also call λ.

The local system of lattices Γ begins with a certain subquotient of $H_1(\Sigma_u; \mathbb{Z})$ and then extends it so that Γ_g is self-dual. This extension depends on a choice of a set $\mathcal{L}$ of maximal mutually local line defects and determines a precise global structure of the gauge group. In the A_1 case we begin with the sublattice $H_1(\Sigma_u; \mathbb{Z})^-$ odd under the deck transformation and then, depending on the choice of $\mathcal{L}$, we extend it to a self-dual lattice. (For details see [3].) The central charge function Z_γ is

$$Z_\gamma = \frac{1}{\pi} \oint_\gamma \lambda. \tag{7.2}$$

Finally we come to the most interesting part of the story: the $\Omega(\gamma; u)$ are the 4d BPS degeneracies, described in [45, 2]. They count certain special networks on C, defined as follows: First, define a *WKB curve of type ij and phase* ϑ to be an oriented real curve on

C, along which $e^{-\mathrm{i}\vartheta}\langle(\lambda_i - \lambda_j), \partial_t\rangle$ is real and positive for some pair of vacua i, j. Here ∂_t is a positively oriented tangent vector along the path. [35] A *WKB network of phase* ϑ is by definition a network of curves on C whose legs are WKB curves of phase ϑ. The network can have three-legged junctions where legs of types ij, jk, ki come together. A leg of type ij is also allowed to end on a branch point where $\lambda_i - \lambda_j = 0$, i.e. the i-th and j-th sheets of Σ merge.

Following any leg of a WKB network in either direction, there are four possible outcomes: either we reach a junction, we reach a branch point, we return to where we started, or we spiral into one of the punctures of C. We call the network *finite* if this fourth possibility never occurs. This last condition makes finite WKB networks rather special, since a generic WKB curve spirals into punctures in both directions. See Figure 13.



Figure 13: A possible finite WKB network in the A_3 theory. All legs are WKB ij-curves, with ij as indicated. Similarly the branch points (orange crosses) are ij-branch points as indicated.

Each finite WKB network has a canonical lift to a union of closed curves on Σ, obtained by lifting a leg of type ij to a pair of curves on sheets i and j of Σ, with opposite orientations. The $\Omega(\gamma; u)$ count finite WKB networks with phase $\vartheta = \arg Z_\gamma$, for which the homology class of the lift is γ.[36] Isolated finite WKB networks contribute $+1$ to Ω. The contributions from networks with moduli are more complicated and have not been analyzed systematically, except in the case of the A_1 theory (see below).

7.2 2d-4d data

Now we describe the new data attached to the surface defect. Most of what we say could be guessed just by looking for the most obvious generalizations of the 4d data to incorporate the surface defect, consistent with the general structure we have described in the rest of this paper. A fuller justification of our claims here could be given using a string theory realization of the A_{n-1} theory. Alternatively, in the A_1 case, our claims are justified by the WKB analysis which we sketch later in this section.

---

[35] In [2] WKB curves for the A_1 case were defined to be unoriented. This is compatible with the present definition, because we only require orientation with respect to *some* pair of vacua i, j.

[36] This is actually equivalent to counting finite WKB networks with arbitrary phase — indeed, if a finite WKB network has phase ϑ and lifts to a class γ, then $\vartheta = \arg Z_\gamma$.

1. **Vacua**: The finite set $\mathcal{V}(\mathbb{S}_z)$ of vacua of the surface defect is just the set of preimages of z in Σ_u; call them x_i, $i = 1, \ldots, n$. Often we will simply write i for the vacuum x_i. (This follows from the identification [17] between the twisted chiral ring of the 2d theory on the surface defect and the Seiberg-Witten curve of the 4d theory.)

2. **Soliton charge torsors**: The torsors Γ_{ij} are taken to be the Γ-torsor of (relative) homology classes of open paths from x_i to x_j. (Thus, we choose one connected path from x_i to x_j and then consider all translates of the relative homology class in $H_1(\Sigma_u, \{x_i, x_j\}; \mathbb{Z})$ by Γ.) The central charge function $Z_{\gamma_{ij}}$ is then [37]

$$Z_{\gamma_{ij}} = \frac{1}{\pi} \int_{\gamma_{ij}} \lambda. \tag{7.3}$$

3. **2d-4d BPS degeneracies** ω: The most naive guess for $\omega(\gamma, \gamma_{ij})$ would be

$$\omega(\gamma, \gamma_{ij}; u) = \Omega(\gamma; u) \langle \gamma, \gamma_{ij} \rangle. \tag{7.4}$$

   This equation is indeed correct, but it needs some clarification: choosing different representatives for the class γ would *a priori* give different values for the intersection number $\langle \gamma, \gamma_{ij} \rangle$, since representatives of γ_{ij} are open curves. Which representative should we use?

   First consider the special case where the only finite WKB networks in class γ are isolated. In this case, the lift of each such network defines a canonical representative of the class γ. In particular this representative defines a definite class $\tilde{\gamma}$ in the homology group $H_1(\Sigma_u - \{x_i, x_j\}; \mathbb{Z})$ (which pairs with $H_1(\Sigma_u, \{x_i, x_j\}; \mathbb{Z})$). Having made this definition, we can say that this lifted network contributes $\langle \tilde{\gamma}, \gamma_{ij} \rangle$ to $\omega(\gamma, \gamma_{ij})$. To get the full $\omega(\gamma, \gamma_{ij})$ we sum over the networks contributing to $\Omega(\gamma)$.

   If the networks contributing to $\Omega(\gamma)$ are not isolated, then the job of computing $\omega(\gamma, \gamma_{ij})$ is more complicated. We do not give a general prescription here, but we do indicate what should be done in the first nontrivial example, the A_1 theory. In this case the networks we have been discussing in fact just consist of single WKB curves — there are no nontrivial junctions. The new case we need to discuss is the case where the WKB curve is closed: in this case it is not isolated but rather lies in a 1-parameter family, sweeping out an annulus on C. In this case, we focus our attention on the two closed WKB curves which make up the inner and outer boundaries of the annulus: letting $\tilde{\gamma}, \tilde{\gamma}'$ denote the lifts of these two boundaries, we define

$$\omega(\gamma, \gamma_{ij}) = -\langle \tilde{\gamma}, \gamma_{ij} \rangle - \langle \tilde{\gamma}', \gamma_{ij} \rangle. \tag{7.5}$$

   In the case where z is not in the annulus both terms contribute equally and this simplifies $\omega(\gamma, \gamma_{ij}) = -2\langle \tilde{\gamma}, \gamma_{ij} \rangle$. This is our original naive formula (7.4), since in this case $\Omega(\gamma) = -2$.

---

[37] In Seiberg-Witten theory it is often said that the Seiberg-Witten differential is ambiguous by shifts $\lambda \to \lambda + \mathrm{d}f$. This would affect the central charges here. In the theories of class $\mathcal{S}$ there is a distinguished Seiberg-Witten differential. This is the one which appears in the chiral ring, and is the one we should employ here.

4. **2d-4d BPS degeneracies** μ: Finally, the $\mu(\gamma_{ij}; u)$ count *finite open WKB networks of type* ij on C. An open WKB network is just like the WKB networks we defined above, except that it has one special leg which is of type ij and has one end at the point z. The lift of any finite open WKB network to Σ defines an element $\gamma_{ij} \in \Gamma_{ij}$. When the moduli space of these networks is an isolated set of points, $\mu(\gamma_{ij})$ is a signed sum over these points. In general we do not have a prescription to fix the sign. In the A_1 case we do know the answer: in this case there is at most a single finite open WKB network in each class, and it contributes $\mu(\gamma_{ij}) = +1$.

5. **Twisting function** σ: We do not know the tricky ±1-valued twisting function σ in general. In the A_1 case we do know it: it is

$$\sigma(\gamma, \gamma') = (-1)^{\langle \gamma, \gamma' \rangle}, \tag{7.6}$$

$$\sigma(\gamma, \gamma_{ii}) = (-1)^{\langle \gamma, \gamma_{ii} \rangle}, \tag{7.7}$$

$$\sigma(\gamma, \gamma_{ij}) = \sigma(\gamma), \tag{7.8}$$

$$\sigma(\gamma_{ij}, \gamma_{ji}) = -\sigma(\gamma_{ij} + \gamma_{ji}), \tag{7.9}$$

   where the σ appearing on the *right* side is the "canonical quadratic refinement" $\sigma : \Gamma \to \mathbb{Z}_2$ which appeared in §7.7 of [2]. In particular recall that $\sigma(\gamma) = -1$ if γ supports a hypermultiplet and $= +1$ if γ supports a vectormultiplet. The above sign prescription is justified in Appendix F.

   We will sometimes use the signs $\sigma(\gamma_{ij}, \gamma_j)$ as well; these involve gauge choices which we do not discuss here.

6. **Supersymmetric interfaces**: In order to extend the wall-crossing data to include interfaces we must describe a suitable class of such interfaces. Suppose we pick two points z and z' of C. We then have two surface defects $\mathbb{S} = \mathbb{S}_z$ and $\mathbb{S}' = \mathbb{S}_{z'}$ Interfaces L between $\mathbb{S}$ and $\mathbb{S}'$ correspond to paths $\wp$ in C from z to z'. One can justify this by considering Janus configurations in the field theory, where z is regarded as a parameter of the surface defect. Alternatively, one can take a geometric engineering viewpoint and construct the interface between $\mathbb{S}_z$ and $\mathbb{S}_{z'}$ by considering three open M2 branes. The first ends on the surface $x^1 = x^2 = 0$ with $x^0 \leq 0$ at $z \in C$. The third ends on the surface $x^1 = x^2 = 0$ with $x^0 \geq 0$ at $z' \in C$, and the second ends at $x^1 = x^2 = x^0 = 0$ in $\mathbb{R}^{1,3}$ times a path $\wp$ on C from z to z'. In a discussion analogous to that of [3], §7.4, the partial twisting of the theory along C means that the defect only depends on the homotopy class of $\wp$ (with fixed endpoints). [38] The Hilbert space $\mathcal{H}_{\mathbb{S}_z L_{\zeta,\wp} \mathbb{S}_{z'}}$ will be graded by a charge lattice $\Gamma_{ij'}$, which will consist of a Γ torsor of homology classes of open paths from x_i (above z) to $x_{j'}$ (above z'). Once again we take

$$Z_{\gamma_{ij'}} = \frac{1}{\pi} \int_{\gamma_{ij'}} \lambda. \tag{7.10}$$

---

[38] One can also introduce supersymmetric interfaces corresponding to the "laminations" discussed in [3], but we will leave that for another occasion.

7. **Framed BPS States**: These should be given by some analog of the millipede construction of §10.8 of [3], but we have not developed the details here.

Let us give one simple example of how the above geometric prescription is compatible with the general wall-crossing formulae of §2.4. In discussing wall-crossing we can vary both u and z, the basepoint on C of the projection of the open curves γ_{ij} on Σ. When we vary z we sometimes write $\gamma_{ij}(z)$ to emphasize the basepoint. When we vary z, it can cross finite WKB networks, and this can change $\langle\tilde{\gamma},\gamma_{ij}\rangle$ thus inducing wall-crossing in $\omega(\gamma,\gamma_{ij})$. To illustrate this, consider a family of basepoints z following a path which crosses a finite WKB curve connecting two branch points of type (ij). This finite curve has lift $\tilde{\gamma}$ with homology class γ. By our rules $\Omega(\gamma)=1$ and $\sigma(\gamma)=-1$. Suppose that the finite WKB curve has phase ϑ_*. When $z=z_*$ is a point on the image of $\tilde{\gamma}$, that is, when it is a point on the finite WKB network, there are two open finite WKB curves $\gamma_{ij}(z_*)$ and $\gamma_{ji}(z_*)$, such that $\gamma_{ij}(z_*)+\gamma_{ji}(z_*)=\tilde{\gamma}$. Now let $z_\pm$ be displaced above and below $\tilde{\gamma}$ as in Figure 14. By continuity, the curves $\gamma_{ij}(z_\pm)$ and $\gamma_{ji}(z_\pm)$ continue to support finite open WKB curves with phase slightly displaced from ϑ_*. By our rules $\mu(\gamma_{ij})=\mu(\gamma_{ji})=1$ and $\sigma(\gamma_{ij},\gamma_{ji})=1$, where γ_{ij} applies to any of the three values of $z=z_\pm,z_*$. At $z=z_*$ the three central charges $Z_{\gamma_{ij}(z_*)}$, $Z_{\gamma_{ji}(z_*)}$, and Z_γ are aligned, so as z passes through the finite WKB network the BPS rays of $\ell_{\gamma_{ij}},\ell_\gamma,\ell_{\gamma_{ji}}$ change order, exactly as in the wall-crossing formula (2.44). Plugging $\Sigma_{ij}=\Sigma_{ji}=\Sigma'_{ij}=\Sigma'_{ji}=1$ into (2.45) we find that the wall-crossing formula demands that



Figure 14: z crossing a finite WKB curve with lift $\tilde{\gamma}$. When $z=z_+$, the intersection $\langle\tilde{\gamma},\gamma_{ij}(z_+)\rangle=+1$ (left). When $z=z_-$, $\langle\tilde{\gamma},\gamma_{ij}(z_-)\rangle=-1$ (right). This leads to a change of $\omega(\gamma,\gamma_{ij})$ as we pass from $z=z_+$ to $z=z_-$; this change is consistent with the 2d-4d wall-crossing formula, as explained in the text.

$$\begin{aligned}\Pi'_i&=\Pi_j\\ \Pi'_j&=\Pi_i\\ \Pi_i&=(1-X_\gamma)\Pi_j\end{aligned}\tag{7.11}$$

and hence, if primed quantities refer to the z_+ side of the wall,

$$\begin{aligned}\omega(\gamma,\gamma_{ij}(z_+))&=1\\ \omega(\gamma,\gamma_{ij}(z_-))&=-1\\ \omega(m\gamma,\gamma_{ij}(z_\pm))&=0\qquad\qquad m>1.\end{aligned}\tag{7.12}$$

These intersection numbers can be verified directly in Figure 14.

One important general lesson we learn from the above example is that *the walls of marginal stability for the parameter z of $\mathbb{S}_z$ include the finite and separating WKB curves* *with phase $\vartheta = \vartheta_*$, where ϑ_* is the phase of some 4d BPS state.* The reason is that, if z lies on a finite or separating WKB curve of phase ϑ_*, there is at least one finite open network with phase ϑ_* (indeed this network just consists of a single finite open WKB curve from z to a turning point). Thus we have at least two BPS states with the same phase (one 2d, one 4d), which means z is on a wall of marginal stability. We will sometimes refer to the finite and separating WKB curves at these special values ϑ_* as *critical WKB curves.*

The details of what happens at the wall depend on whether z is crossing a finite or a separating WKB curve. We showed above how ω can change as z crosses a finite WKB network, while μ is invariant. On the other hand, as we will see in §8, when z crosses a separating WKB curve μ can change, while ω is invariant. This corresponds to the identity (2.36) of type $\prod \mathcal{S} \cdot \mathcal{K} = \mathcal{K} \cdot \prod \mathcal{S}$.

7.3 3d compactification and Hitchin's equations

Now, we consider the theory compactified on S^1 to 3 dimensions.

The moduli space $\mathcal{M}$ of the theory on $\mathbb{R}^{2,1} \times S^1_R$ is the moduli space of solutions of Hitchin's equations [2]. Recall [61] that these are equations for a connection A on a rank n complex vector bundle $E \to C$ and a "Higgs field" $\varphi \in \Omega^{1,0}(\mathrm{End}\ E)$:

$$\begin{aligned} F + R^2[\varphi, \bar{\varphi}] &= 0, \\ \bar{\partial}_A \varphi &= 0, \\ \partial_A \bar{\varphi} &= 0. \end{aligned} \tag{7.13}$$

Here the gauge group is $SU(n)$.[39] We impose the boundary conditions $(A, \varphi) = (A_0, \varphi_0) +$ (regular) near the punctures z_a of C, where (A_0, φ_0) is a suitable singular solution near z_a.[40] These equations can be interpreted as the moment maps for a hyperkähler quotient construction of $\mathcal{M}$ as $\mathcal{M} = \mathcal{N}///\mathcal{G}$. Here $\mathcal{N}$ is the space of (A, φ), not necessarily satisfying (7.13), but satisfying the boundary conditions. $\mathcal{G}$ is the group of gauge transformations which preserve the singular solutions (A_0, φ_0) near each point z_a. Thus, at each puncture z_a there is generically a reduction of structure group from $SU(n)$ to the stabilizer $K_a \subset SU(n)$ of (A_0, φ_0).

The map $\mathcal{M} \to \mathcal{B}$ is the "Hitchin fibration,"

$$\phi_k = \mathrm{Tr}\, \varphi^k. \tag{7.14}$$

Each solution $(A, \varphi) \in \mathcal{M}$ induces a family of flat connections parameterized by $\zeta \in \mathbb{C}^\times$,

$$\mathcal{A} = R\frac{\varphi}{\zeta} + A + R\zeta\bar{\varphi}. \tag{7.15}$$

Indeed Hitchin's equations can be interpreted as the statement that $\mathcal{A}$ is flat for all ζ. For any fixed $\zeta \in \mathbb{C}^\times$, (7.15) gives an identification between $\mathcal{M}^\zeta$ and a moduli space of flat $SL(n, \mathbb{C})$ connections over C, again with appropriate singularities at the punctures.

---

[39] More precisely $\mathcal{M}$ is a certain finite quotient of the moduli space of solutions with gauge group $SU(n)$. The precise quotient we pick depends on some fine details of the 4d theory, like whether we take the gauge group to be simply connected or not; see §§6.3 and 7.3 of ref. [3] for some more discussion of this point.

[40] See §4.1 of [2] for more details about the singular solution.

The family of solutions (A, φ) can be packaged together into a "universal" solution to Hitchin's equations, living on a principal $PSU(n)$ bundle $\mathcal{U}$ over $\mathcal{M} \times C$. Restricted to any $[(A, \varphi)] \times C$, $\mathcal{U}$ is the principal bundle on which (A, φ) is defined. Restricting instead to the locus $\mathcal{M} \times \{z\} \subset \mathcal{M} \times C$ gives a principal bundle $\mathcal{U}_z$ over $\mathcal{M}$, attached to the surface defect $\mathbb{S}_z$.

A nice way to think about $\mathcal{U}_z$ is to consider the "pointed" gauge group $\mathcal{G}_z \subset \mathcal{G}$ consisting of gauge transformations which are the identity at z. We can then split the hyperkähler quotient construction of $\mathcal{M}$ into two steps: first take $\mathcal{N}_z := \mathcal{N}///\mathcal{G}_z$, and then $\mathcal{M} = \mathcal{N}_z///SU(n)$. The center $\mathbb{Z}_n \subset SU(n)$ acts trivially on $\mathcal{N}_z$, but the quotient $PSU(n) = SU(n)/\mathbb{Z}_n$ acts freely. Thus the zero locus of the hyperkähler moment map for the $su(n)$-action on $\mathcal{N}_z$ is a principal $PSU(n)$-bundle over $\mathcal{M}$. This principal bundle is $\mathcal{U}_z$. In particular, this way of building $\mathcal{U}_z$ makes it clear that it carries a canonical hyperholomorphic connection, using general properties of the hyperkähler quotient [37].

To define the "vector bundle" $V_{\mathbb{S}_z}$, we would like to take the bundle associated to $\mathcal{U}_z$ by the n-dimensional representation of $SU(n)$. There is a difficulty here: $\mathcal{U}_z$ is a $PSU(n)$ bundle, which does not have a canonical lift to an $SU(n)$ bundle. (In fact, in similar situations it is known that it does not lift at all [62, 63, 64].) Therefore, $V_{\mathbb{S},z}$ only exists as a twisted bundle over $\mathcal{M}$, twisted by a B-field of order n. This obstruction is consistent with the fact that the surface defect $\mathbb{S}_z$ carries a $\mathbb{Z}_n$ flavor monodromy, and hence, as we discuss in Appendix G, $V_{\mathbb{S}_z}$ should be twisted by a $\mathbb{Z}_n$-valued B-field.

Because there is a reduction of structure group of E at z_a to $K_a \subset SU(n)$, the fiber of E must decompose into irreducible representations of K_a. We write $E_a \simeq \oplus_s L_a^{(s)}$. In particular for $n = 2$ we have $E_a \simeq L_a \oplus L_a^{-1}$ for a complex line L_a. Now, the twisting of $V_{\mathbb{S}_z}$ over $\mathcal{M}$ is independent of z, and therefore $\mathrm{Hom}(V_{\mathbb{S}_{z'}}, V_{\mathbb{S}_z})$ should exist as an honest vector bundle over $\mathcal{M}$. Therefore, again restricting to $n = 2$, the lines $L_a^{\pm 1}$ should define twisted line bundles over $\mathcal{M}$ with the same twisting as $V_{\mathbb{S}_z}$.

Remarks

1. The fact that $\mathrm{Hom}(V_{\mathbb{S}_z}, V_{\mathbb{S}_{z'}})$ is a true bundle and not a twisted bundle has an important physical interpretation. The physical interpretation of sections of this bundle is related to expectation values of supersymmetric interfaces. Indeed, suppose we have a supersymmetric interface $L_\wp$ between surface defects $\mathbb{S}_z$ and $\mathbb{S}_{z'}$. There will be corresponding hyperholomorphic bundles V and V' over $\mathcal{M}$. In close analogy with [3], equation (7.13), the expectation value of the supersymmetric interface $\langle L_\wp \rangle$ should be identified with the parallel transport operator of $\mathcal{A}$ along $\wp$ from z to z' (or vice versa, depending on the orientation of $\wp$):

$$\langle L_\wp \rangle = \mathrm{Pexp}\left(- \int_\wp \mathcal{A} \right). \tag{7.16}$$

   Note that this equation only makes sense if we interpret $V \to \mathcal{M} \times C$ as the universal bundle. Then both the left hand side and the right hand side of this equation can be understood as elements of the vector space $\mathrm{Hom}(V_{\mathcal{A},z}, V_{\mathcal{A},z'})$, (when $\wp$ is oriented from z to z'), so it makes sense to equate them.

2. The (twisted) line bundles L_a over $\mathcal{M}$ also admit hyperholomorphic connections. Indeed, we can take the limit $z \to z_a$ in the construction of §5.6. Then $Z_{\gamma_{ij}(z)} \sim m_a \log(z - z_a) + \text{regular}$ so that the semiflat coordinates $\mathcal{Y}^{\text{sf}}_{\gamma_i}$ will require renormalization. These divergences suppress the integral terms in (5.25) (physically, the soliton masses go to infinity), and hence there is no mixing between line bundles. The connection ∇ on L_a obtained from the construction remains hyperholomorphic. We expect that the periods of $\frac{F(L_a,\nabla)}{2\pi}$ will be in $\frac{1}{n}\mathbb{Z}$, reflecting the twisting, but we have not checked this.

3. The bundles L_a should admit interpretations in terms of physical surface defects. We leave this interesting question to the future.

7.4 Lagrangian descriptions of A_1 theories and their surface defects

At this point we restrict attention from the general A_{n-1} theories to the A_1 case. In this class of theories $\Sigma \to C$ is a 2:1 covering. These theories have nice Lagrangian descriptions and are amenable to the kind of WKB analysis we discuss in §7.5. We consider theories associated to genus g Riemann surfaces C, carrying ℓ regular punctures. These theories have Lagrangian descriptions which are in one-to-one correspondence with pair-of-pants decompositions of C; each Lagrangian description involves $3g - 3 + \ell$ $SU(2)$ gauge groups corresponding to the tubes, and $2g - 2 + \ell$ hypermultiplets in the $(\mathbf{2}, \mathbf{2}, \mathbf{2})$ associated to the trinions [44]; call these hypermultiplets "hypertrinions." By taking careful decoupling limits of these theories, one can produce other A_1 theories, including asymptotically free Lagrangian theories but also non-Lagrangian, Argyres-Douglas-like theories. These limiting theories correspond to Riemann surfaces C carrying irregular punctures. See [2] for a detailed discussion.

Given the four-dimensional gauge theory description of the A_1 theories we can define the surface defects $\mathbb{S}_z$ as Gukov-Witten defects, at least in certain weak-coupling regimes. An explicit example of this was discussed in §3.4.2. More generally, $\mathbb{S}_z$ can be defined as a Gukov-Witten defect for any of the $SU(2)$ factors of the gauge group. This description is valid when the $SU(2)$ in question is weakly coupled; in that case one should think of C as developing a long tube, and the point of C labeling the surface defect as sitting somewhere on that tube. There is a natural coordinate t on the tube; the position of the surface defect in that coordinate system gives the (UV) Gukov-Witten parameter. As the surface defect moves around C from one long tube to another, 2d and 4d strong coupling effects mediate between different weakly coupled descriptions, as illustrated in the discussion of §3.4.2. In that discussion we noted that the 2d-4d instanton expansion breaks down due to light states when z hits a branch point. These light states are represented by the WKB curves from z to the nearby branch point.

We can now fill in a gap left open in our discussions of the twistings associated with Γ_i and $V_{\mathbb{S}_z}$. To do so we need to investigate the flavor symmetries in this class of theories. Associated to each puncture z_a of C is a flavor group $SU(2)_a$, so the full flavor group is $\mathcal{F} \simeq SU(2)^{\ell}$. The mass parameter of each factor $SU(2)_a$ is related to the residue m_a of the Higgs field at z_a. The flavor group $\mathcal{F}$ does not act effectively on the hypertrinions modulo gauge transformation. Indeed, identifying the center $Z(\mathcal{F}) \simeq \mathbb{Z}_2^\ell$, any two of the $\mathbb{Z}_2$ generators are equivalent up to a gauge transformation.[41] Therefore, letting $\mathcal{Z} \subset \mathbb{Z}_2^\ell$ be the index 2 subgroup consisting of products of even numbers of generators, the effective flavor group is $\bar{\mathcal{F}} = \mathcal{F}/\mathcal{Z}$, which sits in a sequence

$$1 \to \mathbb{Z}_2 \to \bar{\mathcal{F}} \to SO(3)^\ell \to 1. \tag{7.17}$$

The distinguished $\mathbb{Z}_2$ subgroup is the center $Z(\bar{\mathcal{F}})$, and can be represented (in many ways) as acting by -1 on an odd number of hypertrinions. Now, the chiral operator in the 4d $\mathcal{N} = 2$ theory which is not mutually local with respect to $\mathbb{S}_z$ is formed by taking the product of all the hypertrinions and contracting gauge indices [60]. Clearly this operator is projected out by gauging the central subgroup $Z(\bar{\mathcal{F}})$ and hence we conclude that if we want the troublesome (but desirable) surface defect $\mathbb{S}_z$ then we must gauge $Z(\bar{\mathcal{F}})$ and the flavor monodromy carried by $\mathbb{S}_z$ is the nontrivial element of this distinguished $\mathbb{Z}_2$ subgroup.

Now, following the discussion in §3.6.1, if we gauge $Z(\bar{\mathcal{F}})$, then there will be an extension of the group of gauge charges as in (3.71), with $\hat{\mathcal{D}} \simeq \mathbb{Z}_2$. We claim that this sequence, with $\mathcal{D} = Z(\bar{\mathcal{F}})$, can be identified with the sequence (3.82) given in Example 3.6.2, where z can be taken to be any of the z_a. Therefore, according to our discussion of that example, this UV flavor symmetry allows us to cancel the anomalies in Aharonov-Bohm phases arising from splitting the Γ_{ij}.

It remains to compare the twisting of $V_{\mathbb{S}_z}$ arising from the fractional shifts of Γ_{ij}, as described in §5.2.3, with that derived from the description in terms of hyperkähler quotients in §7.3. Consistency of our description of 2d-4d systems requires that they be the same, but we lack a direct proof. Two ways to approach the problem would be

1. Consider the $R \to \infty$ limit. Then the connection becomes diagonal, but a $\mathbb{Z}_n$ gerbe cannot depend on R, so the identification should be established in the semiflat geometry. It might be possible to combine the $R \to \infty$ limit with the $z \to z_a$ limit and thereby relate the V_i (in the $z \to z_a$ limit) with the L_a.

2. One could also relate the curvature $F(L_a, \nabla)$ to the B-field associated with the flavor symmetry $SU(2)_a$ as in Appendix G. Note that there are three real mass parameters associated with $SU(2)_a$: these are the complex residue m_a of the Higgs field and the real flavor Wilson line parameter $m_{a,3}$. In complex structure $\zeta = 0$, $m_{a,3}$ is a Kähler parameter and m_a a complex structure parameter, so ω_I is affine-linear in $m_{a,3}$, while $\omega_J + \mathrm{i}\omega_K$ are affine-linear and holomorphic in m_a. In real coordinates, with a proper normalization of $m_{a,i}$, we have

$$\frac{\partial \omega_i}{\partial m_{a,j}} = \delta_{ij} F(L_a, \nabla). \tag{7.18}$$

---

[41] To prove this, draw the trivalent graph representing the pants decomposition of C. The gauge transformation by the nontrivial element of the center of an $SU(2)$ factor associated to an edge is equivalent to an action by -1 on the two hypertrinions associated with the two vertices of that edge. Now use the fact that the graph is connected.

On the other hand, according to Appendix G, $\frac{\partial \omega_I}{\partial m_{a,3}}$ is the B-field associated with the flavor symmetry $SU(2)_a$. With this link between the twisting of L_a and the flavor symmetries established one might be able to prove the desired identity of twistings.

7.5 A brief review of the WKB analysis and its extension to 2d-4d

According to our general discussion in §5.8, the parallel transport operators $\langle L_\wp \rangle$ of (7.16) admit an expansion (5.43) in terms of canonically defined sections $\mathcal{Y}_{\gamma_{ij'}}$ of $\mathrm{Hom}(V, V')$. The coefficients of this expansion are the framed 2d-4d BPS degeneracies. The wall-crossing properties of these framed degeneracies contain the combinatorial data of the BPS degeneracies μ and ω.

For A_1 theories and their canonical surface defects, we can get our hands on the $\mathcal{Y}_{\gamma_{ij'}}$ very explicitly. The main tool is a slight amplification of the WKB analysis of [2], which we used in the pure 4d case to give a formula for the functions $\mathcal{Y}_\gamma$. Its extension similarly gives a formula for the sections $\mathcal{Y}_{\gamma_{ij'}}$. It also gives the expansion (5.43) of the $\langle L \rangle$ (at $y = -1$) in terms of framed 2d-4d BPS degeneracies (generalizing [3]). One can then verify directly that the wall-crossing of these framed degeneracies is indeed governed by the bulk 2d-4d BPS degeneracies μ and ω which we described above. In the rest of this section we will sketch how to construct the $\mathcal{Y}_{\gamma_{ij'}}$, assuming a certain degree of familiarity with the properties of A_1 Hitchin systems and with the results of [2]. We will leave a more detailed analysis, possible extension to other surface defects and to higher rank theories to a future publication.

7.5.1 4d review

The main player in our WKB analysis is the family of flat connections $\mathcal{A}$ on C given in (7.15). We have specialized to A_1 theories, so $\mathcal{A}$ is a flat $sl(2, \mathbb{C})$-valued connection on C, with singularities at various points of C, which we call "regular punctures" or "irregular punctures" depending on the nature of the singularities. We assume that there is at least one puncture.

Given a point $u \in \mathcal{B}$ and a choice of phase $\vartheta \in \mathbb{R}/2\pi\mathbb{Z}$, in [2] we defined the "WKB foliation" $F_{\mathrm{WKB}}(u, \vartheta)$. This foliation in particular determines a decomposition of C into a finite collection of simply connected cells C_{ab}. Here the indices a and b are labeling the possible asymptotic ends for generic WKB curves: these are either regular punctures or Stokes sectors around irregular punctures. Each cell C_{ab} is a union of WKB curves of phase ϑ which run from a to b. ($a = b$ is allowed, giving a "degenerate cell.") See Figure 25 in [2].

The WKB analysis of [2] relied strongly on the concept of a *small flat section* associated with a singularity z_a on C. This will be important in the 2d-4d extension also, so let us review the definition. Given a branch λ of the covering $\Sigma \to C$, WKB curves have a standard orientation, determined by $e^{-\mathrm{i}\vartheta}\langle \lambda, \partial_t \rangle > 0$ where ∂_t is a positively oriented tangent vector. Near a singularity z_a, choose the branch so that the WKB curve with its standard orientation flows *into* z_a. There then exists a flat section, $(\mathrm{d} + \mathcal{A}) s_a^\vartheta(z) = 0$, such that $s_a^\vartheta(z(t))$ becomes exponentially small as $t \to \infty$ and $z(t) \to z_a$. (Thus, s_a^ϑ also grows as z moves away from z_a along a WKB curve.) Moreover, this condition determines the section s_a^ϑ uniquely up to an overall scale. Such a small flat section will be variously denoted as $s_a^\vartheta(z;u,\zeta)$, $s_a^\vartheta(z)$, or just s_a^ϑ. Our main constructions involving s_a^ϑ do not require us to choose the scale. Now suppose that ζ is in the half-plane $\mathbb{H}_\vartheta$ centered on the ray $\zeta \in e^{\mathrm{i}\vartheta}\mathbb{R}_+$. Then, the WKB formula says that for the small flat section the expression

$$s_a^\vartheta(z)\exp\left(\frac{R}{\zeta}\int_{z_a(\epsilon)}^{z}\lambda\right), \tag{7.19}$$

where the integral is along the curve with $e^{-\mathrm{i}\vartheta}\langle\lambda,\partial_t\rangle \in \mathbb{R}$ from $z_a(\epsilon)$ to z, has a finite limit as $\zeta \to 0$ in the half-plane $\mathbb{H}_\vartheta$. Here $z_a(\epsilon)$ is infinitesimally close to the singular point z_a.

One of the main results of [2] was a direct construction of the "Darboux functions" $\mathcal{Y}_\gamma$ given the data of a WKB triangulation and its small flat sections. A somewhat streamlined version of the construction can be summarized as follows. Given ϑ we construct antisymmetric inner products $T_{ab}^\vartheta := (s_a^\vartheta, s_b^\vartheta)$ of the sections associated to a cell C_{ab}. Here the inner product of two sections (s,s') is defined to be $\frac{s\wedge s'}{\mathrm{vol}}$ where vol is the $SL(2,\mathbb{C})$ invariant "volume" form. We stress that it is z-independent. The asymptotics of s_a^ϑ imply similar asymptotics for T_{ab}^ϑ, controlled by $\int_{z_b(\epsilon)}^{z_a(\epsilon)}\lambda$. More precisely, let $P_a^\pm$ be the lifts of the singular points z_a on Σ.[42] Then, for each ordered pair a,b there is a unique oriented lift $\hat{E}_{ab}\subset\Sigma-\{P_a^\pm\}$ of the unoriented edge $e_{ab}\subset C$ which is odd under the deck transformation and satisfies

$$T_{ab}^\vartheta e^{-\frac{R}{\zeta}\oint_{\hat{E}_{ab}}\lambda} \sim \text{finite} \tag{7.20}$$

as $\zeta\to 0$ in the half-plane $\mathbb{H}_\vartheta$. (Once again we need to use the ϵ regularization.) Note that since $T_{ab}^\vartheta = -T_{ba}^\vartheta$ we have $\hat{E}_{ab} = \hat{E}_{ba}$.

Now, let n_{ab} be any matrix of integers such that the diagonal elements vanish and for every a,

$$\sum_b (n_{ab}+n_{ba}) = 0. \tag{7.21}$$

To such a matrix we associate

$$\mathcal{X}_n := \prod_{a,b}\left(T_{ab}^\vartheta\right)^{n_{ab}}. \tag{7.22}$$

The condition (7.21) guarantees that each small flat section s_a enters the product $\mathcal{X}_n$ as many times in the numerator and denominator and hence is independent of the normalization of s_a.[43] Thus $\mathcal{X}_n$ are well-defined on $\mathcal{M}$. Moreover, by (7.20) they have $\zeta\to 0$ asymptotics of the form

$$\mathcal{X}_n \sim \mathcal{N}_n \exp\left(\frac{R}{\zeta}\oint_{\gamma(n)}\lambda\right) \tag{7.23}$$

where $\mathcal{N}_n$ has a well-defined limit for $\zeta\to 0$ and $\gamma(n) = \sum_{ab} n_{ab}\hat{E}_{ab}$. The expression $\gamma(n)$ defines, *a priori*, a class in the relative homology $H_1(\Sigma,\{P_a^\pm\};\mathbb{Z})^-$. However, thanks to

---

[42] The notation presupposes regular singularities, but the generalization to irregular singularities is clear.

[43] Combinations $\prod T_{ab}^{n_{ab}}$ have been considered many times in the literature — in particular the "Fock-Goncharov coordinates" [65] are of the form $\frac{T_{ab}T_{cd}}{T_{bc}T_{da}}$ where a,b,c,d are the four vertices of some quadrilateral in an ideal triangulation of C.

(7.21) that class is in the image of $H_1(\Sigma;\mathbb{Z})^-$. Moreover the map $n \to \gamma(n)$ is linear, onto, and has kernel consisting of antisymmetric matrices.[44] Finally, $\mathcal{Y}^\vartheta_\gamma$ will be defined by

$$\mathcal{Y}^\vartheta_\gamma := \varepsilon(\gamma, n)\mathcal{X}_n, \tag{7.24}$$

where n on the right side obeys $\gamma = \gamma(n)$, and $\varepsilon(\gamma, n) = \pm 1$. Since the map $n \to \gamma(n)$ has a kernel, it is not immediately obvious that this "definition" is well defined. Fortunately, if $k_{ab} = -k_{ba}$ is antisymmetric then $\mathcal{X}_{n+k} = (-1)^{\sum_{a<b} k_{ab}}\mathcal{X}_n$, so (7.24) is indeed well defined provided we take $\varepsilon(\gamma, n+k) = (-1)^{\sum_{a<b} k_{ab}}\varepsilon(\gamma, k)$. To fix ε completely, we need a little more input. In [2] we fixed it by determining it on a basis of Γ and then using the multiplicative law of the $\mathcal{Y}_\gamma$. This is briefly reviewed in Appendix F.

Finally, the Darboux functions are defined by specializing ϑ, $\mathcal{Y}_\gamma := \mathcal{Y}^{\vartheta=\arg\zeta}_\gamma$.

There is a well-known algorithm [66, 65, 67, 68], reviewed in our context in [2, 3], for decomposing the trace of the parallel transport around a loop $\wp$ as a linear combination of the $\mathcal{Y}_\gamma$. Physically this gives the "Darboux expansion" (5.42) of the line defect vev $\langle L_\wp\rangle$ in terms of the "IR line defect vevs," as explained at length in [3].

As we have recalled in §§4 and 5.7, the functions $\mathcal{Y}_\gamma$ are supposed to have discontinuities when (u,ζ) meet any "BPS wall" W_γ, defined in (4.4), with the jump across W_γ identified with a transformation $\mathcal{K}^{\Omega(\gamma)}_\gamma$. This jump reflects the effect of 4d BPS particles of charge γ. In the A_1 theories, these discontinuities arise from jumps in the topology of the WKB foliation $F_{\mathrm{WKB}}(u, \vartheta = \arg\zeta)$ as we vary (u,ζ). The topological content of this foliation is captured by the "decorated WKB triangulation" $T_{\mathrm{WKB}}(u,\vartheta)$. The vertices of $T_{\mathrm{WKB}}(u,\vartheta)$ are the asymptotic ends a, decorated by the sections s_a. The edges of $T_{\mathrm{WKB}}(u,\vartheta)$ are in 1-1 correspondence with the cells C_{ab}. The 4d spectrum of the A_1 theory follows directly from the combinatorics of how $T_{\mathrm{WKB}}(u,\vartheta)$ evolves as ϑ is varied.

More precisely, as ζ varies, $T_{\mathrm{WKB}}(u,\vartheta = \arg\zeta)$ may jump in three different ways, which we call the *flip*, *juggle* and *pop*:

- A flip is a very simple change: we remove one edge E, leaving behind a quadrilateral Q_E, and put back another edge along the other diagonal of Q_E. This jump of T_{WKB} leads to a corresponding transformation of the $\mathcal{Y}_\gamma$: it is $\mathcal{K}_{\gamma_E}$, where γ_E is the lift to Σ of a cycle running around the pair of turning points inside the quadrilateral. See §7.1 of [2] for further details. The occurrence of a flip is signaled by the presence of a finite WKB curve (necessarily of phase $\vartheta = \arg -Z_{\gamma_E}$) joining the two turning points in Q_E; if such a finite WKB curve exists, then T_{WKB} undergoes a flip of edge E, at the BPS ray $Z_{\gamma_E}/\zeta \in \mathbb{R}_-$. This flip corresponds to a BPS hypermultiplet of charge γ_E, i.e. to $\Omega(\gamma_E) = 1$.

- A pop replaces the chosen small flat section s_a at some regular singularity a with the opposite monodromy eigenvector $\tilde{s}_a$. When T_{WKB} undergoes a pop, the functions $\mathcal{Y}_\gamma$ are not changed, but for a rather subtle reason: the functions T_{ab} change, but the recipe for building $\mathcal{Y}_\gamma$ from the T_{ab} also changes, in just such a way that the $\mathcal{Y}_\gamma$ are

---

[44] To prove this we use the dual pairing of $\hat{E}_{ab}$ with the cycles γ_E associated to the edges of the WKB triangulation $T_{\mathrm{WKB}}(\vartheta)$. See §7.1.1 of [2].



Figure 15: The effect of a flip on T_{WKB}.

left invariant. (See §7.6.3 of [2].) A pop occurs at the BPS ray corresponding to a pure flavor central charge γ_a^f, $Z_{\gamma_a^f}/\zeta \in \mathbb{R}_-$. The triangulation $T_{\mathrm{WKB}}(u,\zeta)$ for (u,ζ) near this BPS ray is always *degenerate*: it has only a single edge coming out of the vertex a. The pop is signaled by the appearance of a closed WKB curve surrounding the singularity z_a. (The cycle γ_a^f is an odd lift of this closed WKB curve.) In the language of the 4d theory, this closed WKB curve corresponds to a BPS state carrying only flavor charges.

- A juggle is a subtler transformation: it does not relate two triangulations in the ordinary sense, but rather two *limits* of triangulations, each obtained from an infinite sequence of flips of the common edges of two triangles which form an annulus on C. The effect of the juggle on the limiting coordinates $\mathcal{Y}_\gamma$ is a $\mathcal{K}$-transformation $\mathcal{K}_{\gamma_0}^{-2}$, where γ_0 is an odd lift of a closed loop going around the annulus once. This transformation appears sandwiched between the two infinite sequences of transformations coming from the flips. The juggle happens at the BPS ray $Z_{\gamma_0}/\zeta \in \mathbb{R}_-$. It is signaled by the appearance of a family of closed WKB curves running around the annulus, corresponding to a BPS vector multiplet of charge γ_0.



Figure 16: A juggle arises as the limit of an infinite sequence of flips.

7.5.2 Extension to 2d-4d

Now let us construct the Darboux sections $\mathcal{Y}_{\gamma_{ij'}}$ for the 2d-4d case. These have been defined in §5 in a way analogous to the definition of the $\mathcal{Y}_\gamma$ in [1]. However, as in [2], for A_1 theories there should also be a direct expression in terms of small flat sections, analogous to (7.24). Now we should have $\mathcal{Y}^{\vartheta}_{\gamma_{ij'}} \in \mathrm{Hom}(V_{\mathbb{S}_z}, V_{\mathbb{S}_{z'}})$ where one end of $\gamma_{ij'}$, x_i, lies above z and the other end, $x_{j'}$, lies above z'. [45]

The $\mathcal{Y}^{\vartheta}_{\gamma_{ij'}}$ should be piecewise holomorphic in z, z', u, ζ with wall-crossing discontinuities discussed above and moreover should satisfy reality constraints and, importantly, the asymptotic condition that

$$\mathcal{Y}^{\vartheta}_{\gamma_{ij'}} e^{-\frac{R}{\zeta}\int_{\gamma_{ij'}}\lambda} \tag{7.25}$$

---

[45] Unfortunately, we have reversed conventions here relative to §5.6, where we would have taken $\mathcal{Y}_{\gamma_{ij'}}$ to be in $\mathrm{Hom}(V_{\mathbb{S}_{z'}}, V_{\mathbb{S}_z})$.

has a finite limit as $\zeta \to 0$ in $\mathbb{H}_\vartheta$. Finally, these sections should satisfy

$$\mathcal{Y}_{\gamma_{ij'}}\mathcal{Y}_{\gamma_{j'k''}} = \sigma(\gamma_{ij'}, \gamma_{j'k''})\mathcal{Y}_{\gamma_{ij'}+\gamma_{j'k''}} \tag{7.26}$$

for the twisting function defined in (7.6).

In analogy to the $\mathcal{X}_n$, we again take n_{ab} to be a matrix of integers with zero on the diagonal. Then we define the analog of (7.22):

$$\mathcal{X}^\vartheta_{n,c,d}(z, z') := \prod_{a,b}(T^\vartheta_{ab})^{n_{ab}} s^\vartheta_c(z) \otimes s^\vartheta_d(z'). \tag{7.27}$$

Here z lies in some cell C_{a_0,b_0} and c is one of a_0 or b_0, while z' lies in some cell $C_{a'_0,b'_0}$ and d is one of a'_0 or b'_0. Here and in several places below, we use the $SL(2,\mathbb{C})$-invariant volume form to identify V with V^*. To be explicit, $s^\vartheta_a(z) \otimes s^\vartheta_b(z') \in \mathrm{Hom}(V_{\mathbb{S}_z}, V_{\mathbb{S}_{z'}})$ is defined by

$$s^\vartheta_a(z) \otimes s^\vartheta_b(z') : v \mapsto (v, s^\vartheta_a)s^\vartheta_b(z') \tag{7.28}$$

and thus (7.27) indeed determines a section of $\mathrm{Hom}(V_{\mathbb{S}_z}, V_{\mathbb{S}_{z'}})$. It will be independent of the choice of normalization of the s^ϑ_a provided that n_{ab} satisfies the analog of (7.21),

$$\sum_b (n_{ab} + n_{ba}) = \begin{cases} -1 & a = c, d, \\ 0 & \text{otherwise.} \end{cases} \tag{7.29}$$

Now we need the analog of the map $n \to \gamma(n)$ we found in the case when γ is closed. As before this will be determined by $\zeta \to 0$ asymptotics. As we have explained, to a singularity z_a we can associate a branch λ, such that WKB curves with their standard orientation flow into z_a. Given $z \in C_{a,b}$, let $x_{a,z} \in \Sigma$ be the lift of z to this branch. Then there is a lift of the WKB curve between z_a and z to an oriented path $\hat{E}_{a,x_{a,z}}$ between a lift of z_a and $x_{a,z}$, such that

$$s^\vartheta_a(z) \exp\left(-\frac{R}{\zeta}\int_{\hat{E}_{a,x_{a,z}}} \lambda\right) \tag{7.30}$$

is finite as $\zeta \to 0$ in $\mathbb{H}_\vartheta$. It follows that

$$\mathcal{X}^\vartheta_{n,c,d}(z, z') \exp\left(-\frac{R}{\zeta}\int_{\gamma_{ij'}(n,c,d)} \lambda\right) \tag{7.31}$$

is finite for $\zeta \to 0$ in $\mathbb{H}_\vartheta$, where we have defined[46]

$$\gamma_{ij'}(n, c, d) := -\sigma^*(\hat{E}_{c,x_{c,z}}) + \sum_{a,b} n_{ab}\hat{E}_{ab} + \hat{E}_{d,x_{d,z'}}. \tag{7.32}$$

As with the closed curves, we can now define $\mathcal{Y}^\vartheta_{\gamma_{ij'}}$ by

$$\mathcal{Y}^\vartheta_{\gamma_{ij'}} := \varepsilon(\gamma_{ij'}, n)\mathcal{X}^\vartheta_{n,c,d}(z, z') \tag{7.33}$$

where $\gamma_{ij'} = \gamma_{ij'}(n, c, d)$. As before, $\varepsilon(\gamma_{ij'}, n)$ is a slightly delicate sign; a systematic way of fixing it is given in Appendix F.



Figure 17: z is in the cell C_{ab} and also in the triangle abc, with abc in counterclockwise order. The first sheet is such that with orientation $e^{-i\vartheta}\langle\lambda,\partial_t\rangle > 0$ the WKB curves flow from a to b, counterclockwise on the triangle. The green curves are the projections of open paths γ_{12} and $\tilde{\gamma}_{21}$ connecting sheets 1 and 2 above z so that $\gamma_{12} + \tilde{\gamma}_{21} = -\gamma_E$.

Finally, the $\mathcal{Y}^{\vartheta=\arg\zeta}_{\gamma_{ij'}}$ are to be identified with the $\mathcal{Y}_{\gamma_{ij'}} \in \mathrm{Hom}(V_{\mathbb{S}_z}, V_{\mathbb{S}_{z'}})$ which we are after.

Example. Suppose z is in the cell C_{ab} as in Figure 17, and we define sheet 1 to be the root λ so that the WKB curve with $e^{-i\vartheta}\langle\lambda,\partial_t\rangle > 0$ through z is oriented to flow counterclockwise around the triangle, i.e. from a to b. Then $x_{b,z}$ lies on sheet 1 and $x_{a,z}$ lies on sheet 2. Now consider $\gamma_{11} = 0 \in \Gamma_{11}$. This is associated with

$$\mathcal{Y}_{\gamma_{11}=0} = \frac{s_a(z) \otimes s_b(z)}{(s_b, s_a)}, \tag{7.34}$$

while $\gamma_{22} = 0 \in \Gamma_{22}$ is given by

$$\mathcal{Y}_{\gamma_{22}=0} = \frac{s_b(z) \otimes s_a(z)}{(s_a, s_b)}. \tag{7.35}$$

Now consider the simple path $\gamma_{12}(z)$ from sheet 1 to sheet 2 and projecting to a simple curve from z to the turning point in the triangle abc, as in Figure 17. Then

$$\mathcal{Y}_{\gamma_{12}} = \frac{(s_a, s_c)}{(s_b, s_a)(s_b, s_c)} s_b(z) \otimes s_b(z). \tag{7.36}$$

As a check, if we compose with a similar expression for the open curve $\tilde{\gamma}_{21}$ whose projection is shown in Figure 17, so that $\gamma_{12} + \tilde{\gamma}_{21} = -\gamma_E$, we find indeed that

$$\begin{aligned} \mathcal{Y}_{\gamma_{12}} \mathcal{Y}_{\tilde{\gamma}_{21}} &= \frac{(s_f, s_b)(s_a, s_c)}{(s_a, s_f)(s_b, s_c)} \cdot \frac{s_a(z) \otimes s_b(z)}{(s_b, s_a)} \\ &= -\mathcal{Y}_{-\gamma} \mathcal{Y}_{\gamma_{11}}, \end{aligned} \tag{7.37}$$

where $\gamma_{11} = 0 \in \Gamma_1$. This is summarized in the equation $\mathcal{Y}_{\gamma_{12}} \mathcal{Y}_{\tilde{\gamma}_{21}} = \sigma(\gamma_{12}, \tilde{\gamma}_{21}) \mathcal{Y}_{\gamma_{12}+\tilde{\gamma}_{21}}$.

---

[46] We use σ for both the deck transformation and the twisting function. We hope these will not be confused.



Figure 18: The parameter z of a surface defect is changed so that it moves from a cell C_{ab} to a cell C_{ac} across a separating WKB curve. Blue dots are singularities, and orange crosses are branch points, aka turning points. A similar picture applies if z is held fixed but (u, ζ) are changed so that a separating WKB curve sweeps across z.

To complete the demonstration that the $\mathcal{Y}_{\gamma_{ij'}}$ are correctly defined, we must demonstrate how the full 2d-4d BPS spectrum is captured by the discontinuities of $\mathcal{Y}_{\gamma_{ij'}}$. In this case the combinatorial data that goes into $\mathcal{Y}_{\gamma_{ij'}}$ is slightly more than what we had in the pure 4d case: we need the WKB triangulation T_{WKB} *plus* the information of to which cell C_{ab} the points z, z' belong. As we vary $\vartheta = \arg \zeta$ these combinatorial data might jump, and we ask how these jumps induce jumps in $\mathcal{Y}_{\gamma_{ij'}}$. We have to consider flips, pops and juggles of T_{WKB} as before, as well as the new possibility that z or z' moves from one cell to another:

- Suppose z moves from a cell C_{ab} to a cell C_{ac} as in Figure 18. In this case we can use the simple identity $s_a T_{bc} + s_b T_{ca} + s_c T_{ab} = 0$ to relate the $\mathcal{Y}^{\vartheta}_{\gamma_{ij'}}$ before and after the jump. The resulting transformation turns out to be an $\mathcal{S}$-factor, with $\mu_{\gamma_{ij}} = 1$, where i and j refer to the two preimages of z, and γ_{ij} is the path which starts from one preimage of z, runs to the branch point in the middle of the triangle (abc), and comes back to the other preimage of z. The detailed demonstration of this can be found in Appendix F. If we hold z fixed and let ϑ vary, such a jump will occur whenever there is a WKB curve of phase ϑ which connects z to the branch point. Such a finite WKB curve represents a 2d soliton with charge γ_{ij}. A very similar story also holds with z replaced by z'.
- A flip of T_{WKB} occurs when, as ϑ is varied, one of the cells of C shrinks to zero size. It follows that at the ϑ where the flip occurs, z and z' are (at least generically) not contained in this collapsing cell: even if they were in that cell for some value of ϑ, they will fall out before the flip is reached. From this it follows that the flip affects $\mathcal{Y}_{\gamma_{ij'}}$ only through its action on the functions T_{ab}. Indeed, the action turns out to be

precisely given by a transformation $\mathcal{K}^{\omega}_{\gamma_{ij'}}$, where $\omega(\gamma_E, \gamma_{ij'}) = \langle \gamma_E, \gamma_{ij'} \rangle$ (in the sense we defined in §7.2 above.)

- A pop of T_{WKB} occurs when the WKB curves going into a regular singularity a start winding around a more and more and finally become closed curves. In particular, near the ϑ where the pop occurs, T_{WKB} contains a degenerate cell, which contains a in its interior and also contains a WKB curve which runs from a turning point to a. As ϑ approaches the critical value, this WKB curve winds more and more around a. In particular, if z lies in this degenerate cell, there are infinitely many values of ϑ for which this WKB curve meets z. These values accumulate at the critical ϑ where the pop occurs. As we have explained above, each time the WKB curve meets z, the $\mathcal{Y}_{\gamma_{ij'}}$ undergo an $\mathcal{S}$ transformation: so we get an infinite sequence of such transformations, corresponding to solitons with charges $\gamma_{ij} + n\gamma_a^f$, as $n \to \infty$. On the other side of the critical ζ we have a second infinite sequence of $\mathcal{S}$ transformations corresponding to solitons with charges $\gamma_{ji} + n\gamma_a^f$ as $n \to \infty$. We may then ask whether there is a further transformation sandwiched between these two infinite sequences, exactly at the value of ϑ where the pop occurs. It turns out that the answer is yes: there is a single $\mathcal{K}^{\omega}_{\gamma_a^f}$ transformation, with $\omega(\gamma_a^f, \cdot)$ corresponding to a 2d particle carrying only 4d flavor charges. One could in principle prove this directly by studying the limits of $\mathcal{Y}_{\gamma_i}(\zeta)$ as ζ approaches the critical value from either side, along the lines of what was done for the juggle transformation of the $\mathcal{Y}_\gamma$ in [2]. In practice, it is faster to work indirectly — move the point z just outside the degenerate cell, so that the infinite sequences of transformations disappear, and then use the wall-crossing formula (2.51) to deduce what happens when z is in the degenerate cell. See §8.2 below for a concrete illustration of these remarks.

- A juggle of T_{WKB} occurs when closed WKB curves form an annular region. If z and z' are outside the annular region where the juggle is taking place, then the effect of the juggle on the $\mathcal{Y}_{\gamma_{ij'}}$ is simply a $\mathcal{K}^{\omega}_{\gamma_0}$ transformation with $\omega(\gamma_0, \gamma_{ij'}) = -2\langle \gamma_0, \gamma_{ij'} \rangle$. This is a straightforward extension of our discussion above about the juggle in the pure 4d case. Also as in the pure 4d case, this transformation is surrounded by two infinite sequences of $\mathcal{K}$ corresponding to flips. The more interesting case is when either z or z' lies in the annulus. Then one finds an infinite sequence of $\mathcal{S}$-factors interspersed with these infinite sequences of $\mathcal{K}$, as the WKB curves wind more and more around the annulus. At the critical ζ where the juggle occurs, there is a $\mathcal{K}$-factor with appropriate $\omega(\gamma_0, \cdot)$. Much as we just discussed for the pop, the simplest approach to determining exactly what transformation occurs at the critical ζ is to use a wall-crossing formula. We give an example in §8.3.

Remarks

1. The analog of the Darboux expansion for interfaces $L_\wp$ associated with open curves $\wp$ on C from z to z' is given in (5.43). As in the case of closed curves, for A_1 theories we

can compute the framed BPS degeneracies quite explicitly, using a standard algorithm for writing the parallel transport from z to z' as a linear combination of $\mathcal{Y}^{\vartheta}_{\gamma_{ij'}} \in \mathrm{Hom}(V, V')$. For details see, for example, [2], Appendix A, equation $(A.8)$.

2. It should be possible to define the more subtle quantities $\mathcal{Y}_{\gamma_i}$ in terms of expressions of the form $\prod_{a,b} T_{ab}^{n_{ab}} s_c(z)$ where the n_{ab} are now half-integers, raising issues with the choices of squareroots. We will not attempt to define them here, but we note that doing so would amount to an explicit construction of the twisting B field of the universal bundle, so it might be worthwhile to work this out carefully.

3. Finally let us remark that there exists a 2d-4d analog of the "spectrum generating transformation" written down in §11 of [2]. There we wrote an explicit formula for the $\mathcal{Y}^{\vartheta}_{\gamma}$ in terms of the opposite monodromy eigenvectors $\tilde{s}_{a,b,c,\cdots}$, which is equivalent to working out the transformation relating $\mathcal{Y}^{\vartheta}_{\gamma}$ to $\mathcal{Y}^{\vartheta+\pi}_{\gamma}$. This transformation turned out to be given by a combinatorial recipe depending only on the triangulation $T_{\mathrm{WKB}}(\vartheta)$. On the other hand this transformation is the composition of all the factors $\mathcal{K}_{\gamma}^{\Omega(\gamma)}$ attached to the BPS rays which one encounters in varying $\arg\zeta$ from ϑ to $\vartheta+\pi$, and hence contains complete information about the 4d BPS spectrum. All of that discussion can be generalized to the 2d-4d setting: one can give a combinatorial recipe for the transformation relating $\mathcal{Y}^{\vartheta}_{\gamma_{ij'}}$ to $\mathcal{Y}^{\vartheta+\pi}_{\gamma_{ij'}}$, as an automorphism $\mathbf{S}$ of the vacuum groupoid algebra. This $\mathbf{S}$ has a unique ordered decomposition of the form
$$\mathbf{S} = \ : \prod_{\vartheta < \arg -Z_{\gamma}, \arg -Z_{\gamma_{ij'}} < \vartheta+\pi} \mathcal{K}^{\omega}_{\gamma} \mathcal{S}^{\mu}_{\gamma_{ij'}} : \tag{7.38}$$
which determines the complete 2d-4d BPS spectrum. We do not write the details here. The main new ingredient is a formula for the transformation taking $s_a \to \tilde{s}_a$ for a pop at a single vertex. It involves a rational expression of the Fock-Goncharov coordinates on all the edges of a star neighborhood of the vertex a. We will defer the details to another occasion.

8. Examples of 2d-4d wall-crossing in some A_1 theories

In this section we illustrate the general remarks of §7 with a few simple examples. Many of the more intricate examples of [17] could be reconsidered along the lines below, but we have not done this yet. We expect that some rich phenomena remain to be discovered in those examples.

In the following sections we will be writing formulas for $A(\lhd)$ in various different theories and regions of moduli space. To lighten the notation, we take advantage of the fact that, with our sign convention for σ understood, $\mathcal{S}^{\mu}_{\gamma_{ij}}$ only depends on μ through the value $\mu(\gamma_{ij})$; this value is always $+1$ or zero in our examples, so we drop the exponent and just write $\mathcal{S}_{\gamma_{ij}}$ when $\mu(\gamma_{ij})$ is nonzero. We also sometimes drop the explicit ω in $\mathcal{K}^{\omega}_{\gamma}$ when it is unchanged in a wall-crossing formula.

8.1 Simple Argyres-Douglas-type theories

A particularly simple class of theories for studying wall-crossing are the Argyres-Douglas-type superconformal theories of A_1 type. These theories are obtained by compactifying the A_1 $(2,0)$ theory from six to four dimensions on $C = \mathbb{CP}^1$, with an irregular singularity at infinity, whose strength is specified by an integer $N \geq 1$. (Usually one only considers $N \geq 3$, since the theories with $N = 1, 2$ have rather trivial 4d dynamics. As we will see below, though, even the case $N = 1$ is interesting when coupled to a surface defect.)

Following the discussion of §7, these theories correspond to the Hitchin system on $\mathbb{CP}^1$ with gauge group $SU(2)$ and a single irregular puncture at infinity. The Coulomb branch $\mathcal{B}$ is thus parameterized by meromorphic quadratic differentials $\phi_2(z)$ on $\mathbb{CP}^1$ of the form

$$\phi_2(z) = -P_N(z)\,\mathrm{d}z^2, \tag{8.1}$$

where $P_N(z)$ is a polynomial of degree N. The low-order coefficients of $P_N(z)$ are coordinates on the Coulomb branch, while the higher-order ones are parameters of the theory (see e.g. [2], §9.2).

The BPS spectrum of these theories has been investigated in [69] and revisited from our current perspective in [2]. At any point of $\mathcal{B}$, the 4d BPS spectrum consists of a *finite* set of hypermultiplets. The wall-crossing phenomena which occur when we move in $\mathcal{B}$ can always be understood using only the basic identity "$\mathcal{K}\mathcal{K} = \mathcal{K}\mathcal{K}\mathcal{K}$" (2.14).

Below we consider the 2d-4d BPS spectrum that we get by coupling these theories to their canonical surface defects $\mathbb{S}_z$. As we will see, the answer is again very simple: we will find a finite set of 2d solitons between the two vacua. Studying these solitons will provide examples of the various phenomena we have discussed in this paper. For $N = 1$ there is no wall-crossing for the solitons, but the solitons do induce wall-crossing for framed 2d BPS states. For $N > 1$ the soliton spectrum undergoes wall-crossing, which provide nice examples of the finite 2d-4d wall-crossing formulae. The essential formulae we need are summarized by special cases of (2.36) and (2.44). In the A_1 theories the only identities we will need are:

1. First, the result when $\mathbb{S}_z$ crosses a separating WKB curve:

$$\mathcal{K}_a \mathcal{S}_b = \mathcal{S}_b \mathcal{S}_{b+a} \mathcal{K}_a \qquad \omega(a,b) = -1 \tag{8.2a}$$

$$\mathcal{S}_b \mathcal{K}_a = \mathcal{K}_a \mathcal{S}_{b+a} \mathcal{S}_b \qquad \omega(a,b) = +1 \tag{8.2b}$$

   Note that this does not change the ω degeneracy, so we have suppressed the superscript.

2. Second, the result when $\mathbb{S}_z$ crosses a finite WKB curve:

$$\mathcal{S}_b \mathcal{K}_a^{\omega} \mathcal{S}_{a-b} = \mathcal{S}_{a-b} \mathcal{K}_a^{\omega'} \mathcal{S}_b \tag{8.3}$$

   which leaves the soliton spectrum unchanged but reverses the sign $\omega(a,b) = -1$ to $\omega'(a,b) = +1$. As discussed in the example at the end of §7.2 above, what has happened is that the meaning of the charge "b" has changed since the point z has moved. The prescription for computing ω in terms of intersection numbers remains in force.

8.1.1 $N = 1$

The $N = 1$ theory is obtained by taking $P(z) = z$. In this theory the 4-dimensional IR physics is trivial, so there are no 4d gauge fields; there are also no flavor symmetries, so the charge lattice Γ is trivial, and there are no 4d BPS states. Although the four-dimensional theory is trivial there can still be a nontrivial two-dimensional theory. Indeed we will see that this theory corresponds to a Landau-Ginzburg theory with $W = \frac{1}{3}x^3 - zx$, which we have already studied in §6.3 and Appendix B.[47]

When we include the canonical surface defect $\mathbb{S}_z$ associated to a point $z \in C$, the theory does have 2d BPS states. To see them explicitly we can use our description from §7.2: a 2d BPS state corresponds to a finite WKB curve running from the turning point at $z = 0$ to z. The finite WKB curves emerging from the turning point in this case are just three straight rays from 0 to ∞, with inclinations $\frac{2}{3}\vartheta$, $\frac{2}{3}(\vartheta + 2\pi)$, $\frac{2}{3}(\vartheta + 4\pi)$. As we vary ϑ through a phase π (say from 0 to π), the three rays rotate, and one of them sweeps across the point z. Hence there is exactly one 2d BPS state with phase between 0 and π. The full BPS spectrum consists of this particle plus its antiparticle (which has phase between π and 2π). These BPS states persist for all values of z, which is the only modulus around, so there is no wall-crossing for the ordinary 2d BPS spectrum. The spectrum of vacua and BPS states coincide with those of a Landau-Ginzburg model with superpotential $\frac{1}{3}x^3 - zx$. From this point of view, our statements will just be a rewriting of standard statements from tt^* lore [58, 32].



Figure 19: WKB sectors and interface paths for three values of ζ. The separating WKB curves emerging from the turning point are illustrated for three values of $\vartheta = \arg \zeta$. In (A) ζ is a positive real number. As the phase of ζ is increased the picture rotates counterclockwise to produce figures (B) and (C).

Although there is no wall-crossing for the ordinary BPS states, there is still some wall-crossing in the story: indeed the presence of 2d BPS states implies that there is wall-crossing for 2d *framed* BPS states. Let us explore this a bit. We consider an interface $L_{\wp,\zeta}$ between two canonical surface defects $\mathbb{S}_z$, $\mathbb{S}_{z'}$ associated to a path $\wp$ from z to z' as shown in Figure 19. We initially assume that z and z' are "close" in the sense that the angle

---

[47] Although the mathematics is the same, there is an important conceptual difference. The Hitchin system in §6.3 is on the u-plane $\mathcal{B} = \mathbb{C}$ and there is a 4d $U(1)$ gauge theory. Here there is no 4d gauge theory, $\mathcal{B}$ is a point, and the Hitchin system is on the Riemann surface C.



Figure 20: Using the principal branch of the logarithm to define $z^{1/2}$ determines a sheet of the covering $\Sigma \to C$, which we denote $+$. The standard orientation of WKB curves is shown here for WKB curves on sheet $+$, at $\vartheta = 0$. This orientation allows us to fix a small flat section given a point $z \in C$ and a choice of sheet above z.

between them is less than $2\pi/3$, and discuss the case when the angle is between $2\pi/3$ and π later.

First of all, it is straightforward to express $\langle L_{\wp,\zeta} \rangle$. The small flat sections are defined first in the neighborhood of singular points and then defined for other values of z by parallel transport. Using the convention (7.28) we can write

$$\langle L_{\wp,\zeta} \rangle = \frac{s_1(z) \otimes s_3(z')}{(s_3, s_1)} + \frac{s_3(z) \otimes s_1(z')}{(s_1, s_3)} \tag{8.4a}$$

$$= \frac{s_1(z) \otimes s_2(z')}{(s_2, s_1)} + \frac{s_2(z) \otimes s_1(z')}{(s_1, s_2)} \tag{8.4b}$$

$$= \frac{s_2(z) \otimes s_3(z')}{(s_3, s_2)} + \frac{s_3(z) \otimes s_2(z')}{(s_2, s_3)}. \tag{8.4c}$$

All three of these expressions are valid for any z, z'. Which one of them is *useful* for extracting framed BPS states depends on the sectors to which z, z' belong. Recall that the expressions $\mathcal{X}_{n,c,d}(z, z')$ with good asymptotics are only defined for $c \in \{a, b\}$ when $z \in C_{ab}$ and similarly for d. Therefore in case (A) of Figure 19 we should use (8.4a). In accordance with our discussion of equations (7.34) and (7.35) we define

$$\begin{aligned} \mathcal{Y}_{\gamma_{++'}} &:= \frac{s_3(z) \otimes s_1(z')}{(s_1, s_3)} \\ \mathcal{Y}_{\gamma_{--'}} &:= \frac{s_1(z) \otimes s_3(z')}{(s_3, s_1)} \end{aligned} \tag{8.5}$$

because for $z, z' \in C_{3,1}$ the $+$ branch of λ is associated with z_1 and the $-$ branch with z_3. (See Figure 20.) Therefore in case (A) of Figure 19 we can write

$$\langle L_{\wp,\zeta} \rangle = \mathcal{Y}_{\gamma_{++'}} + \mathcal{Y}_{\gamma_{--'}}. \tag{8.6}$$

There are therefore two framed BPS states associated to the open paths indicated above.

Now let us increase the phase of ζ so that the WKB curves rotate to case (B) of Figure 19. Now z remains in $C_{3,1}$, but $z' \in C_{1,2}$, and given our rules for defining $\mathcal{X}_{n,c,d}$ we must use

$c \in \{1,3\}$ and $d \in \{1,2\}$. Thus, none of the expressions (8.4) is directly useful. However, we can use the relation

$$s_1(s_2,s_3) + s_2(s_3,s_1) + s_3(s_1,s_2) = 0 \tag{8.7}$$

to eliminate $s_3(z')$ in (8.4a) in favor of $s_1(z'), s_2(z')$ to get [48]

$$\begin{aligned} \langle L_{\wp,\zeta} \rangle &= \frac{s_3(z) \otimes s_1(z')}{(s_1,s_3)} + \frac{(s_2,s_3)}{(s_1,s_2)(s_1,s_3)} s_1(z) \otimes s_1(z') + \frac{s_1(z) \otimes s_2(z')}{(s_2,s_1)} \\ &= \mathcal{Y}_{\gamma_{++'}} - \mathcal{Y}_{\gamma_{-+'}} + \mathcal{Y}_{\gamma_{--'}} \end{aligned} \tag{8.8}$$

The path $\gamma_{--'}$ is the same as that used in (8.5), although its expression in terms of the s_a's has changed. Also note that a third term, similar to (7.36), has emerged, so we now have three framed BPS states. This is an example of the wall-crossing phenomenon for framed BPS states.

Now we can check the framed wall-crossing formula. We have

$$\mathcal{Y}_{\gamma_{++'}} - \mathcal{Y}_{\gamma_{-+'}} + \mathcal{Y}_{\gamma_{--'}} = (\mathcal{Y}_{\gamma_{++'}} + \mathcal{Y}_{\gamma_{--'}})(1 + \mathcal{Y}_{\gamma_{-'+'}}) \tag{8.9}$$

which is the expected wall-crossing transformation by $\mathcal{S}^{\mu}$.

If we further increase the phase of ζ so that we come to case (C) of Figure 19 then we use (8.4b) and recover once again (8.6). Now the wall-crossing is given by

$$(1 - \mathcal{Y}_{\gamma_{-+}})(\mathcal{Y}_{\gamma_{++'}} - \mathcal{Y}_{\gamma_{-+'}} + \mathcal{Y}_{\gamma_{--'}}) = \mathcal{Y}_{\gamma_{++'}} + \mathcal{Y}_{\gamma_{--'}} \tag{8.10}$$

which again is the expected transformation by $\mathcal{S}^{\mu}$.



Figure 21: When z and z' are separated by an angle between $2\pi/3$ and π there are six different Darboux expansions (up to monodromy).

Let us now briefly indicate what happens when z and z' are separated by an angle between $2\pi/3$ and π, as in Figure 21. The Darboux expansion for case (A) is

$$\begin{aligned} \langle L_{\wp,\zeta} \rangle &= \frac{s_1(z) \otimes s_2(z')}{(s_2,s_1)} + \frac{(s_2,s_3)}{(s_1,s_2)(s_1,s_3)} s_1(z) \otimes s_1(z') + \frac{s_3(z) \otimes s_1(z')}{(s_1,s_3)} \\ &= \mathcal{Y}_{\gamma_{--'}} + \mathcal{Y}_{\gamma_{-+'}} + \mathcal{Y}_{\gamma_{++'}} \end{aligned} \tag{8.11}$$

---

[48] One can show that consistency of the wall-crossing formulae given below fixes all signs up to the choices of $\sigma(+-,-+')$ and $\sigma(+-,--')$. We will make the choice $\sigma(+-,-+') = +1$ and $\sigma(+-,--') = -1$, which simplifies the formulae.

Denoting this expansion by $\mathcal{E}_A$, etc. the wall-crossings are

$$\begin{aligned}(1-\mathcal{Y}_{+-})\mathcal{E}_A &= \mathcal{E}_B = \mathcal{Y}_{--'} + \mathcal{Y}_{+-'} + \mathcal{Y}_{-+'} \\ \mathcal{E}_B(1+\mathcal{Y}_{-'+'}) &= \mathcal{E}_C = \mathcal{Y}_{--'} + \mathcal{Y}_{+-'} + \mathcal{Y}_{++'} \\ (1-\mathcal{Y}_{-+})\mathcal{E}_C &= \mathcal{E}_D = \mathcal{Y}_{+-'} + \mathcal{Y}_{++'} + \mathcal{Y}_{-+'} \\ \mathcal{E}_D(1+\mathcal{Y}_{+'-'}) &= \mathcal{E}_E = \mathcal{Y}_{++'} + \mathcal{Y}_{-+'} + \mathcal{Y}_{--'} \\ (1-\mathcal{Y}_{+-})\mathcal{E}_E &= \mathcal{E}_F = \mathcal{Y}_{-+'} + \mathcal{Y}_{--'} + \mathcal{Y}_{+-'} \\ \mathcal{E}_F(1+\mathcal{Y}_{-'+'}) &= M \cdot \mathcal{E}_A = \mathcal{Y}_{--'} + \mathcal{Y}_{+-'} + \mathcal{Y}_{++'}\end{aligned} \tag{8.12}$$

In the last line M is the monodromy operator and for brevity we have shortened the notation to $\mathcal{Y}_{\gamma_{++'}} = \mathcal{Y}_{++'}$, etc.

As a final remark note that since the bulk theory is so trivial, and in particular since there is no degree of freedom charged under the $\mathbb{Z}_2$ flavor monodromy for $\mathbb{S}_z$, we should be able to consider boundaries for the surface defect. The small flat sections $\pm s_a(z)$ are perfectly good candidates for the vevs of such boundary line defects. Because they can be canonically normalized (up to sign) by the condition $(s_a, s_{a+1}) = 1$, they are actual holomorphic sections of $V_{\mathbb{S}}$. When z is in an appropriate cell, they will define $\mathcal{Y}_+$, $\mathcal{Y}_-$, $-\mathcal{Y}_+ - \mathcal{Y}_-$.

8.1.2 $N = 2$

We next consider the case $N = 2$, where the polynomial P is of the form $P_2(z) = z^2 + 2m$. The IR dynamics are still trivial, so there are no gauge fields and no gauge charges, $\Gamma_g = 0$. The Coulomb branch $\mathcal{B}$ is just a single point, so there is no possibility of wall-crossing for the 4d BPS spectrum. However, there is a 1-dimensional flavor charge lattice $\Gamma_f = \Gamma$, and the 4d BPS spectrum is nonempty — there is a single BPS particle, carrying a flavor charge γ, corresponding to a finite WKB curve which connects the two zeroes of P. So $\Omega(\pm\gamma) = 1$, while all $\Omega(n\gamma) = 0$ for $n \neq \pm 1$.

The story becomes more interesting when we introduce the canonical surface defect $\mathbb{S}_z$. Then we get a local model for the general behavior of a 2d-4d theory with two vacua, a light bulk particle and light 2d particles. (See the end of this section for further remarks on how to interpret the model.)

Unlike the $N = 1$ example where the 2d BPS spectrum was independent of z, we now find four distinct regions in the z-plane, with different 2d BPS spectra. Indeed, we can use the general principle mentioned at the end of §7.2. The regions are cut out by the finite and separating WKB curves emanating from the two turning points, at the value of $\vartheta = \vartheta_*$ for which the finite WKB curve is present; see Figure 22. Recall that when z lies on one of these curves, there is an obvious finite open WKB network from z to the turning point, with phase ϑ_*. This finite WKB network corresponds to a 2d BPS particle whose phase is ϑ_*. But ϑ_* is also the phase of the 4d BPS particle; so for this value of z the central charges of the 2d and the 4d particle are aligned. Thus, the finite and separating WKB curves of phase ϑ_* shown in Figure 22 are the walls of marginal stability in the parameter z.



Figure 22: The cell decomposition induced by the WKB foliation $F_{\mathrm{WKB}}(\vartheta)$ in the $N=2$ Argyres-Douglas-type theory, when ϑ is set to the critical value $\vartheta_* = \arg Z_\gamma$. The WKB curves shown here are also the walls of marginal stability for the parameter z of the canonical surface defect. We label the regions R, U, L, D. Here m is real and negative.



Figure 23: When $\vartheta \neq \vartheta_*, \vartheta_* + \pi$ the separating WKB curves look as shown. As ϑ increases the three prongs around each turning point both rotate with the same sense. As ϑ crosses ϑ_* the curve of type S_1 jumps to a curve of type S_4 and the curve of type S_2 jumps to a curve of type S_3.

The spectrum μ of 2d solitons can be determined geometrically as follows. In the regions L and R which have a single turning point on their boundary, the 2d spectrum consists of a single soliton (and its antiparticle) between the two vacua: it corresponds to a single finite WKB curve running from z to the turning point at the boundary of the region. In the regions U and D which have two turning points on the boundary, the 2d spectrum consists of two 2d particles, corresponding to two finite WKB curves from z to the two turning points on the boundary. This can be proved as follows. For $\vartheta \neq \vartheta_*, \vartheta_* + \pi$ the separating WKB curves look like those shown in Figure 23. As ϑ rotates the curves rotate around the turning points. The region between S_1 and S_2 or between S_3 and S_4 rotates and fills the upper and lower regions of Figure 22. For any point in these regions there will be values of ϑ such that that point lies on each of the curves of type S_1 and S_3 or S_2 and S_4. Because the critical separating curves (i.e. those at $\vartheta = \vartheta_*$) are walls of marginal stability we know that the curves S_i cannot enter into the left and right regions. Similarly, for any point in the left region there will be some value of ϑ such that this point lies on a separating curve of type S_5 or S_6.



Figure 24: A choice of cycles so that $\omega(\gamma, \gamma_{ij}(z)) = -1$. Here we have taken $P_2 = z^2 - \alpha^2$ where α is a positive real number, and we have placed the canonical surface operator $\mathbb{S}_z$ on the ray (α, ∞).

In order to illustrate the wall-crossing formulae explicitly we now describe the 2d soliton spectrum in some detail. Our technique will be to determine the periods in some easily accessible region and then derive the spectrum in the remaining regions using wall-crossing. Because the local system Γ_{12} has monodromy around the two roots of P_2 we will find different spectra when comparing using homotopically different paths.

We begin with z in the region R and call the charge of the single 2d particle $\gamma_{12}(z)$, or sometimes just γ_{12} with z understood. In order for this to be unambiguous we should choose cuts for the double cover $\Sigma \to C$. It is convenient to define $2m = -\alpha^2$ and write $\sqrt{P} = \sqrt{z+\alpha}\sqrt{z-\alpha}$, choosing the principal branch of the logarithm to define each factor. When α is real $\sqrt{P} \sim z$ at large z and has a cut running along the segment $[-\alpha, \alpha]$. For definiteness we take $\alpha > 0$. We choose orientations as in Figure 24 so that $\omega(\gamma, \gamma_{12}) = -1$. With these choices we can determine the local system Γ_{12}. A simple path taking z around the root α leads to monodromy $\gamma_{12}(z) \to -\gamma_{12}(z) =: \gamma_{21}(z)$, and a simple path taking z around the root $-\alpha$ leads to monodromy $\gamma_{12}(z) \to -\gamma_{12}(z) + 2\gamma$. This follows most easily by noting that $Z_{\gamma_{12}(z)-\gamma}$ has a zero at $z = -\alpha$ and has a square-root branch cut there.

It is also useful to note that for z large and positive the WKB curve clearly runs along the positive axis so $Z_{\gamma_{12}(z)}$ is positive. For large $|z|$ it behaves like $Z_{\gamma_{12}(z)} \sim \pi^{-1} z^2$. (In fact, in this extremely simple model we can write the periods explicitly:

$$\begin{aligned} Z_\gamma &= -\mathrm{i}\alpha^2, \\ Z_{\gamma_{12}(z)} &= \frac{1}{\pi}\left(z\sqrt{z^2-\alpha^2} - \alpha^2 \log\left(z + \sqrt{z^2-\alpha^2}\right) + \alpha^2 \log \alpha\right) \end{aligned} \tag{8.13}$$

but we will phrase our arguments so that such explicit formulae are not needed, since in general explicit formulae for periods are not available.)

Using the above description of the periods, we can form the product $A(\sphericalangle)$, taking $\sphericalangle$ a sector of opening angle 2π, i.e. the whole plane:

$$A(\sphericalangle; R) = \mathcal{K}_\gamma \mathcal{S}_{\gamma_{12}} \mathcal{K}_{-\gamma} \mathcal{S}_{-\gamma_{12}} \tag{8.14}$$

Although we have derived it for m real and negative and for z large and positive, this will hold for all z in the region R. Moreover, changing the phase of m merely rotates the whole picture. (We will return to the wall-crossing in m at the end of this section.)



Figure 25: The central charges for z in the region R on the real axis, and $m \in \mathbb{R}_-$. If m is fixed and z is continued into the region U, then $Z_{\gamma_{12}(z)}$ rotates counterclockwise, and aligns with $Z_{-\gamma}$ at the $R \to U$ wall of marginal stability. If z is continued into the region D then the central charge $Z_{\gamma_{12}(z)}$ rotates clockwise, and aligns with Z_γ at the $R \to D$ wall of marginal stability.

Before proceeding let us make a few comments about the notation (8.14). First, in general, products of wall-crossing factors are only expected to make sense for $\sphericalangle$ of opening less than π. However, in the AD examples the product is always finite, so the full 2π product is sensible. Moreover, it is sometimes useful to write the product for the entire range 2π since one might wish to split it into different half-planes. Of course, one can cycle the factors in these expressions. Second, the spectrum does not depend just on the region R, U, L, D but rather on a homotopy class of paths used to continue from R to the region. We will suppress this in our simple discussion below. Third, the expression (8.14) summarizes the configuration of central charges of occupied charges shown in Figure 25 above. We will not draw the analogous figures for the other regions; rather, the reader is urged to draw pictures of the central charges in order to make the subsequent formulae comprehensible.

We are now ready to proceed with the wall-crossing analysis. We will first take $|z|$ large and move counterclockwise through regions $R \to U \to L \to D$. Then we will compare with moving clockwise from $R \to D$, as well as moving z through the cut $[-\alpha, \alpha]$ directly from $U \to D$.

If we first move z from R to U then $Z_{\gamma_{12}(z)}$ becomes parallel to $Z_{-\gamma}$. (This can be easily seen by making $|z|$ large. Then the marginal stability wall is at $\arg z \approx \frac{\pi}{4}$ and $Z_{\gamma_{12}} \sim \pi^{-1} z^2$ has argument approximately $\pi/2$.) We can therefore apply (8.2b) with $a = -\gamma$ and $b = \gamma_{12}(z)$ to produce the spectrum in region R:

$$A(\sphericalangle; U) = \mathcal{K}_\gamma \mathcal{S}_{\gamma_{21}+\gamma} \mathcal{S}_{\gamma_{21}} \mathcal{K}_{-\gamma} \mathcal{S}_{\gamma_{12}-\gamma} \mathcal{S}_{\gamma_{12}} \tag{8.15}$$

If we continue to increase the phase of z so that we encounter the wall-crossing $U \to L$ then the periods continue rotating and at the $U \to L$ wall $Z_{\gamma_{12}}$ becomes parallel to Z_γ. We now apply (8.2a) with $a = \gamma$ and $b = \gamma_{12} - \gamma$ to produce the spectrum in L:

$$A(\lhd; L) = \mathcal{K}_{\gamma} \mathcal{S}_{\gamma_{12}-\gamma} \mathcal{K}_{-\gamma} \mathcal{S}_{\gamma_{21}+\gamma}. \tag{8.16}$$

Continuing to increase the phase of z at the $L \to D$ wall $\arg z \approx \frac{5\pi}{4}$ and $Z_{\gamma_{12}-\gamma}$ becomes parallel to $Z_{-\gamma}$. We now apply (8.2b) to produce

$$A(\lhd; Dl) = \mathcal{K}_{\gamma} \mathcal{S}_{\gamma_{21}+2\gamma} \mathcal{S}_{\gamma_{21}+\gamma} \mathcal{K}_{-\gamma} \mathcal{S}_{\gamma_{12}-2\gamma} \mathcal{S}_{\gamma_{12}-\gamma} \tag{8.17}$$

We write Dl to indicate that this is the spectrum obtained from continuation from the region L. (Recall that $A(\lhd)$ actually depends on a homotopy class of paths from region R.)

Going back to the region R we could instead have continued z into the D region by moving z clockwise. In this case $Z_{\gamma_{12}(z)}$ rotates clockwise and at the marginal stability wall becomes parallel to Z_{γ} and (8.2a) produces

$$A(\lhd; Dr) = \mathcal{K}_{\gamma} \mathcal{S}_{\gamma_{21}} \mathcal{S}_{\gamma_{21}-\gamma} \mathcal{K}_{-\gamma} \mathcal{S}_{\gamma_{12}} \mathcal{S}_{\gamma_{12}+\gamma} \tag{8.18}$$



Figure 26: The soliton spectrum for the $N = 2$ AD theory. We have chosen two cuts, on the complement of which the torsor Γ_{12} can be trivialized. These cuts divide the "down region" D into three subregions. The spectra in these three regions are related by the monodromy transformations associated with the two turning points. We have only given half the spectrum; the other half consists of the antiparticles.

Finally, we could start with the spectrum summarized by $A(\lhd; U)$ in (8.15) and continue z directly into the region D. Note that when $z \in (-\alpha, \alpha)$ the period $Z_{\gamma_{12}}$ is pure imaginary. Moreover, expanding the period around $z = \alpha$ say, $z = \alpha + w$, with w small we find $Z_{\gamma_{12}} \sim w\sqrt{2\alpha w}$. From this it follows that as z goes through the marginal stability line $U \to D$ the central charges $Z_{\gamma_{12}}$ and $Z_{\gamma_{21}+\gamma}$ simultaneously align with Z_{γ}. Thus we should apply the wall-crossing formula (8.3) to obtain:

$$A(\lhd; Dc) = \mathcal{K}_{\gamma} \mathcal{S}_{\gamma_{12}} \mathcal{S}_{\gamma_{12}-\gamma} \mathcal{K}_{-\gamma} \mathcal{S}_{\gamma_{21}} \mathcal{S}_{\gamma_{21}+\gamma}. \tag{8.19}$$

In this way we arrive at the full spectrum shown in Figure 26. The fact that we found three different spectra in the region D is the result of the monodromy of the local system Γ_{12}

around the two zeroes of P. As we have seen, when comparing $Dl \to Dc$ we should take $\gamma_{12} \to \gamma_{21} + 2\gamma$ (since $Z_{\gamma_{12}-\gamma}$ vanishes at the left turning point), while to compare $Dr \to Dc$ we should take $\gamma_{12} \to \gamma_{21}$. It is easy to check that with these monodromy transformations the spectrum is nicely consistent.



Figure 27: Walls of marginal stability in the m-plane for fixed z. The wall emanating to the left is of type $\mathcal{SKS} = \mathcal{SKS}$, while the two walls emanating from $m = 0$ are of type $\mathcal{KS} = \mathcal{SSK}$ and $\mathcal{SK} = \mathcal{KSS}$ respectively. There is a branch cut emanating from $m = 0$. When $z \to 0$ the 4d and 2d periods Z_γ and $Z_{\gamma_{12}}$ simultaneously vanish. (This corresponds to a singularity in the semiflat geometry, which is resolved by quantum effects; the resolved structure looks like a hyperholomorphic bundle on periodic NUT space.)

Finally, it is interesting to study the behavior of the model in the m plane, since it serves as a useful local model for the general 2d-4d wall-crossing structure near a locus of $\mathcal{B}$ where $Z_\gamma = Z_{\gamma_{ij}} = 0$. Working at fixed z, there are two singularities in the m plane: $m = 0$, where $Z_\gamma = 0$, and $m = m_0 = -\frac{z^2}{2}$, where $Z_{\gamma_{12}} = 0$. Rotating the phase of m, the pattern of walls in Figure 22 rotates, so that if m rotates by 2π then the figure rotates in the same sense by π. When $|m| > \frac{1}{2}|z^2|$ the segment between the two roots of P_2 rotates through z and we apply the wall-crossing formula (8.3). When $|m| < \frac{1}{2}|z^2|$ as the phase of m increases by 2π two of the separating WKB curves pass through z. As we have seen, these correspond to wall-crossings of type (8.2a) and (8.2b). Hence at fixed z we will find two walls of marginal stability emanating from $m = 0$, and the three types of walls meet at m_0. The m-plane is thus divided into two regions, an inner region with a single 2d soliton, and an outer region with two 2d solitons. The two regions are separated by a closed wall of marginal stability running through both singularities. Another wall, of the $\mathcal{SKS} \to \mathcal{SKS}$ type goes from m_0 all the way to infinity. See Figure 27.

Remarks

1. The example of this section is actually a member of a larger family of local models,

$$P_N(x) = zQ_{N_f}(x) \tag{8.20}$$

with $N > N_f$, which correspond to theories with N_f flavor charges only, no gauge charges, and describe the general behavior of 2d-4d A_n theories near loci where a certain number of mutually local bulk particles and domain walls between N vacua become simultaneously light. Indeed $x^2 = z^2 + 2m$ is equivalent to $x^2 - 2m = 2zx$. It would be interesting to study this general class, but we will leave it for future work.

2. Although we said that a bulk particle of flavor charge γ is present, this is rather immaterial, unless one gauges that flavor symmetry, as would happen if this were a local model for a larger theory. Rather, what matters is that there will be non-zero $\omega(\gamma, \gamma_{ij})$. It is perfectly fine to interpret the results of this subsection as concerning the behavior of a certain 2d theory with 2 vacua, with 2d particles carrying a flavor charge γ. It should not be difficult to engineer a theory, with the correct twisted chiral ring relations as to reproduce the spectral curve of this Hitchin system. We can get pretty close if we consider a $U(1)$ gauged linear sigma model, with a chiral multiplet of charge 1 and a chiral multiplet of charge -2. Integrating away the massive chiral multiplets gives us a twisted superpotential

   $$2\sigma \log \sigma - (\sigma + m) \log(\sigma + m) - 2\mathrm{i}\pi t\sigma \tag{8.21}$$

   which is minimized if $\sigma^2 = e^{2\pi \mathrm{i} t}(\sigma + m)$: if we were to set $z = e^{\mathrm{i}\pi t}$ and $\sigma = xz$, so that $2\mathrm{i}\pi\sigma \mathrm{d}t \sim x\mathrm{d}z$ then we get the desired curve. It would be interesting to compare the BPS spectrum of this theory with the one we discuss below.

3. We have not systematically investigated the framed BPS degeneracies in this model, but we have given all the techniques needed to do so.

8.1.3 Larger N

At larger N, the picture becomes more intricate, but no essentially new wall-crossing phenomena happen: all wall-crossings involving 2d particles can be understood using the basic identities (8.2a), (8.2b), and (8.3) we used above.

Let us describe a sample of the behavior one finds. We begin from a region of the Coulomb branch $\mathcal{B}$ where all roots of $P(z)$ are real. Here the 4d spectrum is easily described [69]: finite WKB curves appear only between adjacent roots. We denote their charges as γ^s, $s = 1, \ldots, N-1$, where $s = 1$ corresponds to the *rightmost* root. The intersection products are $\langle \gamma^s, \gamma^{s-1} \rangle = 1$ and $\langle \gamma^s, \gamma^t \rangle = 0$ if $|s-t| > 1$. The corresponding BPS rays ℓ_{γ^s} all lie either along the real or the imaginary ζ axis, for even and odd s respectively. See, for example, Figure 28 for the case $N = 6$. (We have here a rather degenerate situation, where several BPS rays of different nonintersecting charges coincide. Nevertheless, this is not really a wall of marginal stability since the corresponding charges have zero symplectic product with one another.)

Now how about the spectrum with the surface defect $\mathbb{S}_z$ included? If z is well to the right of all the roots, this spectrum just consists of a single 2d particle, corresponding to a finite WKB curve running from z to the rightmost root. Call its charge γ_{12}. So we have an additional BPS ray $\ell_{\gamma_{12}}$ carrying the transformation $\mathcal{S}_{\gamma_{12}}$. Furthermore there are no other open finite WKB networks starting from z. In order to see this we use reasoning similar to that used near Figure 23. At $\vartheta = \vartheta_*$ which supports a 4d BPS state the walls will look like those in Figure 28. For $\vartheta \neq \vartheta_*, \vartheta_* + \pi$, the separating WKB curves will rotate but there will always be a region far to the right containing the large positive real axis and sitting between the rightmost separating WKB curves. Now, WKB rays can never cross, and therefore, if z is a large positive number then for any ϑ there can be at most one open



Figure 28: The walls of marginal stability in C for 2d-4d BPS states in the $N = 6$ Argyres-Douglas theory, when u is in the region of the Coulomb branch where all roots of P_6 are real. Finite WKB curves join successive roots. The separating WKB curves asymptote to the fourteen lines of phase $2\pi n/16$ not parallel to the x-axis. The red walls correspond to wall-crossings involving BPS rays with phase $\vartheta = \pm\pi/2$ and the blue walls correspond to those rays with $\vartheta = 0, \pi$. We have shown $\gamma_{12}(z)$, γ^1 and γ^2 in some cut system, and indicated the $\mathcal{K}$-rays which become parallel with the $\mathcal{S}$-ray $\ell_{\gamma_{12}(z)}$ for the first few walls.

finite WKB network. On the other hand, the walls of marginal stability are determined by the pattern at ϑ_*. Therefore, if z is any point in the rightmost region of Figure 28 there is one and only one open finite WKB network connecting z to the turning point z_1. Moreover, $\omega(\gamma^s, \gamma_{12}) = 0$ for all s, except for $s = 1$, for which $\omega(\gamma^1, \gamma_{12}) = 1$.


$Z_{\gamma^1}, Z_{-\gamma^3}, \ldots$
$Z_{\gamma^2}, Z_{-\gamma^4}, \ldots$
$Z_{-\gamma^2}, Z_{\gamma^4}, \ldots$
$Z_{\gamma_{12}+\gamma^1+\gamma^2}$
$Z_{\gamma_{12}+\gamma^1}$
$Z_{\gamma_{12}}$
$Z_{-\gamma^1}, Z_{\gamma^3}, \ldots$

Figure 29: When z is large and lies in the region in the upper half plane between the separating WKB curves associated to γ^2 and $-\gamma^1$, the soliton spectrum is that shown here.

Let us consider how the configuration of BPS rays evolves as z is varied. As we move z counterclockwise around the configuration of roots of P, staying a large distance away, the single $\mathcal{S}_{\gamma_{12}}$ ray first meets a BPS ray carrying the transformation $\mathcal{K}_{\gamma^1}$ (and also carrying various other $\mathcal{K}$ transformations corresponding to other 4d BPS states, but those transformations commute with $\mathcal{S}_{\gamma_{12}}$, so they play no role at this moment.) Applying the WCF (8.2b), one sees that after these two rays cross, the $\mathcal{S}_{\gamma_{12}}$ ray is replaced by two rays, carrying $\mathcal{S}_{\gamma_{12}}$ and $\mathcal{S}_{\gamma_{12}+\gamma^1}$. As we continue to move z, these two rays next cross $\mathcal{K}_{\gamma^2}$. Since γ_{12} has zero intersection with $\tilde{\gamma}_2$ there is no wall-crossing as the ray for γ_{12} passes through, but this is not so for $\gamma_{12}+\gamma^1$. Indeed, using again (8.2b) with $a=\gamma^2$ and $b=\gamma_{12}+\gamma^1$ wall-crossing produces three rays, $\mathcal{S}_{\gamma_{12}}$, $\mathcal{S}_{\gamma_{12}+\gamma^1}$, $\mathcal{S}_{\gamma_{12}+\gamma^1+\gamma^2}$, leading to the configuration of rays shown in Figure 29. Next, z crosses a ray for $-\gamma^1$ and γ^3. In our degenerate situation with large z these happen almost simultaneously. First the rays for $\gamma_{12}(z)+\gamma^1$ and $\gamma_{12}(z)$ cross the ray with factor $\mathcal{K}_{-\gamma^1}$ and we apply the wall-crossing formula (8.2a):

$$\mathcal{S}_{\gamma_{12}+\gamma^1+\gamma^2}\mathcal{S}_{\gamma_{12}+\gamma^1}\mathcal{S}_{\gamma_{12}}\mathcal{K}_{-\gamma^1}\mathcal{K}_{\gamma^3}=\mathcal{S}_{\gamma_{12}+\gamma^1+\gamma^2}\mathcal{K}_{-\gamma^1}\mathcal{S}_{\gamma_{12}+\gamma^1}\mathcal{K}_{\gamma^3}. \tag{8.22}$$

Next, since $\omega(-\gamma^1,\gamma_{12}(z)+\gamma^1+\gamma^2)=0$, when the ray with charge $\gamma_{12}+\gamma^1+\gamma^2$ passes through $\mathcal{K}_{-\gamma^1}$ we simply commute factors:

$$\mathcal{S}_{\gamma_{12}+\gamma^1+\gamma^2}\mathcal{K}_{-\gamma^1}\mathcal{S}_{\gamma_{12}+\gamma^1}\mathcal{K}_{\gamma^3}=\mathcal{K}_{-\gamma^1}\mathcal{S}_{\gamma_{12}+\gamma^1+\gamma^2}\mathcal{S}_{\gamma_{12}+\gamma^1}\mathcal{K}_{\gamma^3}. \tag{8.23}$$

Next as z crosses the separating line for γ^3, $\mathcal{S}_{\gamma_{12}+\gamma^1}$ commutes through (again because intersection products vanish) and we apply (8.2b) to produce

$$\mathcal{K}_{-\gamma^1}\mathcal{S}_{\gamma_{12}+\gamma^1+\gamma^2}\mathcal{S}_{\gamma_{12}+\gamma^1}\mathcal{K}_{\gamma^3}=\mathcal{K}_{-\gamma^1}\mathcal{K}_{\gamma^3}\mathcal{S}_{\gamma_{12}+\gamma^1+\gamma^2+\gamma^3}\mathcal{S}_{\gamma_{12}+\gamma^1+\gamma^2}\mathcal{S}_{\gamma_{12}+\gamma^1}. \tag{8.24}$$

Thus the 2d particle of charge γ_{12} decays into a 2d particle of charge $\gamma_{12}+\gamma^1$ plus a 4d particle of charge $-\gamma^1$, and almost simultaneously a new 2d particle of charge $\gamma_{12}+\gamma^1+\gamma^2+\gamma^3$ is formed as a bound state of a 2d $\gamma_{12}+\gamma^1+\gamma^2$ and a 4d γ^3. The pattern continues as z moves across the upper half of the plane: we encounter a series of pairs of walls at which one state decays and a new one is born. At the last two walls we have only decays with no new states created, so when z reaches a point far to the left of all of the roots of P, we find just a single BPS state. We can then bring z back around the circle to its original position: we encounter a sequence of walls very similar to what we just described. See Figure 28 for a picture of the walls in the z-plane in the case $N=6$. As in the discussion of $N=2$ above, when z comes back to the original point we must have only the original soliton, but to see this explicitly as the outcome of a composition of wall-crossings requires us to take account of the monodromy of Γ_{12} around the loop. We emphasize that this is only the picture at one specific locus in the Coulomb branch; as we vary $u\in\mathcal{B}$ the walls would deform and even change topology. One could encounter more complicated patterns of wall-crossing in these cases. As a simple illustration of this we plot the walls of marginal stability for the $N=6$ theory for $\lambda^2=(z^6-1)(\mathrm{d}z)^2$ in Figure 30. Still, we believe that one could study all the wall-crossing phenomena systematically using the same basic wall-crossing formulas we used above.

It is worthwhile discussing the case $N=3$ a bit further. In this example the Coulomb branch $\mathcal{B}$ is one-dimensional and has a simple singularity structure. We take the polynomial $P_3(z)=z^3-3\Lambda^2 z+u$. The Coulomb branch for the bulk theory has two singular loci $u_\pm$,



Figure 30: Walls of marginal stability for the $N = 6$ AD theory at a particular point u in $\mathcal{B}$ as described in the text.

where an "electron" and a "monopole", of charges γ_e and γ_m, $\langle \gamma_e, \gamma_m \rangle = 1$, are respectively massless. There is a single wall of marginal stability, separating an inner region where the spectrum consists of the two particles only, and an outer region where the spectrum consists of three particles, of charges γ_e, γ_m, $\gamma_e + \gamma_m$. Notice that the monodromy at infinity which arises from the combination of the monodromies around the two singular points is $(\gamma_e, \gamma_m, \gamma_e + \gamma_m) \to (\gamma_e - \gamma_m, \gamma_m, \gamma_e) \to (-\gamma_m, \gamma_m + \gamma_e, \gamma_e)$, a cyclic permutation of the three particles.

If we set $\Lambda = 0$ the two singular loci coalesce into a AD point, the inner region disappears, and the setup has an exact $\mathbb{Z}_3$ symmetry acting on the z plane which coincides with the cyclic permutation of the populated charges γ_e, γ_m, $\gamma_e + \gamma_m$. If we set the surface defect parameter at the origin, $z = 0$, the spectrum of 2d solitons must also enjoy this symmetry. Indeed, it is not difficult to argue that there are three 2d solitons, whose charges correspond to three straight WKB segments from $z = 0$ to the three turning points $z = u^{1/3}$. Even if $\Lambda \neq 0$, this is the general spectrum for sufficiently large u. It is not difficult to fill in the full spectrum for large $|z|$ as a function of u, interpolating from very large to small $|u|$, but we will not describe the details here. We simply note that the three 2d solitons at very large $|u|$ can be continuously connected to the soliton of charges γ_{ij}, $\gamma_{ij} + \gamma_e$, $\gamma_{ij} + \gamma_e + \gamma_m$ which we encountered at the third step of the general N analysis.

8.2 The $\mathbb{CP}^1$ sigma model

Now let us consider another example where the 4d dynamics is trivial, namely the 2d $\mathbb{CP}^1$ sigma model. This model nicely illustrates the examples related to degenerate cells and pop transitions discussed in §7.5. It is also useful preparation for the rather more challenging example of the four-dimensional $SU(2)$ $N_f = 0$ theory coupled to the the $\mathbb{CP}^1$ sigma model, to be discussed in §8.3.

The BPS spectrum in this model has been studied by Dorey [21]. Let us briefly recall the salient results. The chiral ring with twisted mass parameter m is

$$x^2 = \Lambda^2 e^t + m^2, \tag{8.25}$$

where t is the Kähler parameter of the target $\mathbb{CP}^1$. The twisted mass defines two vacua which we can take to be the north and south poles of the target space $\mathbb{CP}^1$. This mass breaks the $SU(2)$ global symmetry (coming from the isometries of the target space) to $U(1)$. We write the root lattice of $SU(2)$ as $\mathbb{Z}\alpha$ and measure $U(1)$ charges in terms of α. In the strong coupling region, defined by $|m^2| \ll |\Lambda^2 e^t|$, the effect of the twisted mass is negligible and there are two solitons interpolating between the vacua. At $m = 0$ they form a doublet and hence have $U(1)$ charges $\pm\frac{1}{2}\alpha$. On the other hand, in the weak coupling region, defined by $|m^2| \gg |\Lambda^2 e^t|$, the twisted mass localizes the dynamics to one of the two vacua at the north or south poles of $\mathbb{CP}^1$. The massive sigma-model fluctuations around those vacua give 2 BPS particles of central charge $Z = \mathrm{i}m$ together with their antiparticles. There is also an infinite tower of semi-classical solitons interpolating between the two vacua. These solitons (and their antiparticles) carry global $U(1)$ charge $(n + \frac{1}{2})\alpha$ for all integers $n \in \mathbb{Z}$, with degeneracy $\mu = 1$ for each n. Evidently, there is a marginal stability transformation in the BPS spectrum, very reminiscent of the strong-weak coupling transition in the four-dimensional $SU(2)$ theory [21, 23].

It is instructive to see how these results are reproduced in the geometric formulation of this paper. We take $C = \mathbb{CP}^1 - \{0, \infty\}$. ($C$ is not related to the $\mathbb{CP}^1$ target of the sigma model!) The chiral ring corresponds to the spectral cover equation

$$\lambda^2 = \left(\frac{\Lambda^2}{z} + \frac{m^2}{z^2}\right) \mathrm{d}z^2 \tag{8.26}$$

where $z = e^t$ and $\lambda = x\,\mathrm{d}t$. This covering is the spectral cover of an A_1 Hitchin system on C with the weakest possible irregular singularity at infinity, and a regular singularity with residue $m\sigma^3$ at the origin. The Hitchin equations in this system coincide with the tt^* equations for the $\mathbb{CP}^1$ sigma model. The double covering has two branch points: one at $z = z_0 = -\frac{m^2}{\Lambda^2}$ and one at $z = \infty$, so Σ is a thrice-punctured sphere. Let $P^\pm$ be the two lifts of $z = 0$. The homology of Σ is generated by γ, a small curve winding once around P^+, and $\sigma^*(\gamma)$, a small curve winding once around P^-. The class $\gamma_f := \gamma - \sigma^*(\gamma)$ generates the odd homology lattice $\Gamma = H_1(\Sigma; \mathbb{Z})^-$ and corresponds to a closed curve around $z = \infty$. We will identify Γ with the flavor lattice $\alpha\mathbb{Z}$ and hence α with γ_f.

Let us now derive the BPS spectrum geometrically. As in the previous examples this can be inferred from the structure of the WKB foliations, shown in this case in Figure 31. At the critical values $\vartheta = \vartheta_*, \vartheta_* + \pi$ given by $e^{\mathrm{i}\vartheta_*} = \pm\mathrm{i}\frac{m}{|m|}$ there is a finite WKB curve joining z_0 to itself, as in the middle of Figure 31. As we have explained above the finite and separating WKB curves at this critical value of ϑ are the marginal stability lines in z. The finite WKB curve at ϑ_* separates C into two regions, with the inner region near $z = 0$ corresponding to weak coupling and the outer region corresponding to strong coupling.

If z is any point in the strong coupling region there are precisely two special values of ϑ for which there is an open finite WKB curve connecting z to z_0. Moreover, the open paths



$\vartheta < \vartheta_*$ $\vartheta = \vartheta_*$ $\vartheta > \vartheta_*$

Figure 31: At $\vartheta = \vartheta_*$ there is a closed WKB curve connecting the turning point z_0 to itself. For $\vartheta \neq \vartheta_*$ the curves in red are separating WKB curves connecting z_0 to a singularity.

γ_{12} and $\tilde{\gamma}_{21}$ corresponding to these two curves glue together to form $\gamma_{12} + \tilde{\gamma}_{21} = \gamma_f$. These BPS states, with $\mu(\gamma_{12}) = \mu(\tilde{\gamma}_{21}) = 1$, correspond to the two strong-coupling solitons with flavor charges $\frac{1}{2}\alpha$. Together with their antiparticles we have two $SU(2)$ doublets. Note that $\tilde{\gamma}_{21} = \gamma_{21} + \gamma_f = -\gamma_{12} + \gamma_f$. Thus we can identify both Γ_{12} and Γ_{21} with the torsor $\frac{1}{2}\alpha + \Gamma$.

Now suppose z is any point in the weak coupling region. In this case the open path γ_{12} has intersection 2 with γ_f (where we choose a specific representative for γ_f, namely the difference of lifts of the closed WKB curve forming the boundary of the punctured disc.) We thus have $\omega(\gamma_f, \gamma_{12}) = 2$. Dividing this up symmetrically requires us to assign $\omega(\gamma_f, \gamma_1) = 1$ and $\omega(\gamma_f, \gamma_2) = -1$. So we find one particle of charge γ_f in each vacuum. Similarly we find one particle of charge $-\gamma_f$ in each vacuum. This fits with the field-theoretic expectation: each of the two vacua supports a particle of charge γ_f, as well as its antiparticle of charge $-\gamma_f$. Incidentally, there is another way of picturing these BPS states: rather than thinking about the intersection between γ_{12} and a closed WKB curve at the boundary of the disc, which seems a bit indirect, we observe that z itself is lying on a closed WKB curve. The two BPS particles in vacuum 1 can be naturally identified with the two possible orientations of the lift of this closed WKB curve to sheet 1 of Σ, and similarly for vacuum 2.

Moreover, for any point z in the weak coupling region there will be infinitely many values of ϑ, accumulating at the critical value ϑ_* from above and from below, such that the spiraling WKB curves join z to z_0. These correspond to the infinite tower of particles with flavor charges $\gamma_{12} + n\gamma_f$ and $\gamma_{21} + n\gamma_f$ where $n \in \mathbb{Z}$ corresponds to the winding number around the singularity at $z = 0$.

Exactly at $\vartheta = \vartheta_*$ there is a separating WKB curve joining z_0 to $z = \infty$. However, the relevant intersection product is zero, so this WKB curve does not contribute a BPS state.

We have thus reproduced the BPS spectrum of the model. Moreover, the wall-crossing in the z parameter takes place along the finite WKB curve with $e^{i\vartheta} = \pm i m/|m|$. The corresponding wall-crossing formula is in perfect correspondence with (2.51) if we translate $\gamma_{ij} \to \gamma_{12}$, $\gamma_{ji} \to \tilde{\gamma}_{21}$ and $\gamma \to \gamma_f$.

With a bit more patience, we could also explore the spectrum of framed BPS states for simple line defects in the model. We will only sketch the analysis. As we have an irregular singularity of odd degree, we can consider the associted small flat section $s(z)$, canonically normalized by the requirement $(s, Ms) = 1$. Here M is the monodromy matrix which represents parallel transport once around the cylinder. To be precise, we need to pick a path from the irregular singularity to z in order to define $s(z)$. Other choices of paths winding n more times around the cylinder will give sections $s_n(z) = M^n s(z)$ for all integer n. Notice that $(s_n, s_{n+1}) = 1$.

Standard Stokes theory always implies linear relations of the form

$$s_{n-1} + x_n s_n + s_{n+1} = 0 \tag{8.27}$$

for some scalar x_n. In this simple setup, all the x_n are equal, and

$$x_0 s = -\left(M + M^{-1}\right) s = -\left(\mathrm{Tr} M\right) s. \tag{8.28}$$

But the matrix M is just the monodromy matrix around the regular singularity, and has eigenvalues $\mathcal{Y}_{\pm\frac{1}{2}\gamma_f}$. Hence

$$x_n = x_0 = -\mathcal{Y}_{\frac{1}{2}\gamma_f} - \mathcal{Y}_{-\frac{1}{2}\gamma_f}. \tag{8.29}$$

Finally, notice that

$$M\left(Ms - \mathcal{Y}_{\pm\frac{1}{2}\gamma_f} s\right) = \mathcal{Y}_{\mp\frac{1}{2}\gamma_f} Ms - s. \tag{8.30}$$

Hence the vectors $s_{\mp} = Ms - \mathcal{Y}_{\pm\frac{1}{2}\gamma_f} s$ are monodromy eigenvectors with eigenvalue $\mathcal{Y}_{\mp\frac{1}{2}\gamma_f}$.

We can naturally conjecture that the $s_n(z,\zeta)$ are vevs of a boundary line defect L_n in the $\mathbb{CP}^1$ sigma model, i.e. a brane in the 2d sigma model. We can make an educated guess on the nature of the L_n brane: a space-filling brane (Neumann boundary conditions for the sigma model) with a Chan-Paton $U(1)$ bundle with n units of flux. A first check is the observation that a shift of the sigma model B-field $z \to e^{2\pi i} z$ sends $s_n \to s_{n+1}$, and hence it should induce a monodromy $L_n \to L_{n+1}$. But a B-field shift is indeed gauge equivalent to a shift of one unit in the Chan-Paton $U(1)$ bundle flux of all branes.

A more refined check is to derive the spectrum of framed BPS degeneracies associated to the $\mathbb{CP}^1$ sigma model on a segment, with boundaries L_0 and L_n, by expanding (s_0, s_n) recursively:

$$(s_0, s_1) = 1, \qquad (s_0, s_2) = \mathcal{Y}_{\frac{1}{2}\gamma_f} + \mathcal{Y}_{-\frac{1}{2}\gamma_f}, \qquad (s_0, s_3) = \mathcal{Y}_{\gamma_f} + 1 + \mathcal{Y}_{-\gamma_f}. \tag{8.31}$$

Generally

$$(s_0, s_n) = \frac{\mathcal{Y}_{\frac{n}{2}\gamma_f} - \mathcal{Y}_{-\frac{n}{2}\gamma_f}}{\mathcal{Y}_{\frac{1}{2}\gamma_f} - \mathcal{Y}_{-\frac{1}{2}\gamma_f}}. \tag{8.32}$$

Hence the framed BPS Hilbert space contains n states, with the quantum numbers of a irreducible representation of the $SU(2)$ flavor symmetry of the model. We can give a physical interpretation of this calculation. We can reduce the $\mathbb{CP}^1$ sigma model on the segment to the supersymmetric quantum mechanics of the zeromode, in the presence of the n units of Chan-Paton $U(1)$ bundle flux on the $\mathbb{CP}^1$ target space. It is natural for the ground states of this problem, the framed BPS states, to form an irreducible $SU(2)$ multiplet of n lowest Landau levels.

Finally, the cells of the WKB triangulation are associated to pairs of sections with good asymptotics, either of the form $(s_\pm; s_n)$ or of the form (s_n, s_{n+1}), which should be identified with the $\mathcal{Y}_{\gamma_i}$ for appropriate cycles γ_i. One can easily expand $s_{n'}(z)$ in any of those bases, and derive the framed BPS spectrum for the theory on the half line, with $L_{n'}$ boundary conditions. It would be interesting to match it to an explicit calculation. Also, it is conceivable that $s_\pm$ could themselves be identified with the vevs of certain boundary line defects $L_\pm$. The relation $s_\mp = Ms - \mathcal{Y}_{\pm\frac{1}{2}\gamma_f} s$ would seem to relate $L_\pm$ to the "difference" of space-filling brane branes with one or zero units of flux. This suggests the identification of the tentative $L_\pm$ with point-like branes (Dirichlet boundary conditions for the sigma model), possibly located at the North and South poles of $\mathbb{CP}^1$ because of the twisted mass in the Lagrangian.

8.3 The canonical surface defect in pure $SU(2)$ gauge theory

We are now ready to deal with the most complicated example in this paper, the canonical surface defect $\mathbb{S}_z$ in the pure $SU(2)$ gauge theory. The spectral curve in this case is

$$\lambda^2 = \left(\frac{\Lambda^2}{z^3} + \frac{2u}{z^2} + \frac{\Lambda^2}{z} \right) \mathrm{d}z^2. \tag{8.33}$$

Since the analysis to follow is long and technical let us summarize the basic points and lessons first here. We will first analyze the soliton spectrum at strong four-dimensional coupling. We will find that in the z-plane there are a finite number of domains in each of which there is a finite soliton spectrum. We will derive the spectrum of the BPS degeneracies μ and ω at fixed u as functions of z. We will check that the 2d-4d wall-crossing formulae are consistent with the monodromy of the local system Γ_{12} over the z plane. Then we will find a convenient regime in which to fix z and continue u from the strong coupling domain to the weak coupling domain. Since the 4d spectrum changes (dramatically) the walls of marginal stability in the z plane also change (dramatically). There is now an extremely complicated pattern of walls of marginal stability in the z plane (because there are infinitely many 4d bulk particles). We will describe some aspects of the resulting chambers in the z-plane qualitatively and check several nontrivial aspects of the wall-crossing phenomenon and soliton spectrum without giving a full description of μ and ω for u in this weak-coupling domain. In particular, at weak four-dimensional coupling there are two very different regimes. The weak two-dimensional coupling regime is defined by taking z to be in the "vectormultiplet annulus," that is, the annulus foliated by the finite WKB curves arising when ϑ is the phase of the vectormultiplet central charge. The strong two-dimensional coupling regime is then the complementary region in the z-plane. In the strong two-dimensional regime the soliton spectrum is finite in any chamber, but there are a countably infinite number of chambers and the spectrum is unbounded. In the weak two-dimensional regime the soliton spectrum is infinite and chamber-independent. Nevertheless, there is an uncountable number of chambers, and the BPS degeneracies ω depend on these chambers.



Figure 32: A basis for $H_1(\bar{\Sigma};\mathbb{Z})$ at $u=0$. We choose branch cuts emanating from $z=\mathrm{i}$, $z=0$ and $z=-\mathrm{i}$ and running along the imaginary axis. $Z_\gamma = K$ and $Z_{\gamma'} = -\mathrm{i}K$ for K a positive constant. The cycles γ and γ' have $\langle\gamma,\gamma'\rangle = 2$.

8.3.1 Preliminaries on local systems

Let us begin with some technical preliminaries. Some aspects of the theory have already been discussed in §3.4.2. In comparing with §3.4.2 one should take $u \to -u$ and $z = e^t$. Above, we described the local system Γ_{12} over the u-plane for fixed u. We also described the local system Γ over the u-plane $\mathcal{B}^*$. However, in this section it will be convenient to describe the local system $\Gamma \to \mathcal{B}^*$ in a slightly different way. We begin with $u=0$ (strong 4d coupling) and choose cuts for the covering (8.33) as in Figure 32. As shown in Figure 32 we have chosen cycles γ and γ' with $\langle\gamma,\gamma'\rangle = 2$. (We can map to §3.4.2 by identifying $\gamma_m = \gamma'$ and $\gamma_e - \sigma^*(\gamma_e) = (\gamma \pm \gamma')$. The choice of sign depends on how we continue $t_\pm$ to the imaginary axis; the fact that there is a choice reflects the fact that monodromy of the local system on the u-plane can take one form into the other.) The 4d spectrum for u in the strong coupling regime has $\Omega(\pm\gamma) = \Omega(\pm\gamma') = 1$, and all other degeneracies vanish. We will continue into the weak coupling regime; the weak coupling spectrum consists of a vector multiplet of charge $\pm(\gamma+\gamma')$ and $\Omega = -2$ and two infinite towers of hypermultiplets of charge $\gamma_n := \gamma + n(\gamma+\gamma')$ and $\gamma'_n := \gamma' + n(\gamma+\gamma')$, $n \in \mathbb{Z}$. Note that $-\gamma_n = \gamma'_{-(n+1)}$.



Figure 33: We choose $\gamma_{12}(z)$ as shown for z in the region U. The local system is trivialized on the complement of the cuts running down the imaginary axis, starting from $z = \mathrm{i}, 0, -\mathrm{i}$.

We will also need some facts about the local system Γ_{12} in the z-plane. We define a



Figure 34: The local system Γ_{12} in the z-plane is determined by the monodromy around the three paths $\mathcal{P}_{\mathrm{i}}$, $\mathcal{P}_0$, $\mathcal{P}_{-\mathrm{i}}$ shown.

class $\gamma_{12}(z)$ as in Figure 33. We can trivialize Γ_{12} on the complement of the cuts shown there. Using these pictures one can easily check that for $u = 0$ and Im z sufficiently large and positive we have $\omega(\gamma, \gamma_{12}(z)) = \omega(\gamma', \gamma_{12}(z)) = +1$.

The monodromy of Γ_{12} around the three paths shown in Figure 34 is

$$\begin{aligned} \mathcal{P}_{\mathrm{i}} &: \gamma_{12} \to \gamma_{21}, \\ \mathcal{P}_0 &: \gamma_{12} \to \gamma_{12} - \gamma - \gamma', \\ \mathcal{P}_{-\mathrm{i}} &: \gamma_{12} \to \gamma_{21} + \gamma' - \gamma. \end{aligned} \tag{8.34}$$

The most straightforward way to prove these equations is simply to transport $\gamma_{12}(z)$ around the respective paths and compute intersection products with γ, γ'. Alternatively, if we continue z along $\mathcal{P}_0$ to $z \approx 0$, we will find that $Z_{\gamma_{12}(z)+\frac{1}{2}(\gamma+\gamma')} \to 0$ as $z \to 0$ with a $\mathbb{Z}_2$ branch cut. Therefore, a cycle that only encircles $z = 0$ leads to a reflection

$$\left(\gamma_{12}(z) + \frac{1}{2}(\gamma + \gamma')\right) \to -\left(\gamma_{12}(z) + \frac{1}{2}(\gamma + \gamma')\right). \tag{8.35}$$

In a similar way we find that continuation along a path $\mathcal{P}_{-\mathrm{i}}$ gives $Z_{\gamma_{12}(z)+\gamma} \to 0$ for $z \to -\mathrm{i}$.

8.3.2 The soliton spectrum at strong coupling

Given the strong coupling 4d spectrum, the walls of marginal stability in the z-plane are obtained from the finite and separating WKB curves at the critical values $\vartheta = \arg \pm Z_{\gamma}$ and $\vartheta = \arg \pm Z_{\gamma'}$. For the case $u = 0$ these are shown in Figure 35. The complement of the union of these critical curves is the union of connected regions R, U, L, D, and their respective reflected images R', U', L', D'. The wall-crossing analysis will be very similar to that in §8.1.2, but now with *two* flavor charges γ and γ'.

It is easiest to begin the analysis of the soliton spectrum for z in regions U, D. As shown in Figure 37, in this case there are 3 distinct finite open WKB curves ending at z, implying the existence of 3 solitons (plus their antiparticles) in both these regions. We begin our analysis with z in the region U. In this case one of the open finite WKB curves is the projection of a path in the class $\gamma_{12}(z)$ defined in Figure 33. By carefully computing the intersection numbers of the three finite open curves one can establish that



Figure 35: The critical WKB curves for the monopole and dyon, of charges $\pm\gamma$ and $\pm\gamma'$ respectively. The turning points are at $z = \pm\mathrm{i}$. The separating WKB curves asymptote to horizontal lines.



Figure 36: Lines of marginal stability in the z-plane, for u in the strong coupling regime ($u = 0$ in this figure). The region R' is the image of R under $z \to 1/z$.

the charges of the three solitons in region U are $\gamma_{12}(z), \gamma_{12}(z) - \gamma', \gamma_{12}(z) + \gamma$ (together with their antiparticles). To find the proper ordering of the corresponding BPS rays, we must compute periods. One finds by direct computation that $Z_\gamma = K$ and $Z_{\gamma'} = -\mathrm{i}K$ where $K > 0$. [49] Now, for $Z_{\gamma_{12}(z)}$ it is useful to take $|z|$ large. Then when $z \in U$ one finds $Z_{\gamma_{12}(z)} \approx \frac{2}{\pi} z^{1/2}$ where we use the principal branch of the log. Thus ordering of central charges is as shown in Figure 38. As in §8.1.2, we will not give the analogous figures for the other seven strong coupling regions, but will simply write the corresponding products:

$$A(\sphericalangle; U) = \mathcal{K}_{\gamma'} \mathcal{K}_{\gamma} \mathcal{S}_{\gamma_{12}+\gamma} \mathcal{S}_{\gamma_{12}} \mathcal{S}_{\gamma_{12}-\gamma'}. \tag{8.36}$$

Here (unlike §8.1.2) we will choose $\sphericalangle$ to be an appropriate sector of width π (rather than 2π). We take it to be the right half-plane, in this and all subsequent expressions (for the

---

[49] In fact $K = 8\sqrt{2\pi}\frac{\Gamma(3/4)}{\Gamma(1/4)}$.


$\vartheta = \frac{\pi}{4}$
$\vartheta = \frac{\pi}{2}$
$\vartheta = 0$
$\vartheta = \frac{3\pi}{4}$

Figure 37: The separating WKB curves as a function of ϑ at $u = 0$. The critical WKB curves for 4d particles are shown at $\vartheta = 0$ and $\vartheta = \frac{\pi}{2}$. The unoriented WKB curves are the same for ϑ and $\vartheta + \pi$. Therefore, the figure only shows the curves as ϑ varies from 0 to π. The separating WKB lines going to infinity eventually become parallel. Therefore, from the above configurations of these separating WKB lines we can deduce that when z is in the region U or D there are precisely three finite open WKB curves connecting z to the turning point. When $|z|$ is large, the corresponding phases ϑ of the three finite open WKB curves are close together.

strong coupling region).

Now we consider the wall-crossings as z moves along a path $U \to L \to D$. As z crosses from U to L the phase of $Z_{\gamma_{12}(z)}$ becomes close to $\pi/2$ for $|z|$ large, so the two rays $\ell_{\gamma_{12}(z)}$ and $\ell_{\gamma_{12}(z)-\gamma'}$ in Figure 38 become parallel to $\ell_{-\gamma'}$ and we can apply the formula (8.2a). (Note that $\omega(-\gamma', \gamma_{12}(z)) = -1$.) This leads to a spectrum with only two solitons:

$$A(\triangleleft; L) = \mathcal{K}_{\gamma'} \mathcal{S}_{\gamma_{21}} \mathcal{K}_{\gamma} \mathcal{S}_{\gamma_{12}+\gamma}. \tag{8.37}$$

If we continue to increase the phase of z and cross from L into D then $\ell_{\gamma_{12}+\gamma}$ becomes parallel to $\ell_{-\gamma'}$ and (8.2b) leads to

$$A(\triangleleft; Dl) = \mathcal{K}_{\gamma'} \mathcal{S}_{\gamma_{21}-\gamma+\gamma'} \mathcal{S}_{\gamma_{21}-\gamma} \mathcal{S}_{\gamma_{21}} \mathcal{K}_{\gamma}. \tag{8.38}$$

Now let us compare moving z along a path $U \to R \to D$. Crossing from $U \to R$ at large $|z|$ the wall is at $\vartheta \approx 0$, and $Z_{\gamma_{12}(z)}$ and $Z_{\gamma_{12}(z)+\gamma}$ become parallel to Z_{γ}. We can then apply



Figure 38: Arrangement of the central charges for $z \in U$ in the strong coupling domain. Here we take $u = 0$ and $|z|$ large.

(8.2b) with $a = \gamma$ and $b = \gamma_{12}$ to get

$$A(\sphericalangle; R) = \mathcal{K}_{\gamma'}\mathcal{S}_{\gamma_{12}}\mathcal{K}_{\gamma}\mathcal{S}_{\gamma_{12}-\gamma'}. \tag{8.39}$$

Then, moving from $R \to D$, $Z_{\gamma_{12}-\gamma'}$ aligns with Z_γ and we get

$$A(\sphericalangle; Dr) = \mathcal{K}_{\gamma'}\mathcal{S}_{\gamma_{12}}\mathcal{S}_{\gamma_{12}-\gamma'}\mathcal{S}_{\gamma_{12}+\gamma-\gamma'}\mathcal{K}_{\gamma}. \tag{8.40}$$

Now (8.38) differs from (8.40), but we must take monodromy into account. The cuts in Figure 34 divide D into two regions Dl and Dr and, by (8.34), there is a discontinuity taking $\gamma_{21} \to \gamma_{12} + \gamma - \gamma'$ when passing from Dl to Dr. Thus, the wall-crossing formula is consistent with the monodromy of the local system around $\mathcal{P}_{-\mathrm{i}}$, as expected.

Now let us continue the spectrum from region $L \to L'$ in Figure 36. In this case z crosses the finite WKB curve for the charge γ, and this curve has phase $\vartheta = 0, \pi$. Consistency requires that the BPS rays $\ell_{\gamma_{12}+\gamma}$ and $\ell_{\gamma_{21}}$ move toward ℓ_γ and pass through each other when z is on the marginal stability wall. (This can be checked by noting that $Z_{\gamma_{12}(z)} \to 0$ as $z \to \mathrm{i}$.) Thus, the situation is very analogous to the $D \to U$ transition in the $N = 2$ AD example of §8.1.2. In particular we should apply the wall-crossing formula (8.3). Note that $\omega(\gamma, \gamma_{21}(z)) = -1$ for $z \in L$ (as needed to apply the formula) so that $\omega'(\gamma, \gamma_{21}(z)) = +1$ for $z \in L'$. The spectrum is now

$$A(\sphericalangle; L') = \mathcal{K}_{\gamma'}\mathcal{S}_{\gamma_{12}+\gamma}\mathcal{K}_{\gamma}\mathcal{S}_{\gamma_{21}}. \tag{8.41}$$

Similarly, when continuing from $z \in R$ to $z \in R'$, z crosses the finite WKB curve corresponding to γ', with phase $\vartheta = \pm\pi/2$, and we apply (8.3) with the rays $\ell_{\gamma_{12}}$ and $\ell_{\gamma_{21}+\gamma'}$ sweeping through $\ell_{\gamma'}$ to discover that $\omega'(\gamma', \gamma_{12}(z)) = -1$ for $z \in R'$, and the spectrum is

$$A(\sphericalangle; R') = \mathcal{K}_{\gamma'}\mathcal{S}_{\gamma_{21}+\gamma'}\mathcal{K}_{\gamma}\mathcal{S}_{\gamma_{21}}. \tag{8.42}$$

Now we would like to continue from $L' \to U'$ and from $R' \to U'$ and compare. To do this z must cross a separating WKB curve. The separating WKB curve between L' and U' has phase $\vartheta = \pm\pi/2$, so the central charge $Z_{\gamma_{12}}$ lines up with $Z_{\gamma'}$. In order to apply the wall-crossing formula (8.2b) with $b = \gamma_{12}(z)$ and $a = \gamma'$ we need to know that $\omega(\gamma', \gamma_{12}(z)) = +1$

for $z \in L'$. This is indeed the case, and is compatible with $\omega'(\gamma', \gamma_{12}(z)) = -1$ for $z \in R'$ thanks to the monodromy of the local system Γ_{12} around $z = \mathrm{i}$. In this way we get

$$A(\sphericalangle; U'l) = \mathcal{K}_{\gamma'} \mathcal{S}_{\gamma_{12}+\gamma'} \mathcal{S}_{\gamma_{12}} \mathcal{S}_{\gamma_{12}+\gamma} \mathcal{K}_{\gamma}. \tag{8.43}$$

In an analogous fashion we find that $\omega(\gamma, \gamma_{12}(z)) = +1$ for $z \in R'$ and hence

$$A(\sphericalangle; U'r) = \mathcal{K}_{\gamma'} \mathcal{S}_{\gamma_{21}+\gamma'} \mathcal{S}_{\gamma_{21}} \mathcal{S}_{\gamma_{21}+\gamma} \mathcal{K}_{\gamma}. \tag{8.44}$$

As in our comparison of Dl with Dr, the cuts in Figure 33 divide U' into two regions $U'l$ and $U'r$, and (8.43) agrees with (8.44) once we take into account the discontinuity across the cut summarizing the monodromy around $\mathcal{P}_{\mathrm{i}}$.

To complete the analysis we consider the wall-crossing from $L' \to D'$ and $R' \to D'$. Since $Z_{\gamma_{12}(z)+\gamma} \to 0$ as $z \to -\mathrm{i}$ from the region L' we can see that this central charge must line up with $Z_{\gamma'}$ and we apply (8.2a) with $a = \gamma'$ and $b = \gamma_{12}(z) + \gamma$ to obtain

$$A(\sphericalangle; D'l) = \mathcal{K}_{\gamma'} \mathcal{K}_{\gamma} \mathcal{S}_{\gamma_{21}} \mathcal{S}_{\gamma_{21}-\gamma} \mathcal{S}_{\gamma_{21}-\gamma-\gamma'}. \tag{8.45}$$

Finally, to do the wall-crossing from R' to D', since $Z_{\gamma_{12}(z)+\gamma'}$ aligns with Z_γ passing from R' to D' we apply (8.2b) to get

$$A(\sphericalangle; D'r) = \mathcal{K}_{\gamma'} \mathcal{K}_{\gamma} \mathcal{S}_{\gamma_{21}+\gamma+\gamma'} \mathcal{S}_{\gamma_{21}+\gamma'} \mathcal{S}_{\gamma_{21}}. \tag{8.46}$$

In an analogous way to the previous cases (8.45) is compatible with (8.46) once we take into account the monodromy around $\mathcal{P}_0$. This completes our analysis of the soliton spectrum and its compatibility with the wall-crossing formula.

8.3.3 The soliton spectrum for weak 4d coupling and strong 2d coupling



Figure 39: Critical WKB curves associated to γ and γ', when u is positive imaginary in the weak coupling domain.

Let us now consider what happens when z is fixed (in an appropriate region discussed below) and u moves along a path from strong coupling to weak coupling. To be specific, we consider a path so that the central charges Z_γ and $Z_{\gamma'}$ align. Then, the Kontsevich-Soibelman wall-crossing formula produces the weak coupling spectrum,

$$\left(\prod_{n \nearrow 0}^{\infty} \mathcal{K}_{\gamma_n} \right) \mathcal{K}_{\gamma+\gamma'} \left(\prod_{n \searrow \infty}^{0} \mathcal{K}_{\gamma'_n} \right), \tag{8.47}$$



Figure 40: Critical WKB curves associated to γ_n and γ'_n, with some large $n > 0$, when u is positive imaginary in the weak coupling domain. In each figure the finite WKB curve shown winds n times around the origin.



Figure 41: Critical WKB curves associated to the vectormultiplet of charge $\gamma+\gamma'$. The two purple finite WKB curves are the "outer" and "inner' vectormultiplet curves. The annulus between these two curves is foliated by closed WKB curves and is referred to as the "vectormultiplet annulus." This region corresponds to strong 2d coupling. The complement of the annulus in the z plane consists of an "outer region" connected to $z = \infty$ and an "inner region" connected to $z = 0$. These are both weak coupling domains for the 2d coupling.

in agreement with the standard result [25, 70, 71, 1, 2]. In this notation $\prod_{n\nearrow 0}^{\infty}$ means that the product is taken so that as one reads from left to right n increases $0, 1, \ldots, \infty$, while $\prod_{n\searrow\infty}^{0}$ similarly means that as one reads from left to right n decreases from ∞ to 0. We have suppressed the ω superscript but we remind the reader that $\Omega(\gamma+\gamma') = -2$. Because there are many new 4d particles, the walls of marginal stability in the z-plane change dramatically. If, for example, u moves along the positive imaginary axis into the weak coupling region then the critical WKB curves for γ and γ' evolve into those shown in Figure 39. In addition, there is an infinite tower of hypermultiplets with charges γ_n, γ'_n. These lead to critical WKB curves such as those shown in Figure 40. Finally a vectormultiplet emerges with charge $\gamma+\gamma'$ and critical WKB curves shown in Figure 41. The finite WKB curves for the vectormultiplet foliate an annulus. The finite WKB lines associated to the hypermultiplets all wind around the origin and reside inside this annulus. In addition each hypermultiplet contributes two separating WKB curves both outside and inside the annulus, with each pair of lines differing by one unit of winding around the origin. As $n \nearrow \infty$ these separating WKB curves accumulate from the outside on the outer boundary of the annulus and from the inside on the inner boundary of the annulus. The resulting



Figure 42: Taking the union of the critical hypermultiplet WKB curves with the inner and outer vectormultiplet curves produces an intricate pattern of walls of marginal stability. There are an infinite number of chambers accumulating from the outer and inner regions complementary to the annulus onto the outer and inner vectormultiplet curves, respectively. In the interior of the vectormultiplet annulus the walls are dense and there is an uncountable number of "chambers."

pattern of marginal stability lines, taking into account the first few hypermultiplets, is illustrated in Figure 42.

Now, to derive the soliton spectrum we first note that we can follow a path for $u \in \mathcal{B}^*$ so that the lines ℓ_γ and $\ell_{\gamma'}$ in Figure 38 sweep through each other without passing through the soliton lines, thus producing the configuration of central charges shown in Figure 43. This is clearly true if we take $|z|$ to be sufficiently large with large imaginary part. We will call this region U_0. Thus, for z in U_0, the spectrum is given by

$$A(\sphericalangle; U_0) = \left(\prod_{n\nearrow 0}^{\infty} \mathcal{K}_{\gamma_n}\right) \mathcal{K}_{\gamma+\gamma'} \left(\prod_{n\searrow\infty}^{0} \mathcal{K}_{\gamma'_n}\right) \mathcal{S}_{\gamma_{12}(z)+\gamma}\mathcal{S}_{\gamma_{12}(z)}\mathcal{S}_{\gamma_{12}(z)-\gamma'} \tag{8.48}$$

We have suppressed the superscript ω, but we must remember to extend our discussion slightly to find the ω. We consider the element $\gamma_{12}^0 = \gamma_{12} + \frac{1}{2}(\gamma - \gamma')$, for which $\omega(\gamma, \gamma_{12}^0) = \omega(\gamma', \gamma_{12}^0) = 0$ before the wall-crossing. It follows that also $\omega(\cdot, \gamma_{12}^0) = 0$ after the wall-crossing, just by acting with the BPS product on $X_{\gamma_{12}^0}$ (cf. a similar computation in §2.4). Then the affine-linearity of ω shows $\omega(m\gamma + n\gamma', \gamma_{12}) = (n+m)\Omega(m\gamma + n\gamma')$.

We now comment on the wall-crossing of the soliton spectrum, without doing a complete analysis. As we have noted, there are an infinite number of chambers outside the annulus bounded by separating WKB curves for hypermultiplets of charges γ_n and γ'_n. If z moves across such lines then there will be a wall-crossing formula of type (8.2a) or (8.2b). Let us choose one particular path where z begins on the imaginary axis at large positive imaginary part and moves downwards towards the vectormultiplet annulus. Referring to



Figure 43: The configuration of central charges after moving u along an appropriate path into the weak coupling domain, while holding z fixed with large imaginary part, corresponding to the upper region in Figure 42. The $\mathcal{K}$-rays have experienced wall-crossing while staying far from the $\mathcal{S}$-rays. The central charges corresponding to infinitely many more $\mathcal{K}$-rays, with phases accumulating to that of $Z_{\gamma+\gamma'}$, are suppressed.

Figure 42 it is clear that there will be an infinite sequence of wall-crossings as z moves across pairs of walls which intersect along the imaginary axis. We will call the resulting chambers $U_0, U_1, U_2, \ldots$ where U_0 is the noncompact chamber at large $|z|$. These chambers get smaller and accumulate at the intersection of the outer ring of the vectormultiplet annulus with the positive imaginary axis. Now, there is a canonical half-plane we can use to describe the spectrum whose boundaries are the W-boson lines. We will denote this sector as $\sphericalangle_W$. Thus, we can rewrite the spectrum (8.48) in the equivalent form:

$$A(\sphericalangle_W; U_0) = \mathcal{K}_{\gamma_W} \left(\prod_{n \searrow \infty}^{0} \mathcal{K}_{\gamma'_n} \right) \mathcal{S}_{\gamma_{12}(z)+\gamma} \mathcal{S}_{\gamma_{12}(z)} \mathcal{S}_{\gamma_{12}(z)-\gamma'} \left(\prod_{n \nearrow 0}^{\infty} \mathcal{K}_{-\gamma_n} \right) \tag{8.49}$$

Now, as z crosses from the noncompact region U_0 in Figure 42 across the topmost wall the line for $\gamma_{12}(z) - \gamma'$ sweeps counterclockwise across that for $-\gamma$ and simultaneously the line for $\gamma_{12}(z) + \gamma$ sweeps across that for γ'. We therefore have simultaneous wall-crossings of types (8.2b) and (8.2a) respectively, producing

$$\begin{aligned} A(\triangleleft_W;U_1) = \mathcal{K}_{\gamma_W} & \left(\prod_{n\searrow\infty}^{1} \mathcal{K}_{\gamma'_n} \right) \mathcal{S}_{\gamma_{12}+\gamma} \\ & \cdot (\mathcal{S}_{\gamma_{12}+\gamma_W}\mathcal{K}_{\gamma'})\mathcal{S}_{\gamma_{12}}(\mathcal{K}_{-\gamma}\mathcal{S}_{\gamma_{12}-\gamma_W})\cdot \\ & \mathcal{S}_{\gamma_{12}-\gamma'} \left(\prod_{n\nearrow 1}^{\infty} \mathcal{K}_{-\gamma_n} \right) . \end{aligned} \tag{8.50}$$

We are now set up for an inductive process, since once again on the left side of the product the ray with charge $\gamma_{12}+\gamma$ can sweep clockwise across the next $\mathcal{K}$-ray of charge γ'_1, while on the right the ray for $\gamma_{12}-\gamma'$ can sweep counter-clockwise across $-\gamma_1$. After N such steps we find

$$\begin{aligned} A(\triangleleft_W;U_N) = \mathcal{K}_{\gamma_W} & \left(\prod_{n\searrow\infty}^{N} \mathcal{K}_{\gamma'_n} \right) \mathcal{S}_{\gamma_{12}+\gamma} \\ & \left(\prod_{n\searrow N-1}^{0} (\mathcal{S}_{\gamma_{12}+(n+1)\gamma_W}\mathcal{K}_{\gamma'_n}) \right) \mathcal{S}_{\gamma_{12}} \left(\prod_{n\nearrow 0}^{N-1} (\mathcal{K}_{-\gamma_n}\mathcal{S}_{\gamma_{12}-(n+1)\gamma_W}) \right) \cdot \\ & \mathcal{S}_{\gamma_{12}-\gamma'} \left(\prod_{n\nearrow N}^{\infty} (\mathcal{K}_{-\gamma_n}) \right) . \end{aligned} \tag{8.51}$$

Remarks

1. One might wonder whether the path we have described meets other wall-crossings, involving exchanges of other $\mathcal{K}$ and $\mathcal{S}$ factors in (8.51). The easiest way to see that this cannot happen is to note that the $\omega(a,b)$ associated to these other potential crossings would not be compatible with (8.2a) or (8.2b).

2. Other paths from U_0 into the vectormultiplet annulus will produce interesting variants of (8.51) which are not symmetrical between the products in the middle line of this equation. We have not attempted to investigate the full chamber structure and soliton spectrum in detail.

3. Using the $z \to 1/z$ symmetry we can also conclude that there is a similar spectrum of solitons in the regions inside the vectormultiplet annulus.

4. There is a nice physical interpretation of the spectrum we have found. As we have discussed in §8.2, the strongly coupled $\mathbb{CP}^1$ sigma model has only two solitons whereas (8.48) predicts 3 solitons. In order to compare these spectra we should take the limit as the 4d $SU(2)$ gauge coupling becomes infinitely weak. One way to do this is to set $\tilde{z} = \Lambda^2 z$ and take $\Lambda \to 0$ holding $\tilde{z}$ fixed. In this limit the chiral ring equation (8.33) degenerates to (8.25). Moreover, the period $Z_{\gamma_{12}(z)}$ diverges as $\sim -\sqrt{2u}\log\frac{\Lambda^4}{2u}$

while $Z_{\gamma_{12}-\gamma'}$ and $Z_{\gamma_{12}+\gamma}$ have finite limits. [50] It follows that the soliton with charge γ_{12} becomes large. Indeed, identifying $\sqrt{2|u|}=v$, where v is the scale of the vev of the 4d vacuum, this soliton has a mass $\sim \frac{32\pi v}{g^2}$, and in the weak coupling limit $g \to 0$ it decouples from the spectrum. By contrast, the solitons with charges $\gamma_{12}+\gamma$ and $\gamma_{12}-\gamma'$ remain in the spectrum. Similarly, in (8.51) we find two towers of length N of solitons with charges $\gamma_{21}+n(\gamma+\gamma')$ and $\gamma_{12}+n(\gamma+\gamma')$, $n=1,\ldots,N$. These too have divergent central charges going like $\sim -\sqrt{2u}\log\frac{\Lambda^4}{2u}$ and hence should be viewed as bound states of the hypermultiplet dyons with the surface defect. All of these heavy particles should correspond to solitonic field configurations visible in a semiclassical analysis of the $\mathbb{CP}^1$ sigma model coupled to the weakly coupled $SU(2)$ theory. It would be an interesting check to produce these solitons directly. It would also be interesting to test directly the predictions $\omega(\gamma_n,\gamma_{12})=\omega(\gamma'_n,\gamma_{12})=2n+1$. In any case, we conclude that the spectrum (8.48) is physically sensible.

5. As z moves towards the vectormultiplet annulus the product approaches the expression

$$\begin{aligned} A(\sphericalangle_W;U_\infty) &= K_{\gamma_W} S_{\gamma_{12}+\gamma}\left(\prod_{n\searrow\infty}^{-\infty}\left(\mathcal{S}_{\gamma_{12}+(n+1)\gamma_W}\mathcal{K}_{\gamma'_n}\right)\right)\cdot\mathcal{S}_{\gamma_{12}-\gamma'} \\ &= K_{\gamma_W} S_{\gamma_{12}+\gamma}\left(\prod_{n\nearrow-\infty}^{\infty}\left(\mathcal{K}_{-\gamma_n}\mathcal{S}_{\gamma_{12}-(n+1)\gamma_W}\right)\right)\cdot\mathcal{S}_{\gamma_{12}-\gamma'} \end{aligned} \tag{8.52}$$

8.3.4 The soliton spectrum for weak 4d coupling and weak 2d coupling

When z moves into the vectormultiplet annulus both the 4d and 2d theories are weakly coupled. There is an infinite spectrum of 4d hypermultiplets leading to a very intricate pattern of marginal stability lines inside the annulus, as we have already mentioned, and in addition, experience with the $\mathbb{CP}^1$ sigma model in §8.2 suggests that there should also be an infinite spectrum of weakly coupled 2d $\mathbb{CP}^1$ solitons, in addition to those we have already discovered. Indeed, let us return to the limiting expression (8.52). By slightly shifting the half-plane we can write an equivalent product, cycling the factor $\mathcal{S}_{\gamma_{12}-\gamma'}$ on the right to its anti-particle factor $\mathcal{S}_{\gamma_{21}+\gamma'}$ on the left. Then, the product $\mathcal{S}_{\gamma_{21}(z)+\gamma'}\mathcal{K}_{\gamma+\gamma'}\mathcal{S}_{\gamma_{12}(z)+\gamma}$ has the property that $\omega(\gamma+\gamma',\gamma_{21}(z)+\gamma')=0$ and $\omega(\gamma+\gamma',\gamma_{12}(z)+\gamma)=0$, as can be seen by computing intersection products. Therefore, we can apply the $\mathbb{CP}^1$ wall-crossing formula (2.51) to this product to produce a product on the left of the form

$$\left(\prod_{n\nearrow 0}^{\infty}\mathcal{S}_{\gamma_{12}+\gamma_n}\right)\mathcal{K}^{\omega}_{\gamma_W}\left(\prod_{n\searrow\infty}^{0}\mathcal{S}_{\gamma_{21}+\gamma'_n}\right) \tag{8.53}$$

---

[50] To prove these statements note that in this limit one turning point approaches $\tilde{z}=0$ like $\tilde{z}\approx-\Lambda^4/(2u)$ while the other approaches $\tilde{z}\approx-2u$. The WKB curve for γ_{12} goes to the turning point near the origin, and the term $\sqrt{2u}\frac{d\tilde{z}}{\tilde{z}}$ dominates the line integral of λ. On the other hand for the charges $\gamma_{12}+\gamma$ and $\gamma_{12}-\gamma'$ the WKB curve goes to the other turning point $-2u$ and the line integral does not diverge.

where $\omega(\gamma+\gamma', \gamma_{12}+\gamma) = 2$. Using this one can derive a heuristic formula for the spectrum when z lies *exactly on* the outer boundary of the annulus for the vectormultiplet. Written back in the standard half-plane $\sphericalangle_W$ this can be written in several forms, of which two of the more suggestive ones are:

$$\begin{aligned} A(\sphericalangle_W; Ann) &= \mathcal{K}_{\gamma_W} \left(\prod_{n \searrow \infty}^{0} \mathcal{S}_{\gamma_{21}+\gamma'_n} \right) \cdot \left(\prod_{n \searrow \infty}^{-\infty} (\mathcal{S}_{\gamma_{12}+(n+1)\gamma_W} \mathcal{K}_{\gamma'_n}) \right) \cdot \left(\prod_{n \searrow -1}^{-\infty} \mathcal{S}_{\gamma_{21}+\gamma'_n} \right) \\ &= \mathcal{K}_{\gamma_W} \left(\prod_{n \nearrow -\infty}^{-1} \mathcal{S}_{\gamma_{21}-\gamma_n} \right) \cdot \left(\prod_{n \nearrow -\infty}^{\infty} (\mathcal{K}_{-\gamma_n} \mathcal{S}_{\gamma_{12}-(n+1)\gamma_W}) \right) \cdot \left(\prod_{n \nearrow 0}^{\infty} \mathcal{S}_{\gamma_{21}-\gamma_n} \right) \end{aligned} \tag{8.54}$$

Of course these are simply related by using the identity $\gamma_{-(n+1)} = -\gamma'_n$.

The expression (8.54) is slightly unphysical. When z moves into the interior of the vectormultiplet annulus it crosses an infinite number of walls, and hence there will be an infinite number of wall-crossings applied to (8.54). At any interior point there will be some definite ordering of the central charges appearing in (8.54) determining the true expression. Indeed, we claim that the possible soliton spectra and chambers in the vectormultiplet annulus are precisely those given by the bi-infinite words in the $\mathcal{S}$ and $\mathcal{K}$ factors appearing in (8.54) subject to the following rules:

Define

$$\begin{aligned} \mathcal{A}_n &= \mathcal{S}_{\gamma_{21}+\gamma'_n}, \\ \mathcal{C}_n &= \mathcal{S}_{\gamma_{12}+(n+1)\gamma_W}, \\ \mathcal{K}_n &= \mathcal{K}_{\gamma'_n}, \end{aligned} \tag{8.55}$$

where $n \in \mathbb{Z}$. Then the bi-infinite words are of the form

$$\cdots \mathcal{S}\mathcal{K}\mathcal{S}\mathcal{K}\mathcal{S} \cdots \tag{8.56}$$

such that:

1. The $\mathcal{S}$-factor between $\mathcal{K}_n$ and $\mathcal{K}_{n-1}$ can be either $\mathcal{A}_m$ or $\mathcal{C}_m$ for some m.
2. The word therefore determines three sequences of letters of type $\mathcal{K}_n$, $\mathcal{A}_n$ and $\mathcal{C}_n$, respectively. The index on all three sequences decreases to the right in steps of one. Thus we have three sequences $\cdots \mathcal{K}_{n+1}\mathcal{K}_n \cdots$, $\cdots \mathcal{A}_{n+1}\mathcal{A}_n \cdots$ and $\cdots \mathcal{C}_{n+1}\mathcal{C}_n \cdots$ interwoven in the pattern (8.56).

We will give a strong argument for this claim, although we do not insist it is a completely rigorous proof.

Before establishing our claim, let us note that it is nicely consistent with wall-crossing. First, when z crosses the closed WKB curves there is no wall-crossing, which is good since they foliate the annulus. Indeed every point z lies on one such closed curve and, as noted in §8.2, these give the two weakly coupled vacua at the north and south poles of the $\mathbb{CP}^1$

sigma model. Next, note that all the other walls in the interior of the annulus are finite hypermultiplet walls for γ'_n for some $n \in \mathbb{Z}$. Note that the sum of charges for $\mathcal{A}_n$ and $\mathcal{C}_m$ is precisely γ'_{n+m+1}. Therefore, when z crosses the wall for γ'_{n+m+1} there can be a wall-crossing event of type (8.3), involving

$$\mathcal{A}_n \mathcal{K}_{n+m+1} \mathcal{C}_m = \mathcal{C}_m \mathcal{K}_{n+m+1} \mathcal{A}_n. \tag{8.57}$$

It might not at first appear obvious that $\omega(\gamma'_{n+m+1}, \gamma_{21}(z) + \gamma'_n) = -1$ or $\omega(\gamma'_{n+m+1}, \gamma_{12}(z) + (m+1)\gamma_W) = -1$ as is required for (8.3) (which case is realized depends on which side of the wall z is on). This can be shown by noting that as z approaches the WKB curve for γ'_{n+m+1} the phase of the line for $\gamma_{21}(z) + \gamma'_n$ approaches that for γ'_{n+m+1}. On the other hand, WKB walls with the same value of ϑ cannot intersect except at turning points and singularities. Therefore, the topology of the WKB curves must be that shown in the universal situation illustrated in Figure 14. Equation (8.57) now shows how wall-crossings (possibly infinitely many) allow us to pass between any two words of the type (8.56) described above.



Figure 44: In case (a) the corresponding sequence of morphisms is $\cdots \mathcal{K}_n \mathcal{A}_m \mathcal{K}_{n-1} \cdots$ because the slope of the $\mathcal{A}$ ray lies between those of the $\mathcal{K}$ rays. In case (b) the corresponding sequence is $\cdots \mathcal{K}_{n+1} \mathcal{C}_m \mathcal{K}_n \cdots$.

Now, to investigate the chamber structure and establish our claim it is first useful to map the annulus to the w-plane,

$$w := \int_{z_+}^{z} \lambda, \tag{8.58}$$

where we integrate from the turning point z_+ nearest to $z = 0$. It will be useful to work with an approximate formula for w, valid deep in the weak coupling domain. Let us introduce the scaled coordinate

$$\tilde{z} = \left(\frac{\Lambda^2}{u}\right)^{2\alpha} z \tag{8.59}$$

with $0 < 2\alpha < 1$. We consider the weak coupling limit where $u \to \infty$ (or equivalently $\Lambda \to 0$) with $\tilde{z}$ held fixed. In this regime the middle term in the expression (8.33) for

λ^2 dominates the other two and hence we can approximate $\lambda \approx \sqrt{2u}\frac{\mathrm{d}z}{z}$ and hence $w \approx \sqrt{2u}\log z/z_+$. After further rescaling the real and imaginary parts of w we can therefore take the image of the annulus to be the infinite strip between 0 and 1, with inner turning points at $\mathrm{i}N$ and outer turning points at $1+\mathrm{i}N$, $N \in \mathbb{Z}$. In w-coordinates the WKB curves are straight lines of slope proportional to ϑ. The lines of marginal stability are the straight lines joining these turning points, and are hence labeled by pairs of integers, (m,n) and given by $y = (m-n)x + n$. After solving a simple Diophantine equation one can show that every point in the strip with rational coordinates is at the intersection of two such lines. In particular, the walls of marginal stability are dense and the "vertices" of the chambers are the points in the strip with rational coordinates. Now, any two points not on a WKB wall will be separated by such a wall. (To prove this draw a straight line segment between the two points. Choose a point with rational coordinates sufficiently close to that line segment. There will be a WKB wall through that point.) Therefore, the "chambers" — defined to be the connected components of the complement of the walls — will consist of single points. The set of walls is a countable union of sets of measure zero and hence has measure zero, and therefore the chambers have positive measure, and therefore there are uncountably many chambers.

Now suppose that a surface defect is at a point z_0 in the vectormultiplet annulus and z_0 maps to a point w_0. The WKB curves for charges associated with $\mathcal{A}_n$ are the straight lines going from w_0 to the turning points $\mathrm{i}N$ on the inner boundary while the WKB curves for charges associated with $\mathcal{C}_n$ are the straight lines going from w_0 to the turning points $1+\mathrm{i}N$ on the outer boundary of the annulus. (Note that this way two such lines can glue together to give a hypermultiplet WKB curve.) It is now a matter of simple geometry to see that w_0 can lie in one of two different kinds of triangles. In each triangle there is exactly one $\mathcal{S}$ factor with slope between those of $\mathcal{K}_n$ and $\mathcal{K}_{n-1}$. In one type of triangle the slope between $\mathcal{K}_n$ and $\mathcal{K}_{n-1}$ will be of $\mathcal{A}$-type and in the other it will be of $\mathcal{C}$-type. See Figure 44.

Remarks

1. If in Figure 42 z continues down the imaginary axis then there must be an infinite number of wall-crossings, ending up at a product analogous to (8.54) for the inner ring of the annulus, with $\mathcal{A}$ and $\mathcal{C}$ exchanged. This is analogous to a phenomenon we found at strong coupling, where $z \to 1/z$ had the effect of transforming the soliton spectrum by the involution $\gamma_{12} \to \gamma_{21} + \gamma$.

2. Our analysis makes some interesting predictions which would be worth checking with weakly coupled semiclassical field theory methods. The region defined by the scaling parameter (8.59) defines a regime where the surface defect may be treated as a Gukov-Witten defect. This may be justified by using the approximate expression $\lambda \approx \sqrt{2u}\frac{\mathrm{d}z}{z}$ to obtain approximate expressions for $Z_{\gamma_{12}(z)}$, when z is in the domain with $u \to \infty$

and $\tilde{z}$ held fixed. One then finds the IR GW parameter

$$t_{\gamma_1} - t_{\gamma_2} = \frac{\partial Z_{\gamma_{12}(z)}}{\partial a} = \begin{cases} \frac{(2\alpha-1)}{2\pi \mathrm{i}} \log \frac{a}{\Lambda} + \cdots & \text{outer turning point,} \\ \frac{(2\alpha+1)}{2\pi \mathrm{i}} \log \frac{a}{\Lambda} + \cdots & \text{inner turning point.} \end{cases} \tag{8.60}$$

In a semiclassical analysis one would define a moduli space of “ramified monopoles,” solutions of the bulk BPS equations in the presence of GW boundary conditions dictated by (8.60). Semiclassical quantization of this moduli space for various monopole charges should reproduce the weak-coupling spectrum described in this section: in particular, the spectrum of solitons $\mu(\gamma_{ij})$ is independent of z but the spectrum of $\omega(\gamma, \gamma_{ij})$ varies strongly with z, and should grow at most linearly in the charges.

3. It should be interesting to study the framed BPS states in this example, but we leave this for another occasion.

9. An application: solving Hitchin systems by integral equations

Finally let us consider an interesting application of our discussion: we can use it to give concrete formulas for solutions of Hitchin equations on punctured surfaces C.

Indeed, as we explained in §7.3, for theories of class $\mathcal{S}$, the $\mathcal{Y}_{\gamma_i}(\zeta)$ can be thought of as sections of the universal Higgs bundle over $\mathcal{M} \times C$. Allowing z to vary we obtain $\mathcal{Y}_{\gamma_i}(\zeta, z)$, a flat section with respect to the connection $\mathrm{d} + \mathcal{A}(\zeta)$. This follows from comparing (5.43) and (7.16). Alternatively, we can use $\mathcal{Y}$ to define the connection (see below) and then apply an argument analogous to that used in Appendix E to determine that the connection so defined is the desired one. This flat section is concretely computable using the integral equations (5.23), (5.25). The only data one needs to write these equations are the central charges Z_γ and the BPS degeneracies ω, μ.

Suppose we trivialize the ramified cover $\Sigma \to C$ in a neighborhood of z. Then we can label the sheets $i = 1, 2$, and similarly, we can locally trivialize the cover above z' and label the sheets $i' = 1, 2$. We can compute the four sections $\mathcal{Y}_{\gamma_{ii'}}$ in a neighborhood of the points z and z', using the integral equations (5.23) and (5.25) together with (5.32). Any invertible combination of these four sections defines a fundamental solution Ψ of the flatness equations at z. Then $\Psi^{-1} \mathrm{d}_z \Psi = \varphi/\zeta + A + \zeta\bar{\varphi}$ gives the solution to the Hitchin system, in the form of explicit $\varphi, A, \bar{\varphi}$ valid in a neighborhood of z (with the particular solution and gauge depending on a basepoint z'). Different choices of Ψ are possible, in the form of different linear combinations of the various $\mathcal{Y}_{ii'}$. They will give different gauge choices for the solution. For examples, if z and z' are in a common neighborhood with a common trivialization of the double cover, then at least for sufficiently large R, a sum like

$$\Psi = \mathcal{Y}_{\gamma_{12'}} + \mathcal{Y}_{\gamma_{21'}} \tag{9.1}$$

or alternatively a sum like

$$\Psi = \mathcal{Y}_{\gamma_{11'}} + \mathcal{Y}_{\gamma_{22'}} \tag{9.2}$$

will serve as fundamental solutions.

Again, taking z, z' in some common neighborhood we can also apply the discussion around (5.33) of §5.6 where we take the diagonal matrix to be

$$x(z,\zeta;z') = x_{\gamma_{i2'}}(z,\zeta;z')e_{ii} \tag{9.3}$$

Here z' and its lift to sheet $2'$ are serving as basepoints. e_{ii} are matrix units and the sum on $i = 1,2$ is understood. We can then consider $g(\zeta)x(z,\zeta;z')$ to be a fundamental solution to $(\mathrm{d}_z + \mathcal{A})\Psi = 0$. It follows that gauge transformation by $g(\zeta)$ diagonalizes the flat spectral connection for the Hitchin system, to a diagonal connection

$$\mathrm{d}_z \log x_{\gamma_{i2'}} e_{ii}. \tag{9.4}$$

We can make this more explicit using (5.23): the connection we obtain is the same as the semiflat connection

$$\mathrm{d}_z \log \mathcal{Y}^{\mathrm{sf}}_{\gamma_{i2'}} e_{ii} \tag{9.5}$$

plus the correction

$$\mathrm{d}_z \left[\sum_{\gamma} \omega(\gamma,\gamma_i;z) \frac{1}{4\pi \mathrm{i}} \int_{\ell_\gamma} \frac{\mathrm{d}\zeta'}{\zeta'} \frac{\zeta'+\zeta}{\zeta'-\zeta} \log(1-\mathcal{Y}_\gamma(\zeta')) \right] e_{ii}. \tag{9.6}$$

The only z-dependence in the expression in square brackets comes from the discontinuities in $\omega(\gamma,\gamma_i;z)$ due to wall-crossing when z crosses finite WKB curves (see Figure 14 in §7.2). These give delta-function singularities localized on these curves. We have argued on general grounds that our solution to the Hitchin equations should be smooth, so these singularities are expected to cancel against similar discontinuities in $g(\zeta)$. However, we have not checked this point in detail.

Because of the simple ζ dependence of $\mathcal{A}$, the information in the full $\mathcal{Y}_{ab'}(\zeta)$ is rather redundant. It is sufficient to expand asymptotically for small ζ, so that

$$x_{\gamma_i}(\zeta) \sim \mathcal{Y}^{\mathrm{sf}}_{\gamma_i}(\zeta) \exp\left[-\sum_{\gamma} \omega(\gamma,\gamma_i) \frac{1}{4\pi \mathrm{i}} \int_{\ell_\gamma} \frac{\mathrm{d}\zeta'}{\zeta'} (1 + 2\frac{\zeta}{\zeta'} + \cdots) \log(1-\mathcal{Y}_\gamma(\zeta')) \right]. \tag{9.7}$$

and

$$g_i(\zeta) \sim g_i^{\mathrm{sf}} + \sum_{j\neq i,\gamma_{ji}} \mu_{\gamma_{ji}} \frac{1}{4\pi \mathrm{i}} \int_{\ell_{\gamma_{ji}}} \frac{\mathrm{d}\zeta'}{\zeta'} (1+\cdots) g_j(\zeta') x_{\gamma_{ji}}(\zeta'), \tag{9.8}$$

The leading order for g_i and the leading order for x_{γ_i} is sufficient to derive the expression for φ. The next to leading order in x_{γ_i} is required to derive A. Then $\bar{\varphi}$ can be obtained simply by complex conjugation, or from a large ζ expansion.

10. Future directions

In conclusion, we list here some directions for future research which be believe might be fruitful.

1. The exercise of §7 appears to require genuinely new techniques for higher rank groups. We are currently exploring this question.

2. Our construction produces examples of hyperholomorphic bundles over moduli spaces $\mathcal{M}$ of Higgs bundles. Such hyperholomorphic bundles fit naturally into the approach of [72] to the geometric Langlands correspondence: they are examples of "$(B,\ B,\ B)$ branes" in that context. The sections $\tilde{\nu}_i$ which entered our construction as part of the 2d-4d data define a complex Lagrangian multi-section of the dual moduli space $\widetilde{\mathcal{M}}$, i.e. a "$(B,\ A,\ A)$ brane." These two branes are related by mirror symmetry, i.e. Langlands duality. In this paper we have obtained a more explicit picture of the geometry of these $(B,\ B,\ B)$ branes than previously known. It is natural to wonder whether this picture can be of any use in the geometric Langlands correspondence.

3. In §6.1.7 we encountered, but did not really address, the question of systematically describing the structure of our hyperholomorphic bundles near orbifold singularities of $\mathcal{M}$. Via mirror symmetry this seems to be related to the same question for branes supported on complex Lagrangian sections of $\widetilde{\mathcal{M}}$ which run into singularities. For branes supported on a single singular fiber of the Hitchin fibration, the relevant extra structure has been discussed in [57]; it would be desirable to have a discussion analogous to that of [57] which applies to the branes we met here.

4. In §4 we sketched a 2d-4d spin wall-crossing formula. However, some work remains to be done in connecting this formula to the physical halo picture in the case when $y \neq -1$. For example, we have left out the details of how to relate the wall-crossing data $d(i,s)$ etc. to the physical halo particle degeneracies.

5. It would be interesting to generalize one of the main results of [3] from line defects to surface defects. In [3] it was shown that the formal generating functions for line defects $F(L) = \sum_\gamma \overline{\underline{\Omega}}(L_\zeta, y, \gamma) X_\gamma$ satisfy an algebra which is a quantum deformation of the algebra of holomorphic functions on $\mathcal{M}^\zeta$. This algebra can be interpreted as the algebra $\mathcal{O}_{BB}$ of open BB strings of the canonical coisotropic brane on $\mathcal{M}$, a certain type of A-brane. Now, in the present paper we have defined $F(L) = \sum_{\gamma_{ij}} \overline{\underline{\Omega}}(L, \gamma_{ij}) X_{\gamma_{ij}}$. The algebra of these functions can be identified with the algebra of holomorphic sections of $\mathrm{Hom}(V_{\mathbb{S}}, V_{\mathbb{S}})$, which is the algebra of open string states on a B-brane whose Chan-Paton bundle is $V_{\mathbb{S}}$. As in §4 we can introduce a y-deformation $\widehat{F}(L) = \sum_{\gamma_{ij}} \overline{\underline{\Omega}}(L, \gamma_{ij}, y) \widehat{X}_{\gamma_{ij}}$ to produce a y-dependent deformation of the noncommutative — but "classical" — algebra $\Gamma\left(\mathrm{Hom}(V_{\mathbb{S}}, V_{\mathbb{S}})\right)$. It is natural to conjecture that this y-deformed algebra is related to the open string algebra $\mathcal{O}_{B_{\mathbb{S}},B_{\mathbb{S}}}$ where $B_{\mathbb{S}}$ is a higher rank generalization of the canonical coisotropic brane, (an A-brane), which should, moreover have a natural hyperholomorphic connection. (Higher rank coisotropic branes have been investigated in [73].)

6. Recently, Witten has introduced an approach to knot homology based on the Hilbert space associated to surface defects in six-dimensional $(2,0)$ theory [74]. It is natural to ask whether the considerations of this paper can play a role in this approach to knot homology. Let us consider a theory in the class $\mathcal{S}$ on $\mathbb{R}^4 \times C$. Let us label coordinates $x^{0,1,2,3} \in \mathbb{R}^4$ and $z \in C$. Consider a knot $K \subset \mathbb{R} \times C$ described

by an equation $z = z(x^3)$. With a suitable topological twisting we can consider the surface defect $\mathbb{S}_z$ at $x^1 = x^2 = 0$ with worldsheet coordinates x^0 and x^3, with parameter $z = z(x^3)$ varying along x^3. Imagine that $z(x^3)$ is generically constant, but has rapid transitions — like a soliton. Then we can consider the surface defect to have several interfaces. More generally, one could consider a surface defect with various kinds of interfaces and Janus defects. It would appear that one would need to generalize surface defect amplitudes to amplitudes corresponding to several surface defects located at $(z_1, \dots, z_n)$. Naively, these correspond to having several M2-branes end on the M5-brane. It would be interesting to generalize the notion of framed BPS states to such situations. If this can be done it would then be natural to ask if the spaces of framed BPS states associated to this setup are related to knot homologies. As a first step one would want to relate the q-grading of knot homology to the y-grading of spaces of framed BPS states.

7. It might be interesting to apply some of our techniques to the study of branes in Landau-Ginzburg models, a subject which has been extensively discussed in [75]. Indeed it was noted at the very end of [17] that the period amplitudes Π_i^a of [75] should be related to supersymmetric interface amplitudes between a surface defect and the null surface defect. This in turn might shed further shed light on the quantum McKay correspondence [76, 77, 78].

8. The integral equations of [1] for the $\mathcal{Y}_\gamma$ are formally similar to the thermodynamic Bethe ansatz. The new integral equations introduced in §5.6 are very similar to those used in the inverse scattering method of integrable systems theory [79, 80]. The $x_{\gamma_{ij}}$ are analogous to scattering data, and the g_k are used to solve the associated linear problem. In some special cases the link can be made much more explicit. For example, in a recent paper S. Lukyanov and A. Zamolodchikov [81] studied the modified sinh-Gordon equation. This intersects with the Hitchin systems for AD theories for special values of the parameter α of [81]. Equation (5.15) of [81] is related to our proposal for solving the auxiliary linear problem. It would be useful to sharpen this relation and see whether there is room for any interesting technology transfer between the theory of integrable systems and the theory of defects in supersymmetric Yang-Mills theory.

9. Recently the methods of [1, 2] have found some surprising applications to scattering in $\mathcal{N} = 4$ super Yang-Mills theory [82, 83, 84, 85]. We believe that applying the techniques of this paper to that situation should allow one to study the shape of the minimal surfaces in AdS_5 whose area computes the amplitudes. Information about the shape of the surface can be useful in computing mixed correlation functions of a polygonal Wilson loop and local operators in the strong coupling limit: they should correspond to the integral over the surface of an appropriate bulk-to-boundary propagator.

10. Another physical context in which this structure appears is in $\mathcal{N} = 2$ supergravity. Surface defects are replaced by cosmic strings and dyons are replaced by dyonic

black holes [86, 87]. Perhaps the 2d-4d wall-crossing formula can be useful in this context. Related to this, we can make surface defects by wrapping D-branes on Calabi-Yau manifolds. For example, in Type IIA string theory we can consider D4-branes wrapping supersymmetric 3-cycles in a Calabi-Yau manifold. It should be very interesting to apply the present formalism to these surface defects and we intend to return to this subject in the future.

11. Finally, the 2d-4d BPS degeneracies we have described in this paper should presumably be identified mathematically as the "open" analogue of Donaldson-Thomas invariants. It would be very interesting to develop this theory. (Some investigation of such open invariants has been carried out in [88, 89].)

Acknowledgements

We thank David Ben-Zvi, Emanuel Diaconescu, Dan Freed, Martin Rocek, Nathan Seiberg, Karen Uhlenbeck, and Edward Witten for discussions.

We would like to thank the Simons Center for Geometry and Physics for hosting several excellent workshops related to this project and for hospitality at those workshops. We also thank the Aspen Center for Physics for hospitality during the initial writing of this paper. GM would like to thank the SCGP for hospitality during the final stages of writing this paper.

The work of GM is supported by the DOE under grant DE-FG02-96ER40959. The work of AN is supported by the NSF under grant number DMS-1006046. DG is supported in part by the NSF grant PHY-0503584. DG is supported in part by the Roger Dashen membership in the Institute for Advanced Study.

A. Relation of 2d and 4d superalgebras and their multiplets

The preserved supersymmetries of our surface defect are part of a subalgebra of the $\mathcal{N}=2$, $d=4$ superalgebra which can be seen to be isomorphic to a $d=2$, $(2,2)$ superalgebra by mapping

$$Q_+ = {Q_2}^2 = -Q_{21} \qquad\qquad \bar{Q}_+ = \bar{Q}_{\dot{2}2} \tag{A.1}$$

$$Q_- = \bar{Q}_{\dot{1}1} \qquad \bar{Q}_- = {Q_1}^1 = Q_{12} \tag{A.2}$$

One can then check using the $d=4$ $\mathcal{N}=2$ algebra relations that

$$\begin{aligned}
&\{Q_+,\bar{Q}_+\} = -2(P_0+P_3)\\
&Q_+^2 = \bar{Q}_+^2 = 0\\
&\{Q_-,\bar{Q}_-\} = -2(P_0-P_3)\\
&Q_-^2 = \bar{Q}_-^2 = 0\\
&\{Q_+,Q_-\} = \{\bar{Q}_+,\bar{Q}_-\} = 0\\
&\{Q_+,\bar{Q}_-\} = -2\bar{Z}\\
&\{\bar{Q}_+,Q_-\} = -2Z
\end{aligned} \tag{A.3}$$

The $u(1)_V$ symmetry of $(2,2)$ is identified with the action of the generator of $so(2)_{12}$ (which equals that of $u(1)_r$ acting on the supercharges) and the $u(1)_A$ symmetry is identified with $u(1)_R$.

An important point is that one does *not* need to set $P_1 = P_2 = 0$ when deriving (A.3). Therefore the field multiplets of $d = 4$, $\mathcal{N} = 2$ decompose as multiplets of $d = 2$ $(2,2)$ supersymmetry without putting any conditions such are normally employed in dimensional reduction. In particular, with the above identification, the four-dimensional fields in a vectormultiplet can be assembled as

$$\Upsilon = \varphi - \mathrm{i}\sqrt{2}\vartheta^{+}\psi_{21} - \mathrm{i}\sqrt{2}\ \overline{\vartheta^{-}}\psi_{12} + \vartheta^{+}\overline{\vartheta^{-}}(F_{03} - \mathrm{i}(F_{12} - D_{12})) \tag{A.4}$$

to produce a twisted chiral multiplet with values in the adjoint representation of some gauged Lie algebra $\mathfrak{g}$. The remaining degrees of freedom in the 4d vectormultiplet decompose in an infinite tower of "semi-chiral (2,2) multiplets" in the language of [90, 91, 92].

With a surface defect we may restrict the four-dimensional field (A.4) to the surface to produce a twisted chiral multiplet on the worldsheet of the surface defect. The associated vectormultiplet can be used to gauge a flavor $\mathfrak{g}$-symmetry of a $(2,2)$ quantum field theory. In particular the adjoint scalar φ plays the role of a twisted mass parameter.

The full supersymmetric expression for the twisted superpotential contribution to the action is

$$\exp\left[\frac{\mathrm{i}}{2}\int \mathrm{d}x^3\mathrm{d}x^0\left(\frac{\partial\mathcal{W}_i}{\partial a^I}(\mathrm{i}(D_{12} - F_{12}) + F^I_{03}) - \frac{\partial^2\mathcal{W}_i}{\partial a^I\partial a^J}\psi^I_+\psi^J_-\right) + c.c.\right] \tag{A.5}$$

The variation of the holomorphic term is exactly zero under two supersymmetries ${Q_1}^1$ and ${Q_2}^2$, and under the other two varies into total derivatives

$$\sim \int \mathrm{d}x^3\mathrm{d}x^0(\partial_0 \pm \partial_3)\left(\frac{\partial\mathcal{W}_i}{\partial a^I}\psi^I_{\mp}\right). \tag{A.6}$$

Therefore, when placed on a half-space $x^0 \leq 0$ the supersymmetric variation can be cancelled by a line defect of the form (5.17).

Finally, let us comment on the BPS representations and the definition of the index $\mu(\gamma_{ij})$. In our conventions, in a massive representation the little superalgebra is diagonalized by the combinations

$$\begin{aligned}\{Q_+ + e^{\mathrm{i}\rho}Q_-, \bar{Q}_+ + e^{-\mathrm{i}\rho}\bar{Q}_-\} &= 4(E - |Z|),\\ \{Q_+ - e^{\mathrm{i}\rho}Q_-, \bar{Q}_+ - e^{-\mathrm{i}\rho}\bar{Q}_-\} &= 4(E + |Z|),\end{aligned} \tag{A.7}$$

with all other anticommutators vanishing for $e^{\mathrm{i}\rho}Z = |Z|$. Therefore, short multiplets when $E = |Z|$ are spanned by the doublet $|s\rangle$ and $(\bar{Q}_+ - e^{-i\rho}\bar{Q}_-)|s\rangle$, where the Clifford vacuum is defined by

$$(Q_+ + e^{\mathrm{i}\rho}Q_-)|s\rangle = (\bar{Q}_+ + e^{-\mathrm{i}\rho}\bar{Q}_-)|s\rangle = (Q_+ - e^{\mathrm{i}\rho}Q_-)|s\rangle = 0. \tag{A.8}$$

Note that to define the index of [20][51] we must take F to be the $u(1)_V$ symmetry. That is, it must satisfy $[F, Q_+] = Q_+$ and $[F, Q_-] = Q_-$. Note that this leaves F ambiguous by a shift by any operator commuting with the entire superalgebra. The contribution of the short representation to the index $\mathrm{Tr}\, F e^{\mathrm{i}\pi F}$ is

---

[51] A related "new supersymmetric index" was defined earlier in [18]. The relation of this index to the index μ is roughly analogous to the relation between the hyperkähler metric on $\mathcal{M}$ and the indices Ω in [1].

$$(f-(f-1))e^{\mathrm{i}\pi f} = e^{\mathrm{i}\pi f} \tag{A.9}$$

We may take this short multiplet, denoted h, to represent the center of mass degree of freedom in the soliton multiplet. Factoring this out in a BPS representation space so that

$$\mathcal{H}^{BPS}_{\gamma_{ij}} = h \otimes \tilde{\mathcal{H}}^{BPS}_{\gamma_{ij}} \tag{A.10}$$

we may write

$$\mu(\gamma_{ij}) := \mathrm{Tr}_{\tilde{\mathcal{H}}^{BPS}_{\gamma_{ij}}} e^{\mathrm{i}\pi F} \tag{A.11}$$

B. A simple Hitchin system

We consider a rank 2 Hitchin system on $\mathbb{CP}^1$ with a single irregular singularity at $z=\infty$. This singularity is invariantly characterized by the statement that $\mathrm{Tr}\,\varphi^2$ has a fifth-order pole there. Since $\mathrm{Tr}\,\varphi^2$ is regular everywhere else, this actually fixes it (up to shifts of z and overall rescaling):

$$\mathrm{Tr}\,\varphi^2 = 2z\,(\mathrm{d}z)^2. \tag{B.1}$$

The precise boundary condition is given by specifying the form of (A,φ) near the singularity; it is given in §3 of [3], but we will not need it. What we do need to know is that there is a unique solution obeying this boundary condition. In this appendix we briefly summarize some facts about this solution. See §9.4.1 of [3] for its explicit form.

We consider the corresponding flat connection $\nabla(\zeta)$ given by (7.15). For any fixed ζ, the asymptotics of the flat sections of $\nabla(\zeta)$ exhibit Stokes phenomena as $z\to\infty$, with three anti-Stokes rays r_a, $a=1,2,3$ cyclically. The r_a are located at z for which $\frac{1}{\zeta}z^{3/2}\in\mathbb{R}$ (this condition makes sense even though $z^{3/2}$ is defined only up to sign). There are three "small flat sections" $s_a(\zeta,z)$ defined up to scalar multiple by the condition that s_a decays (exponentially fast) as $z\to\infty$ along r_a, indeed $s_a \sim \exp\left[-\frac{\pi R}{\zeta}\frac{2}{3}z_a^{3/2}\right]$, where "$z_a^{3/2}$" means the choice of $z^{3/2}$ making the exponent negative along r_a. This requirement implies that s_a is not a scalar multiple of s_{a+1}, so we can normalize the sections by the requirement that $(s_a,s_{a+1})=1$ where $(,)$ denotes an $SU(2)$-invariant antisymmetric inner product. This normalization fixes the s_a up to an overall sign, and implies that $\sum_a s_a = 0$. The s_a are not quite single-valued as functions of $\zeta\in\mathbb{C}^\times$: rather, varying $\zeta\to e^{2\pi\mathrm{i}}\zeta$ permutes the three anti-Stokes rays, and gives $s_a(e^{2\pi\mathrm{i}}\zeta) = -s_{a+1}(\zeta)$.[52] On the other hand they are perfectly analytic in z for fixed ζ.

Now fix a point z and a single root $z^{3/2}$. From the s_a just discussed, we can assemble two single-valued but piecewise-continuous sections $\mathcal{Y}_\pm(\zeta)$ which have uniform asymptotics as $\zeta\to 0$:

$$\mathcal{Y}_\pm(\zeta) \sim \exp\left[\pm\frac{\pi R}{\zeta}\frac{2}{3}z^{3/2}\right] e_\pm, \tag{B.2}$$

---

[52] From what we have explained so far, one can see easily that $s_a(e^{2\pi\mathrm{i}}\zeta) = \pm s_{a+1}(\zeta)$. The sign is subtler to determine.

where $e_\pm$ are ζ-independent.

We have to take a little care since s_a are multivalued as functions of ζ. Fixing a branch choice is equivalent to fixing a labeling of the three anti-Stokes rays for every ζ. First, for $\zeta \in z^{3/2}\mathbb{R}_+$, one of the three anti-Stokes rays actually contains the point z; label this one r_1, and then label the other two r_2, r_3 going around counterclockwise. We then vary this picture continuously with ζ, until we reach $\zeta \in z^{3/2}\mathbb{R}_-$, where we put a branch cut in the ζ-plane. Having chosen this labeling we now have $s_a(\zeta)$ well defined and single-valued.

We now take $(\mathcal{Y}_+, \mathcal{Y}_-) = (s_3, s_1)$ in the half-plane $\mathrm{Im}\frac{1}{\zeta}z^{3/2} > 0$ and $(\mathcal{Y}_+, \mathcal{Y}_-) = (-s_2, s_1)$ in the half-plane $\mathrm{Im}\frac{1}{\zeta}z^{3/2} < 0$. This gives two piecewise-continuous sections $\mathcal{Y}_\pm$ which have the desired asymptotics as $\zeta \to 0$. What are their discontinuities? As ζ crosses the ray $\frac{1}{\zeta}z^{3/2} \in \mathbb{R}_+$, the relation $-s_2 = s_1 + s_3$ gives

$$\mathcal{Y}_- \to \mathcal{Y}_-, \qquad \mathcal{Y}_+ \to \mathcal{Y}_+ + \mathcal{Y}_-, \tag{B.3}$$

while as ζ crosses $\frac{1}{\zeta}z^{3/2} \in \mathbb{R}_-$, using $-s_3 = s_1 - (-s_2)$ *and* taking account of the discontinuities of the s_i themselves across this cut gives

$$\mathcal{Y}_+ \to \mathcal{Y}_+, \qquad \mathcal{Y}_- \to \mathcal{Y}_- - \mathcal{Y}_+. \tag{B.4}$$

According to the Riemann-Hilbert correspondence, $\mathcal{Y}_\pm(z,\zeta)$ (with z fixed and ζ varying) are discontinuous flat sections for some connection in a rank-2 bundle over $\mathbb{CP}^1$, with irregular singularities at $\zeta = 0$ and $\zeta = \infty$ (and regular everywhere else.) The equations (B.3), (B.4) give the Stokes factors for this connection, at either irregular singularity, as $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and $\begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}$.

C. Two-dimensional bound state radius in Landau-Ginzburg theories

In this appendix we give a quantitative lower bound for how the "bound state radius" of a soliton in a Landau-Ginzburg theory diverges when three critical points become collinear.

If we take, for simplicity, a trivial Kahler metric $K_{\alpha\beta} = \delta_{\alpha,\beta}$ then the soliton equation for an (ij)-soliton is

$$\frac{\mathrm{d}\phi^\alpha}{\mathrm{d}x} = \zeta_{ji}\frac{\partial \bar{W}}{\partial \bar{\phi}^\alpha} \tag{C.1}$$

where $W(\phi)$ is the superpotential, the boundary condition is determined by critical points ϕ_i, ϕ_j:

$$\begin{aligned} \lim_{x\to-\infty} \phi(x) &= \phi_i \\ \lim_{x\to+\infty} \phi(x) &= \phi_j \end{aligned} \tag{C.2}$$

and $\zeta_{ji} = \frac{W(j)-W(i)}{|W(j)-W(i)|}$ is determined by the corresponding critical values. Here and below $W(i) := W(\phi_i)$, etc.

We can derive immediately

$$\frac{\mathrm{d}W}{\mathrm{d}x} = \zeta_{ji}\left|\frac{\partial \bar{W}}{\partial \bar{\phi}^\alpha}\right|^2 \tag{C.3}$$

so the image of the soliton in the W-plane is a straight line of slope ζ_{ji}. Near each critical point — assumed to be Morse — a quadratic approximation to W applies and the solution approaches the critical point with a decaying exponential. Therefore, we expect a rapid transition region from $\phi \approx \phi_i$ to $\phi \approx \phi_j$. This can be estimated from (C.3) to be

$$\Delta x_{ji} \sim \frac{|W(j) - W(i)|}{\left|\frac{\partial \bar{W}}{\partial \bar{\phi}^\alpha}\right|_*^2} \tag{C.4}$$

where in the denominator we use some typical intermediate value from the transition region. The soliton does not have a well-defined center, but it should be located within a region of size (C.4).

Now consider a situation where there are at least three critical points ϕ_i, ϕ_j, ϕ_k and the parameters of W are changed so that the critical values $W(i), W(j), W(k)$ are becoming collinear. We assume moreover that there are ij and jk solitons. As the critical values become collinear (and we approach the wall of marginal stability from a "stable" region) some of the ik solitons can be viewed as bound states of ij and jk solitons. We would like to estimate how the "bound state radius" of these bound states diverges as the critical values become collinear.

Along the x-axis an ik soliton of the bound state type will first have a rapid transition from $\phi \approx \phi_i$ to $\phi \approx \phi_j$ and then will stay close to $\phi \approx \phi_j$ for a long interval L and then transition rapidly from $\phi \approx \phi_j$ to $\phi \approx \phi_k$. The width of the two transition regions can be estimated using (C.4) above, and if they are much smaller than L, we can speak of an approximate bound state radius.

We can put a lower bound on the bound state radius as follows. Only the real part of $\frac{W}{\zeta}$ flows — the imaginary part is fixed. In order to get a lower bound on the distance in space corresponding to a part of the soliton trajectory in the W plane which passes close to the turning point j we can use a quadratic approximation, and integrate the equation in the finite W-plane region of the flow where the quadratic approximation is valid.

As a simple example capturing the spirit of the problem, consider a one-dimensional situation, with critical point ϕ_j at the origin and critical value $W(j) = 0$:

$$W = \frac{1}{2} C \phi^2 + \cdots \tag{C.5}$$

Then we can measure how close the flow is to the critical point by the small value ϵ of the imaginary part of $\frac{W}{\zeta}$. Let us denote the real part of $\frac{W}{\zeta}$ as t. Notice that

$$\left|\frac{\partial \bar{W}}{\partial \bar{\phi}^\alpha}\right|^2 \approx |C|^2 |\phi|^2 \approx 2|C||W| \approx 2|C|\sqrt{t^2 + \epsilon^2} + \cdots \tag{C.6}$$

So the distance in real space L is bounded below by

$$\int_{t_0}^{t_1} \frac{\mathrm{d}t}{2|C|\sqrt{t^2 + \epsilon^2}} \tag{C.7}$$

where t_0, t_1 are some fixed values far from the transition regions of the ij and jk solitons. The integral (C.7) diverges logarithmically as $\epsilon \to 0$:

$$\int_{t_0}^{t_1} \frac{\mathrm{d}t}{2|C|\sqrt{t^2+\epsilon^2}} \sim \frac{1}{|C|}\log\frac{1}{\epsilon} \tag{C.8}$$

If we consider the triangle formed by Z_i, Z_j, Z_k and let ϵ be the height of the perpendicular from Z_j to the segment $[Z_i, Z_k]$, then $\epsilon = \mathrm{Im}(Z_{ij}\frac{\bar{Z}_{ik}}{|Z_{ik}|})$, so we can write the result more invariantly as

$$L \geq \frac{1}{\Lambda}\log\left(\frac{|Z_{ik}|}{\mathrm{Im}(Z_{ij}\bar{Z}_{ik})}\right) \tag{C.9}$$

where Λ is a nonuniversal number. The argument of the logarithm is essentially the same quantity that arises in Denef's bound state radius formula for 4dimensional bound state radii.

If the field space is of higher dimension, the above estimate can be given as it is as long as $|\partial W|^2$ is bounded from above by $|C||W|$ for some constant C near the critical point. This is trivially true as long as the critical point is non-degenerate.

Thus, one can give a halo-type description to the decay of $1+1$ dimensional solitons, but the bound state radius formula is not as precise as in the four-dimensional case. The reason is that the physics responsible for the bound state is rather different, and does not involve the exchange of massless particles.

D. Physical models for the affine linearity of $\omega(\gamma, \gamma_i)$

The change of the number of chiral multiplets on the surface defect discussed in §3.5 is a rather novel phenomenon. In this appendix we offer some physical models for understanding why it happens.

D.1 A supersymmetric quantum mechanics model

As a warmup, we consider the following model problem. Let us thicken the solenoid so that it carries uniform magnetic flux B in a disk of radius R around the origin. We search for the ground states of a supersymmetric quantum mechanical charged particle in the complex plane coupled to this magnetic field. We can choose a gauge

$$A = \begin{cases} \frac{1}{2}B(z\mathrm{d}\bar{z} - \bar{z}\mathrm{d}z) & |z| \leq R \\ \frac{1}{2}BR^2\frac{(z\mathrm{d}\bar{z}-\bar{z}\mathrm{d}z)}{|z|^2} & |z| \geq R \end{cases} \tag{D.1}$$

The wavefunctions lie in $\Omega^{0,*}(\mathbb{C})$ and the supersymmetry operators are identified with $\bar{\partial} + \bar{A} : \Omega^{0,0}(\mathbb{C}) \to \Omega^{0,1}(\mathbb{C})$ and $(\bar{\partial} + \bar{A})^\dagger : \Omega^{0,1}(\mathbb{C}) \to \Omega^{0,0}(\mathbb{C})$. For $B > 0$ we find that the normalizable zero-energy wavefunctions in $\Omega^{0,0}$ have the form

$$\psi = \begin{cases} f(z)e^{-\frac{1}{2}B|z|^2} & |z| \leq R \\ (\frac{z\bar{z}}{R^2})^{-BR^2/2}e^{-\frac{1}{2}BR^2}f(z) & |z| \geq R \end{cases} \tag{D.2}$$

where $f(z)$ is an entire function. The total wavefunction is then only normalizable if $f(z)$ is a polynomial of degree n with $n < BR^2 - 1$. There are no normalizable zero-energy states in $\Omega^{0,1}$. The situation is reversed for $B < 0$ with normalizable wavefunctions in $\Omega^{0,1}$ governed by a polynomial $g(\bar{z})$ of degree $n < |B|R^2 - 1$ and no normalizable wavefunctions of type $\Omega^{0,0}$. One important point brought out by this example is that the number of states depends linearly on the flux through the solenoid, not just the holonomy around the solenoid.

D.2 Probe particle in the presence of a solenoid and dyon

In this section we analyze the quantum mechanics of a halo particle near a dyon interface in a solenoid in order to determine the structure of the one-particle Hilbert space which generates the halo Fock space, as used in §4.7.2.

We model the surface defect with interface as in §4.7.2 as a solenoid with a dyon of charge $\gamma_{i,j'} \in \Gamma_{i,j'}$ inside. One can write the fixed point equations for the preserved supersymmetries. A solution is provided by the attractor-like equations

$$2\mathrm{Im}(\zeta^{-1}Z(\gamma, u(r)) = -\frac{\langle \gamma, \gamma_{i,j'} \rangle}{r} + 2\mathrm{Im}(\zeta^{-1}Z(\gamma, u)). \tag{D.3}$$

In the self-dual formalism the gauge potential is

$$\mathcal{A} = \frac{1}{2}\left(-\cos\theta \mathrm{d}\phi \otimes \gamma_{ij'} - \frac{\mathrm{d}t}{r} \otimes I(\gamma_{ij'})\right), \tag{D.4}$$

where a projection of $\gamma_{ij'}$ to Γ_g is understood and I is the complex structure on $\Gamma_g \otimes \mathbb{R}$ at $u \in \mathcal{B}$. As in the Denef halo solutions (for details in this situation see [3], Appendix C or [49]), a test particle of charge γ^h is bound at a radius given by the same formula in the absence of the surface defect. (After all, outside the solenoid, the electromagnetic fields are zero, and hence the probing dyon feels no extra force from the solenoid.)

The effective Minkowski-signature Lagrangian of a halo particle in the probe approximation, confined to the bound state radius, is of the form

$$\int \frac{1}{2}\mu r^2(\sin^2\theta\dot{\phi}^2 + \dot{\theta}^2) - \int (\kappa_1 \cos\theta + \kappa_2^{\mathrm{left}})\dot{\phi} \tag{D.5}$$

or

$$\int \frac{1}{2}\mu r^2(\sin^2\theta\dot{\phi}^2 + \dot{\theta}^2) - \int (\kappa_1 \cos\theta + \kappa_2^{\mathrm{right}})\dot{\phi}. \tag{D.6}$$

Here θ, ϕ are angular coordinates centered on the dyon. The “left” half of the surface defect is $\theta = 0$ and the “right” half is $\theta = \pi$. The first Lagrangian applies for $0 \le \theta < \pi$ and the second for $0 < \theta \le \pi$. The two Lagrangians define a smooth quantum measure for $\kappa_2^{\mathrm{left}} - \kappa_2^{\mathrm{right}} \in \mathbb{Z}$. Moreover, κ_2^{left} and $\kappa_2^{\mathrm{right}}$ can be shifted by integers.

The Aharonov-Bohm phase of the probe particle of charge γ^h around a small circle far to the left is $\exp[2\pi\mathrm{i}\langle\gamma^h, \gamma_i^0\rangle]$ while that far to the right is $\exp[2\pi\mathrm{i}\langle\gamma^h, \gamma_{j'}^0\rangle]$. Shifting κ_2 by appropriate integers we may take the solution

$$\begin{aligned} \kappa_1 &= \frac{1}{2}\langle \gamma^h, \gamma_{ij'} \rangle, \\ \kappa_2^{\text{left}} &= \frac{1}{2}\left(\langle \gamma^h, \gamma_i^0 + \gamma_{j'}^0 - \gamma^c \rangle \right), \\ \kappa_2^{\text{right}} &= \frac{1}{2}\left(\langle \gamma^h, \gamma_i^0 + \gamma_{j'}^0 + \gamma^c \rangle \right). \end{aligned} \tag{D.7}$$

Note that

$$\kappa_2^{\text{left}} - \kappa_2^{\text{right}} = -\langle \gamma^h, \gamma^c \rangle. \tag{D.8}$$

Now let us analyze the ground state wavefunctions of the quantum particle with Lagrangian

$$\int \frac{1}{2}\mu r^2 (\sin^2\theta \dot{\phi}^2 + \dot{\theta}^2) - \int (\kappa_1 \cos\theta + \kappa_2)\dot{\phi}. \tag{D.9}$$

Such wavefunctions are of the form

$$\Psi = e^{\mathrm{i}m\phi}(1+\cos\theta)^a (1-\cos\theta)^b, \tag{D.10}$$

where $2m \in \mathbb{Z}$, with $2m$ even or odd according to whether the particle is a boson or a fermion. The values of a, b are

$$a = \frac{\text{sign}(\kappa_1)}{2}(\kappa_1 - m'), \qquad b = \frac{\text{sign}(\kappa_1)}{2}(\kappa_1 + m'). \tag{D.11}$$

with $m' = m + \kappa_2$. If we require that the wavefunction be nonsingular everywhere on the sphere, so that $a \geq 0$ and $b \geq 0$, then m' must lie in the interval

$$|\kappa_1| \geq m' \geq -|\kappa_1| \tag{D.12}$$

The energy is $E = \frac{|\kappa_1|}{2\mu r^2}$.

Let N_{κ_1,κ_2} denote the number of solutions of the inequality (D.12). In the case where κ_1 and κ_2 are not half-integers and $\kappa_1 > 0$ its value is given by

N_{κ_1,κ_2}	$\{\kappa_1\} < 1 - \{\kappa_2\}$	$\{\kappa_1\} \geq 1 - \{\kappa_2\}$
$\{\kappa_1\} \geq \{\kappa_2\}$	$2[\kappa_1] + 1$	$2[\kappa_1] + 2$
$\{\kappa_1\} < \{\kappa_2\}$	$2[\kappa_1]$	$2[\kappa_1] + 1$

(D.13)

where $x = [x] + \{x\}$ is the decomposition of a real number into its greatest integer and fractional parts.

In a chamber where there are no 2d bound states of charge γ_{ii}^h or $\gamma_{j'j'}^h$ we can identify

$$\omega(\gamma^h, \gamma_{ij'}) := N_{\kappa_1,\kappa_2}\Omega(\gamma^h). \tag{D.14}$$

Note that shifting $\gamma_{ij'} \to \gamma_{ij'} + \Delta\gamma^c$ leads to a shift in N_{κ_1,κ_2} reproducing the affine-linearity of $\omega(\gamma^h, \gamma_{ij'})$.

As an aside, we note the curious point that if we had only demanded that the wavefunctions be L^2-normalizable with respect to the measure $\mathrm{d}(\cos\theta)\mathrm{d}\phi$, then for generic values of $\{\kappa_1\}$ and $\{\kappa_2\}$ the ground state would in fact be only two-fold degenerate with ground state energy smaller than $|\kappa_1|$ and with the two eigenstates having diverging wavefunctions at the north and south poles, respectively.

E. Twistor construction of hyperholomorphic connections

In this appendix we explain how to construct the hyperholomorphic connection A on the bundle V over $\mathcal{M}$, from the sections $\mathcal{Y}_{\gamma_i}$ of V over $\mathcal{M} \times \mathbb{C}^{\times}$. The construction is local along $\mathcal{M}$, but global along $\mathbb{C}^{\times}$. It seems to be a close cousin to the construction of self-dual Yang-Mills fields [93] and more generally of hyperholomorphic connections from holomorphic vector bundles over twistor space, although we will not use those constructions explicitly.

Our construction depends crucially on the analytic properties of $\mathcal{Y}_{\gamma_i}$ as a function of $\zeta \in \mathbb{C}^{\times}$ at a fixed point of $\mathcal{M}$:

- $\mathcal{Y}_{\gamma_i}$ is *piecewise* holomorphic in ζ, with discontinuities only at the BPS rays;
- The jumps of $\mathcal{Y}_{\gamma_i}$ across BPS rays are of the form $\mathcal{Y}_{\gamma_i} \to \sum_{k,\gamma_k \in \Gamma_k} c_{\gamma_i,\gamma_k} \mathcal{Y}_{\gamma_k}$, with c_{γ_i,γ_k} constants;
- The limit of $\mathcal{Y}_{\gamma_i}(\mathcal{Y}^{\mathrm{sf}}_{\gamma_i})^{-1}$ as $\zeta \to 0$ is finite and smooth;
- Finally, to normalize the sections we put a reality condition on $\mathcal{Y}_{\gamma_i}$ by demanding that in the Hermitian metric h on V, defined by the unitary framing of §5.2.2, we have the unitarity constraints described in equations (5.35), et. seq. in §5.6.

The first three properties are consequences of the integral equations (5.23), (5.25). Together with the unitarity constraints all four properties determine the sections uniquely.

The first step in constructing the hyperholomorphic connection on V is to pass from these properties of the $\mathcal{Y}_{\gamma_i}$ to a system of "Cauchy-Riemann" equations for sections of V. We first recall that the Cauchy-Riemann equations for ordinary functions on $\mathcal{M}$ can be written in the form [1]

$$\frac{\partial}{\partial u}\mathcal{Y} = \mathcal{A}_u(\zeta)\mathcal{Y}, \tag{E.1}$$

$$\frac{\partial}{\partial \bar{u}}\mathcal{Y} = \mathcal{A}_{\bar{u}}(\zeta)\mathcal{Y}, \tag{E.2}$$

where $\mathcal{A}_u(\zeta)$ and $\mathcal{A}_{\bar{u}}(\zeta)$ are two complex vertical vector fields on the torus fibers, depending holomorphically on $\zeta \in \mathbb{C}^{\times}$.[53] In particular, these equations hold for the functions $\mathcal{Y} = \mathcal{Y}_\gamma$. One can describe the ζ dependence of these vector fields more precisely: they are of the form

$$\mathcal{A}_u(\zeta) = \frac{1}{\zeta}\mathcal{A}_u^{(-1)} + \mathcal{A}_u^{(0)}, \tag{E.3}$$

$$\mathcal{A}_{\bar{u}}(\zeta) = \mathcal{A}_{\bar{u}}^{(0)} + \zeta\mathcal{A}_{\bar{u}}^{(1)}, \tag{E.4}$$

where $\mathcal{A}_u^{(-1)}$ are linearly independent at every point, and similarly $\mathcal{A}_{\bar{u}}^{(1)}$. This form of the Cauchy-Riemann equations is one of the main ingredients in the twistorial construction of the hyperkähler metric g.

---

[53] Of course $\mathcal{B}$ can have dimension larger than one, so we should understand u to stand for coordinates u^m, $m = 1, \ldots, \dim \mathcal{B}$, with the index suppressed.

Now choose a local trivialization of V, so that the $\mathcal{Y}_{\gamma_j}$ are represented as vector-valued functions. Also choose a single $\gamma_j^0 \in \Gamma_j$ for each j. The $\mathcal{Y}_{\gamma_j^0}$ are linearly independent vectors, at least for R large enough. We can thus define matrix-valued functions $\mathcal{B}_u(\zeta)$, $\mathcal{B}_{\bar{u}}(\zeta)$ on $\mathcal{M}$ by requiring

$$\frac{\partial}{\partial u}\mathcal{Y}_{\gamma_j^0} = \mathcal{A}_u(\zeta)\mathcal{Y}_{\gamma_j^0} + \mathcal{B}_u(\zeta)\mathcal{Y}_{\gamma_j^0}, \tag{E.5}$$

$$\frac{\partial}{\partial \bar{u}}\mathcal{Y}_{\gamma_j^0} = \mathcal{A}_{\bar{u}}(\zeta)\mathcal{Y}_{\gamma_j^0} + \mathcal{B}_{\bar{u}}(\zeta)\mathcal{Y}_{\gamma_j^0}, \tag{E.6}$$

for all j. Here each $\mathcal{A}$ term is a vector field on $\mathcal{M}$ which acts by differentiation, but is proportional to the identity matrix as an endomorphism of V, while each $\mathcal{B}$ term is a matrix-valued function on $\mathcal{M}$. It is straightforward to see that the $\mathcal{B}$ terms are continuous functions of ζ despite the fact that $\mathcal{Y}_{\gamma_j^0}$ are not, and moreover that the $\mathcal{B}$ terms are independent of our choice of γ_j^0. So (E.5), (E.6) can be thought of as Cauchy-Riemann equations defining a holomorphic structure $\bar{\partial}^{(\zeta)}$ on V.

We want to show that the ζ dependence of $\mathcal{B}$ has the same form as that of $\mathcal{A}$, i.e. that we can decompose

$$\mathcal{B}_u(\zeta) = \frac{1}{\zeta}\mathcal{B}_u^{(-1)} + \mathcal{B}_u^{(0)}, \tag{E.7}$$

$$\mathcal{B}_{\bar{u}}(\zeta) = \mathcal{B}_{\bar{u}}^{(0)} + \zeta\mathcal{B}_{\bar{u}}^{(1)}. \tag{E.8}$$

For this purpose we first consider the semiflat version of (E.5) and (E.6). In that setting we have [1]

$$\mathcal{A}_u^{(-1),\mathrm{sf}} = -\mathrm{i}\pi R\frac{\partial Z}{\partial u}\cdot\frac{\partial}{\partial\theta}, \quad \mathcal{A}_{\bar{u}}^{(1),\mathrm{sf}} = -\mathrm{i}\pi R\frac{\partial \bar{Z}}{\partial \bar{u}}\cdot\frac{\partial}{\partial\theta}, \tag{E.9}$$

$$\mathcal{A}_u^{(0),\mathrm{sf}} = 0, \quad \mathcal{A}_{\bar{u}}^{(0),\mathrm{sf}} = 0, \tag{E.10}$$

and a direct computation using (5.13) then shows that (E.5), (E.6) are satisfied if we take $\mathcal{Y} = \mathcal{Y}^{\mathrm{sf}}$ and $\mathcal{B}$ to have the form (E.7) where $\mathcal{B}_u^{(-1),\mathrm{sf}}$ is diagonal with jj^{th} matrix element given by:

$$\left(\mathcal{B}_u^{(-1),\mathrm{sf}}\right)_{jj} = \pi R\frac{\partial Z_{\gamma_j^0}}{\partial u}, \quad \left(\mathcal{B}_{\bar{u}}^{(1),\mathrm{sf}}\right)_{jj} = \pi R\frac{\partial \bar{Z}_{\gamma_j^0}}{\partial \bar{u}}, \tag{E.11}$$

$$\mathcal{B}_u^{(0),\mathrm{sf}} = 0, \quad \mathcal{B}_{\bar{u}}^{(0),\mathrm{sf}} = 0. \tag{E.12}$$

Here to write a definite formula we needed a trivialization of V, and we picked the one determined by the local sections γ_j^0 of Γ_j; a ζ-independent change of trivialization would affect the detailed form of $\mathcal{B}$ but not the general structure of its ζ dependence.

Away from the semiflat approximation, to see that $\mathcal{B}$ still has the desired form, we use our asymptotic condition on $\mathcal{Y}_{\gamma_j^0}$. An efficient way to get the result is as follows. For any fixed u, we can consider the torus $\mathcal{M}_u$ as the locus of real values of θ_γ, sitting inside its complexification $\mathcal{M}_u^{\mathbb{C}}$ where θ_γ are allowed to be arbitrary. The functions $\mathcal{Y}_\gamma^{\mathrm{sf}}$ have natural continuations to this complexification. Then we can define a map $\Upsilon(\zeta) : \mathcal{M}_u \to \mathcal{M}_u^{\mathbb{C}}$ by requiring $\mathcal{Y}_\gamma = \mathcal{Y}_\gamma^{\mathrm{sf}} \circ \Upsilon$. The asymptotics of $\mathcal{Y}_\gamma$ [1] say that $\Upsilon(\zeta)$ is finite in the limit $\zeta \to 0$ or $\zeta \to \infty$. Our asymptotic condition on $\mathcal{Y}_{\gamma_j^0}$ above just says that the natural extension of $\Upsilon(\zeta)$ to a map of *bundles* is also finite as $\zeta \to 0, \infty$. In particular we have $\mathcal{B}(\zeta) = \Upsilon(\zeta)^* \mathcal{B}^{\mathrm{sf}}(\zeta)$, so we get the desired form for $\mathcal{B}$, with $\mathcal{B}^{(-1)}$ and $\mathcal{B}^{(1)}$ given by the pullbacks of their semiflat values (E.11), (E.12).

Now we are ready to define the unitary connection A. We will engineer A so that (E.5) and (E.6) say

$$[(\mathrm{d} + A)\mathcal{Y}_{\gamma_j^0}]^{(0,1)_\zeta} = 0. \tag{E.13}$$

In other words, (E.5) and (E.6) are the ζ-dependent Cauchy-Riemann equations for sections of a hyperholomorphic bundle. The equivalence between (E.5), (E.6) and (E.13) means

$$A\left(\mathcal{A}_u^{(-1)}\right) = \mathcal{B}_u^{(-1)}, \tag{E.14}$$

$$A\left(-\partial_u + \mathcal{A}_u^{(0)}\right) = \mathcal{B}_u^{(0)}, \tag{E.15}$$

$$A\left(-\partial_{\bar u} + \mathcal{A}_{\bar u}^{(0)}\right) = \mathcal{B}_{\bar u}^{(0)}, \tag{E.16}$$

$$A\left(\mathcal{A}_{\bar u}^{(1)}\right) = \mathcal{B}_{\bar u}^{(1)}. \tag{E.17}$$

These requirements uniquely define A.

Note that

1. Applying this procedure in the semiflat case, i.e. using (E.9)-(E.12), we quickly recover A^{sf} given in (5.4).

2. It would have been impossible to construct an A with the desired property if $\mathcal{B}$ had contained, say, a $1/\zeta^2$ term.

3. The transformations of $\mathcal{Y}_{\gamma_i}$ across the BPS walls W_a involve multiplication by holomorphic quantities and do not lead to discontinuities in A.

4. Our reality conditions show that

   $$A^\dagger = -A \tag{E.18}$$

   i.e. $\mathrm{d} + A$ is a *unitary* connection with respect to the metric h in V.

5. Following the discussion in [1] we expect that the sections $\mathcal{Y}_{\gamma_i}$ also satisfy interesting equations in R and ζ expressing anomalous Ward identities. We leave a full discussion of this for another occasion.

F. $\mathcal{Y}_{\gamma_{ij'}}$ in A_1 theories

In this appendix we give a precise definition of the sections $\mathcal{Y}_{\gamma_{ij'}}$ in A_1 theories. The main point of this discussion is to check carefully the sign σ appearing in the multiplicative law, and to see that the discontinuities of $\mathcal{Y}_{\gamma_{ij'}}$ are as expected.

F.1 Defining $\mathcal{Y}$

We choose (u,ζ) generic (not on any BPS wall), and also choose the edges of the WKB triangulation so that z does not lie on any edge. Let z_i and z_j denote the two preimages of z on Σ.

Recall from [2] that given an edge E of a quadrilateral Q_E of the WKB triangulation, we defined γ_E to be the odd sum of lifts of a path in Q_E connecting the two turning points. Its overall orientation is fixed by requiring $\langle \gamma_E, \hat{E}\rangle = 1$, where $\hat{E}$ denotes either of the lifts of E to Σ, with its standard WKB orientation (we will use this orientation several times below.) Then with vertices 1234 going around Q_E counterclockwise, and using the shorthand ab for $s_a \wedge s_b$, we defined

$$\mathcal{Y}_{\gamma_E} = -\mathcal{X}_{\gamma_E} = \frac{12\ 34}{23\ 41}. \tag{F.1}$$

More general $\mathcal{Y}_\gamma$ were defined using the rule

$$\mathcal{Y}_{\gamma+\gamma'} = (-1)^{\langle\gamma,\gamma'\rangle}\mathcal{Y}_\gamma\mathcal{Y}_{\gamma'}. \tag{F.2}$$

Now let S denote the "sector" where z sits (a triangle bounded by an edge of the WKB triangulation and two separating WKB curves; two vertices of this triangle are vertices of the WKB triangulation, while the third is a turning point.) Let $\gamma_{ij,S}$ denote a path from z_i to z_j, obtained as the odd sum of lifts of a path from z_i to the turning point on the boundary of S. Also let

$$s_{i,S} = s_a(bc) \tag{F.3}$$

where a is the vertex reached by flowing along a lifted WKB curve from z_i, and vertices abc go around the triangle counterclockwise. Similarly define $s_{j,S}$.

Define $\mathcal{Y}_{\gamma_{ij,S}}$ to be an endomorphism of the fiber over z of our rank-2 bundle, which maps $s_{i,S} \mapsto 0$ and $s_{j,S} \mapsto \nu_{i,S}s_{i,S}$, where $\nu_{i,S} = +1$ if the lifted WKB curve through z_i goes around the triangle counterclockwise, and $\nu_{i,S} = -1$ if it goes around clockwise. For more general γ_{ij} we define $\mathcal{Y}_{\gamma_{ij}}$ using the rule

$$\mathcal{Y}_{\gamma_{ij}+\gamma} = \sigma(\gamma)\mathcal{Y}_\gamma\mathcal{Y}_{\gamma_{ij}}, \tag{F.4}$$

where σ is the "canonical quadratic refinement" of [2].

Let $\gamma_{ii,0}$ denote the element in Γ_{ii} corresponding to $0 \in \Gamma$. Define $\mathcal{Y}_{\gamma_{ii,0}}$ to be the endomorphism which maps $s_i \mapsto s_i$ and $s_j \mapsto 0$. For more general γ_{ii} we use the rule

$$\mathcal{Y}_{\gamma_{ii,0}+\gamma} = \mathcal{Y}_\gamma\mathcal{Y}_{\gamma_{ii,0}}. \tag{F.5}$$

The fact that σ is indeed a quadratic refinement implies that the product we have defined is associative.

F.2 Multiplication laws

Directly from the definitions we have the relation

$$\mathcal{Y}_{\gamma_{ij,S}}\mathcal{Y}_{\gamma_{ji,S}} = -\mathcal{Y}_{\gamma_{ii,0}}. \tag{F.6}$$

(The minus sign comes from the fact that $\nu_{i,S}\nu_{j,S} = -1$.)

Multiplying both sides by general $\mathcal{Y}_\gamma$ this becomes

$$\sigma(\gamma)\mathcal{Y}_{\gamma_{ij,S}+\gamma}\mathcal{Y}_{\gamma_{ji,S}} = -\mathcal{Y}_\gamma\mathcal{Y}_{\gamma_{ii,0}}, \tag{F.7}$$

i.e.

$$\mathcal{Y}_{\gamma_{ij,S}+\gamma}\mathcal{Y}_{\gamma_{ji,S}} = -\sigma(\gamma)\mathcal{Y}_{\gamma_{ii,0}+\gamma}. \tag{F.8}$$

Multiplying in another $\mathcal{Y}_{\gamma'}$ gives

$$\sigma(\gamma')\mathcal{Y}_{\gamma_{ij,S}+\gamma}\mathcal{Y}_{\gamma_{ji,S}+\gamma'} = -\sigma(\gamma)\mathcal{Y}_{\gamma'}\mathcal{Y}_{\gamma_{ii,0}+\gamma}, \tag{F.9}$$

i.e.

$$\mathcal{Y}_{\gamma_{ij,S}+\gamma}\mathcal{Y}_{\gamma_{ji,S}+\gamma'} = -\sigma(\gamma)\sigma(\gamma')(-1)^{\langle\gamma,\gamma'\rangle}\mathcal{Y}_{\gamma_{ii,0}+\gamma+\gamma'}, \tag{F.10}$$

which is

$$\mathcal{Y}_{\gamma_{ij}}\mathcal{Y}_{\gamma_{ji}} = -\sigma(\gamma_{ij}+\gamma_{ji})\mathcal{Y}_{\gamma_{ij}+\gamma_{ji}}. \tag{F.11}$$

Finally, multiplying (F.5) by $\mathcal{Y}_{\gamma'}$ gives

$$\mathcal{Y}_{\gamma'}\mathcal{Y}_{\gamma_{ii,0}+\gamma} = \mathcal{Y}_{\gamma'}\mathcal{Y}_\gamma\mathcal{Y}_{\gamma_{ii,0}}, \tag{F.12}$$

i.e.

$$\mathcal{Y}_{\gamma'}\mathcal{Y}_{\gamma_{ii,0}+\gamma} = (-1)^{\langle\gamma,\gamma'\rangle}\mathcal{Y}_{\gamma'+\gamma+\gamma_{ii,0}}, \tag{F.13}$$

i.e. (writing $\gamma_{ii} = \gamma_{ii,0}+\gamma$ and then relabeling $\gamma' \to \gamma$)

$$\mathcal{Y}_\gamma\mathcal{Y}_{\gamma_{ii}} = (-1)^{\langle\gamma_{ii},\gamma\rangle}\mathcal{Y}_{\gamma+\gamma_{ii}} \tag{F.14}$$

where in writing $\langle\gamma_{ii},\gamma\rangle$ we use the isomorphism $\Gamma_{ii} \simeq \Gamma$.

The equtaions (F.2), (F.4), (F.11), (F.14) give the definition of the σ appearing in the multiplicative law for $\mathcal{Y}$:

$$\sigma(\gamma,\gamma') = (-1)^{\langle\gamma,\gamma'\rangle}, \tag{F.15}$$

$$\sigma(\gamma,\gamma_{ii}) = (-1)^{\langle\gamma,\gamma_{ii}\rangle}, \tag{F.16}$$

$$\sigma(\gamma,\gamma_{ij}) = \sigma(\gamma), \tag{F.17}$$

$$\sigma(\gamma_{ij},\gamma_{ji}) = -\sigma(\gamma_{ij}+\gamma_{ji}). \tag{F.18}$$

F.3 Morphism: crossing a separating WKB curve

Now suppose we displace from z in sector S clockwise to z' in sector S', while remaining in the same triangle. In what follows we drop the subscript S, and for quantities which should have subscript S' we just put a prime on top. We are free to choose i to be the sheet with $\nu_i = 1$, i.e. the WKB curves passing through z_i go around counterclockwise. Then $\nu'_i = -1$.

Number the vertices of the triangle so that the WKB curve through z_i runs from vertex 1 to 2. We have

$$\mathcal{Y}_{\gamma_{ij}}(s_i) = 0, \quad \mathcal{Y}_{\gamma_{ij}}(s_j) = s_i \tag{F.19}$$

i.e.

$$\mathcal{Y}_{\gamma_{ij}}(s_2) = 0, \quad \mathcal{Y}_{\gamma_{ij}}(s_1) = s_2 \frac{31}{23}, \tag{F.20}$$

and also

$$\mathcal{Y}_{\gamma_{ji}}(s_j) = 0, \quad \mathcal{Y}_{\gamma_{ji}}(s_i) = -s_j \tag{F.21}$$

i.e.

$$\mathcal{Y}_{\gamma_{ji}}(s_1) = 0, \quad \mathcal{Y}_{\gamma_{ji}}(s_2) = -s_1 \frac{23}{31}. \tag{F.22}$$

Then the WKB curve through z_i' runs from 1 to 3, and we have

$$\mathcal{Y}'_{\gamma_{ij}}(s_i') = 0, \quad \mathcal{Y}'_{\gamma_{ij}}(s_j') = -s_i', \tag{F.23}$$

i.e.

$$\mathcal{Y}'_{\gamma_{ij}}(s_3) = 0, \quad \mathcal{Y}'_{\gamma_{ij}}(s_1) = -s_3 \frac{12}{23}. \tag{F.24}$$

Using the identity

$$(12)s_3 + (23)s_1 + (31)s_2 = 0 \tag{F.25}$$

this becomes

$$\mathcal{Y}'_{\gamma_{ij}}((23)s_1 + (31)s_2) = 0, \quad \mathcal{Y}'_{\gamma_{ij}}(s_1) = \left(\frac{23}{12}s_1 + \frac{31}{12}s_2\right)\frac{12}{23}, \tag{F.26}$$

i.e.

$$\mathcal{Y}'_{\gamma_{ij}}((23)s_1 + (31)s_2) = 0, \quad \mathcal{Y}'_{\gamma_{ij}}(s_1) = s_1 + \frac{31}{23}s_2, \tag{F.27}$$

i.e.

$$\mathcal{Y}'_{\gamma_{ij}}(s_2) = -\frac{23}{31}s_1 - s_2, \quad \mathcal{Y}'_{\gamma_{ij}}(s_1) = s_1 + \frac{31}{23}s_2, \tag{F.28}$$

which means

$$\mathcal{Y}'_{\gamma_{ij}} = \mathcal{Y}_{\gamma_{jj}} - \mathcal{Y}_{\gamma_{ii}} + \mathcal{Y}_{\gamma_{ij}} + \mathcal{Y}_{\gamma_{ji}} \tag{F.29}$$

or finally,

$$\mathcal{Y}'_{\gamma_{ij}} = (1 - \mathcal{Y}_{\gamma_{ji}})\mathcal{Y}_{\gamma_{ij}}(1 + \mathcal{Y}_{\gamma_{ji}}). \tag{F.30}$$

We also have

$$\mathcal{Y}'_{\gamma_{ji}}(s_j') = 0, \quad \mathcal{Y}'_{\gamma_{ji}}(s_i') = s_j', \tag{F.31}$$

i.e.

$$\mathcal{Y}'_{\gamma_{ji}}(s_1) = 0, \quad \mathcal{Y}'_{\gamma_{ji}}(s_3) = s_1 \frac{23}{12}. \tag{F.32}$$

Using (F.25) again this is

$$\mathcal{Y}'_{\gamma_{ji}}(s_1) = 0, \quad \mathcal{Y}'_{\gamma_{ji}}(s_2) = -s_1 \frac{23}{31}, \tag{F.33}$$

i.e.

$$\mathcal{Y}'_{\gamma_{ji}} = \mathcal{Y}_{\gamma_{ji}} \tag{F.34}$$

which is also

$$\mathcal{Y}'_{\gamma_{ji}} = (1 - \mathcal{Y}_{\gamma_{ji}})\mathcal{Y}_{\gamma_{ji}}(1 + \mathcal{Y}_{\gamma_{ji}}). \tag{F.35}$$

So in summary we have the expected transformation law

$$\mathcal{Y}' = \mathcal{S}^{\mu=1}_{\gamma_{ji}} \mathcal{Y}. \tag{F.36}$$

Note γ_{ji} is the path from z_j to z_i through the turning point — in particular, when ζ sits exactly on the BPS ray $\ell_{\gamma_{ji}}$ γ_{ji} can be represented by a lifted WKB path, with the *opposite* of the WKB orientation as expected (since $Z_{\gamma_{ji}}/\zeta \in \mathbb{R}_-$.)

F.4 Morphism: crossing an edge

We can also displace z across an edge E to z' in a neighboring triangle. We keep the labeling as above, so E is the 12 edge, and let the other vertex of the neighboring triangle be 4 (so 142 go around counterclockwise). Then going around the quadrilateral Q_E counterclockwise we have 1423, i.e.

$$\mathcal{Y}_{\gamma_E} = \frac{14\;23}{42\;31}. \tag{F.37}$$

We also have

$$s_i' = s_i \frac{14}{31}, \quad s_j' = s_j \frac{42}{23}, \quad \nu' = -\nu \tag{F.38}$$

which says

$$\mathcal{Y}'_{\gamma'_{ij}} = -\mathcal{Y}_{\gamma_E} \mathcal{Y}_{\gamma_{ij}} = \mathcal{Y}_{\gamma_E + \gamma_{ij}}, \qquad \mathcal{Y}'_{\gamma'_{ji}} = -\mathcal{Y}^{-1}_{\gamma_E} \mathcal{Y}_{\gamma_{ji}} = \mathcal{Y}_{-\gamma_E + \gamma_{ji}}. \tag{F.39}$$

But indeed $\gamma'_{ij} = \gamma_{ij} + \gamma_E$ and $\gamma'_{ji} = \gamma_{ji} - \gamma_E$ so this just says

$$\mathcal{Y}' = \mathcal{Y} \tag{F.40}$$

as it should since crossing an edge does not correspond to any BPS ray.

F.5 Morphism: flips and juggles

If the triangulation undergoes a flip or a juggle somewhere far away from z, then $\mathcal{Y}_{\gamma_{ij,S}}$ is unaffected. Together with the multiplication laws and the known action of flips and juggles on $\mathcal{Y}_\gamma$ from [2], we believe this is enough to show that the action of flips and juggles on $\mathcal{Y}$ is by $\mathcal{K}^\omega_\gamma$ as it should be.

G. A review: hidden flavor symmetries in 3d Coulomb branches

Consider some three-dimensional field theory which admits an effective low energy description as a non-linear sigma model on a smooth moduli space of vacua $\mathcal{M}$. (In this Appendix, $\mathcal{M}$ is a general target space, not necessarily the space (1.2).) Flavor symmetries of the UV theory which are spontaneously broken by the vacuum are realized as isometries of $\mathcal{M}$. Flavor symmetries of the UV theory which are unbroken at all points of $\mathcal{M}$ might still manifest themselves in the IR description, but in a more subtle way.

Let $\varphi : \mathbb{R}^{1,2} \to \mathcal{M}$ and consider a closed 2-form b on $\mathcal{M}$. The pullback of b is a closed 2-form in the 3d spacetime, hence at least classically a conserved current:

$$J = \varphi^* b. \tag{G.1}$$

Notice that two closed 2-forms which differ by an exact form give rise to currents which differ by an "improvement" term, and measure the same conserved charges. Thus, the sigma model has an abelian group of global symmetries whose Lie algebra may be identified with the de Rham cohomology $H^2_{DR}(\mathcal{M})$. The corresponding conserved charges cannot be changed by quantum effects in the IR sigma model, though they very well might not exist in the full UV theory.

Now note that the Hilbert space can be graded by $\pi_2(\mathcal{M})$ because a map $\varphi : \mathbb{R}^2 \to \mathcal{M}$ at a fixed time maps the boundary at infinity in $\mathbb{R}^2$ to the vacuum, i.e. a point in $\mathcal{M}$. Elementary excitations are in the trivial homotopy class and solitonic excitations are in nontrivial homotopy classes. Suppose $\exp : H^2_{DR}(\mathcal{M}) \to \mathcal{F}$ is the exponential map of the flavor group $\mathcal{F}$. If $f = \exp(b)$ is in the connected component of the identity then the action of f on wavefunctions in a sector $[\varphi] \in \pi_2(\mathcal{M})$ is

$$f \cdot \Psi = e^{2\pi \mathrm{i} \int \varphi^* b} \Psi \tag{G.2}$$

The charge thus defines a character in $\mathrm{Hom}(\pi_2(\mathcal{M}), U(1))$, which actually only depends on the image under the Hurewicz map, i.e. it descends to $\mathrm{Hom}(H_2(\mathcal{M}, \mathbb{Z}), U(1))$.

Now, $\mathrm{Hom}(H_2(\mathcal{M}, \mathbb{Z}), U(1)) \simeq H^2(\mathcal{M}, U(1))$ so it is natural to suspect that $\mathcal{F} \simeq H^2(\mathcal{M}, U(1))$ is the correct expression for the full flavor group. $H^2(\mathcal{M}, U(1))$ is a compact abelian group whose Lie algebra may be identified with $H^2_{DR}(\mathcal{M})$. The connected component of the identity of this group can be identified with a torus $\mathcal{T} = H^2_{DR}(\mathcal{M})/H_2(\mathcal{M}, \mathbb{Z})$, in harmony with (G.2). However, $\mathcal{F}$ might well have a nontrivial discrete group $\mathcal{D}$ of components, leading to discrete flavor symmetries not continuously connected to the identity and equation (G.2) generalizes naturally to this case. In general, $\mathcal{F}$ is a semidirect product of the connected torus $\mathcal{T}$ with a discrete group $\mathcal{D}$. (In fact $\mathcal{D}$ is isomorphic to the torsion subgroup of $H^3(\mathcal{M}, \mathbb{Z})$.) If $\mathcal{M}$ also has an isometry group G, it will act on the space of closed 2-forms and more generally on $\mathcal{F}$; hence the full flavor symmetry group of the IR theory will be a semidirect product of G and $\mathcal{F}$.

It is an interesting problem to relate the flavor symmetry group of the UV theory and of the IR sigma-model description. A sufficient amount of supersymmetry, say $\mathcal{N} = 2$ in three dimensions (4 supercharges), makes the task easier. To every $U(1)$ subgroup of the UV flavor group, one has a real mass parameter m in the UV theory, which would be the vev of the scalar superpartner of a background gauge field gauging the $U(1)$ flavor subgroup. In the IR, the parameter m for a flavor symmetry which is unbroken everywhere on the Coulomb branch only affects the metric of the sigma model, and hence enters in a very specific way in the IR Lagrangian. Supersymmetrization of this coupling gives a coupling to the background gauge field, i.e. to the IR conserved current for the $U(1)$ flavor symmetry. In superspace, the kinetic term arises from integrating the Kähler potential

$$\int \mathrm{d}^4\theta \, \mathcal{K}(\Phi, \bar{\Phi}, M). \tag{G.3}$$

Here M is a linear multiplet whose lowest component is the real mass m. (Thus M is a sum $\Psi + \bar{\Psi}$ of chiral multiplets, m is the real part of the complex scalar, and the imaginary part is dualized to a flavor gauge field with fieldstrength F_{fl}.) Expanding, we find a coupling

$$\int (\partial - \bar{\partial})\partial_m \mathcal{K} \wedge F_{fl}. \tag{G.4}$$

As the Kähler potential and its first derivatives are not globally defined, we should integrate by parts to an expression involving the Kähler form:

$$\int \partial_m \omega \wedge A_{fl}. \tag{G.5}$$

Hence the relevant 2-form b on $\mathcal{M}$ is simply the variation of the Kähler form under a change in the real mass parameter.

We can extend the above statements, valid for $U(1)$ subgroups of the flavor group, to discrete flavor symmetry groups which manifest themselves in the IR in non-trivial components of $\mathcal{F}$. If $f \in \mathcal{F}$ then we can modify the path integral on $\mathbb{R}^{1,1} \times S^1$ by using f-twisted boundary conditions around S^1 – that is, we can insert a "flavor Wilson line" corresponding to f. If the circle is sufficiently large that we can use the effective three-dimensional IR theory, but sufficiently small that we can express the partition function as that of an effective two-dimensional theory, then this twisting is equivalent to the insertion of a B-field amplitude in the two-dimensional sigma model. For example, if $f = \exp(b)$ is an element of the connected component of the identity in $\mathcal{F}$ then the twisted partition function has an insertion of an operator

$$\exp 2\pi \mathrm{i} \int_{2d} \varphi^* b \tag{G.6}$$

which survives the limit of taking a small circle. Hence the flavor Wilson line induces a B-field b in the effective 2d sigma model. This makes perfect sense for a $U(1)$ flavor symmetry: the 3d Kähler parameter m becomes a complexified 2d Kähler parameter built out of m and b. For a general $f \in \mathcal{F}$, we get a coupling for the corresponding flat B-field in $H^2(\mathcal{M}, U(1)) = \mathcal{F}$.

Finally, it is useful to consider line defects in the 3d theory which introduce a monodromy given by a flavor group element f in $\mathcal{F}$. Such defects would appear naturally if we were to gauge a (possibly discrete) subgroup of $\mathcal{F}$ containing f. Now, a line defect stretching along $\ell \subset \mathbb{R}^{1,2}$ can be viewed as the boundary of a domain wall S, i.e. $\partial S = \ell$. The fields across the domain wall are related by a flavor group transformation f. If $f = \exp b$ is in the connected component of the identity then the domain wall carries a coupling

$$\exp 2\pi \mathrm{i} \int_{S} \varphi^* b. \tag{G.7}$$

More generally, it has a coupling pairing the class $\varphi^*(f)$ with S. Because S has a boundary, this pairing does not define a complex number, but rather a section of a line bundle on the loop space of spacetime, as in the theory of D-branes in the presence of a B-field. In order to define a good operator in the sigma model we need the line operator at the boundary of S to be a section of the dual line bundle on loop space.

As in the theory of D-branes in the presence of a topologically nontrivial B-field there is a natural class of such line operators related to “connections on twisted bundles.” See [94] for a concrete discussion. The flat B-field $f \in H^2(\mathcal{M}, U(1))$ determines a (torsion) twisting class in $\mathrm{Tors}(H^3(\mathcal{M}, \mathbb{Z}))$. The twisted class can be viewed as an ’t Hooft flux, thus determining — for example — an $SU(N)/\mathbb{Z}_N$ bundle which does not lift to an $SU(N)$ bundle. A connection on the $SU(N)/\mathbb{Z}_N$ bundle has well-defined holonomy in the adjoint representation, but the holonomy in representations of $SU(N)$ transforming nontrivially under the center must be regarded as sections of line bundles over the loop space of $\mathcal{M}$. For an appropriately twisted vector bundle and connection, then, $\mathrm{Hol}(A, \ell)$ – the trace of the holonomy in the fundamental representation – will live in the dual line bundle (over loop space) to that where (G.7) is valued, and hence the product

$$\mathrm{Hol}(A, \ell) \cdot \exp 2\pi \mathrm{i} \int_S \varphi^* b \tag{G.8}$$

will be a well-defined function. In the present context this gives a well-defined line operator.

References

[1] D. Gaiotto, G. W. Moore, and A. Neitzke, “Four-dimensional wall-crossing via three-dimensional field theory,” `arXiv:0807.4723 [hep-th]`.

[2] D. Gaiotto, G. W. Moore, and A. Neitzke, “Wall-crossing, Hitchin Systems, and the WKB Approximation,” `arXiv:0907.3987 [hep-th]`.

[3] D. Gaiotto, G. W. Moore, and A. Neitzke, “Framed BPS States,” `arXiv:1006.0146 [hep-th]`.

[4] A. Kapustin, “Wilson-’t Hooft operators in four-dimensional gauge theories and S-duality,” *Phys. Rev.* **D74** (2006) 025005, `arXiv:hep-th/0501015`.

[5] A. Kapustin and E. Witten, “Electric-magnetic duality and the geometric Langlands program,” `arXiv:hep-th/0604151`.

[6] S. Gukov and E. Witten, “Gauge theory, ramification, and the geometric Langlands program,” `arXiv:hep-th/0612073`.

[7] E. Witten, “Gauge Theory And Wild Ramification,” `arXiv:0710.0631 [hep-th]`.

[8] S. Gukov and E. Witten, “Rigid Surface Operators,” `arXiv:0804.1561 [hep-th]`.

[9] D. S. Freed, “Extended structures in topological quantum field theory,” `arXiv:hep-th/9306045`.

[10] J. C. Baez and J. Dolan, “Higher dimensional algebra and topological quantum field theory,” *J. Math. Phys.* **36** (1995) 6073–6105, `arXiv:q-alg/9503002`.

[11] J. Lurie, “On the classification of topological field theories,” in *Current developments in mathematics, 2008*, pp. 129–280. Int. Press, Somerville, MA, 2009.

[12] D. S. Freed, M. J. Hopkins, J. Lurie, and C. Teleman, “Topological Quantum Field Theories from Compact Lie Groups,” `arXiv:0905.0731 [math.AT]`.

[13] A. Kapustin, “Topological Field Theory, Higher Categories, and Their Applications,” `arXiv:1004.2307 [math.QA]`.

[14] A. Kapustin and N. Saulina, "Surface operators in 3d Topological Field Theory and 2d Rational Conformal Field Theory," `arXiv:1012.0911 [hep-th]`.

[15] A. Kapustin and N. Saulina, "The algebra of Wilson-'t Hooft operators," *Nucl. Phys.* **B814** (2009) 327–365, `arXiv:0710.2097 [hep-th]`.

[16] A. Kapustin, "Holomorphic reduction of N = 2 gauge theories, Wilson-'t Hooft operators, and S-duality," `arXiv:hep-th/0612119`.

[17] D. Gaiotto, "Surface Operators in N=2 4d Gauge Theories," `arXiv:0911.1316 [hep-th]`.

[18] S. Cecotti, P. Fendley, K. A. Intriligator, and C. Vafa, "A New supersymmetric index," *Nucl. Phys.* **B386** (1992) 405–452, `arXiv:hep-th/9204102`.

[19] S. Cecotti and C. Vafa, "Ising model and N=2 supersymmetric theories," *Commun. Math. Phys.* **157** (1993) 139–178, `arXiv:hep-th/9209085`.

[20] S. Cecotti and C. Vafa, "On classification of $\mathcal{N} = 2$ supersymmetric theories," *Commun. Math. Phys.* **158** (1993) 569–644, `arXiv:hep-th/9211097`.

[21] N. Dorey, "The BPS spectra of two-dimensional supersymmetric gauge theories with twisted mass terms," *JHEP* **11** (1998) 005, `arXiv:hep-th/9806056`.

[22] S. Cecotti. Unpublished.

[23] D. Tong, "Quantum Vortex Strings: A Review," *Annals Phys.* **324** (2009) 30–52, `arXiv:0809.5060 [hep-th]`.

[24] A. Hanany and K. Hori, "Branes and N = 2 theories in two dimensions," *Nucl. Phys.* **B513** (1998) 119–174, `arXiv:hep-th/9707192`.

[25] N. Seiberg and E. Witten, "Monopole Condensation, And Confinement In N=2 Supersymmetric Yang-Mills Theory," *Nucl. Phys.* **B426** (1994) 19–52, `arXiv:hep-th/9407087`.

[26] N. Seiberg and E. Witten, "Monopoles, duality and chiral symmetry breaking in $\mathcal{N} = 2$ supersymmetric QCD," *Nucl. Phys.* **B431** (1994) 484–550, `arXiv:hep-th/9408099`.

[27] W. Lerche, "Introduction to Seiberg-Witten theory and its stringy origin," *Nucl. Phys. Proc. Suppl.* **55B** (1997) 83–117, `arXiv:hep-th/9611190`.

[28] L. Alvarez-Gaume and S. F. Hassan, "Introduction to S-duality in N = 2 supersymmetric gauge theories: A pedagogical review of the work of Seiberg and Witten," *Fortsch. Phys.* **45** (1997) 159–236, `arXiv:hep-th/9701069`.

[29] A. de Saint Exupéry, "Le Petit Prince".

[30] E. Witten, "Phases of $\mathcal{N} = 2$ theories in two dimensions," *Nucl. Phys.* **B403** (1993) 159–222, `arXiv:hep-th/9301042`.

[31] D. R. Morrison and M. Ronen Plesser, "Summing the instantons: Quantum cohomology and mirror symmetry in toric varieties," *Nucl. Phys.* **B440** (1995) 279–354, `arXiv:hep-th/9412236`.

[32] S. Cecotti and C. Vafa, "On classification of $\mathcal{N} = 2$ supersymmetric theories," *Commun. Math. Phys.* **158** (1993) 569–644, `arXiv:hep-th/9211097`.

[33] M. Kontsevich and Y. Soibelman, "Stability structures, motivic Donaldson-Thomas invariants and cluster transformations," `arXiv:0811.2435 [math.AG]`.

[34] F. Denef and G. W. Moore, "Split states, entropy enigmas, holes and halos," `arXiv:hep-th/0702146`.

[35] E. Andriyash, F. Denef, D. L. Jafferis, and G. W. Moore, "Wall-crossing from supersymmetric galaxies," `arXiv:1008.0030 [hep-th]`.

[36] N. Seiberg and E. Witten, "Gauge dynamics and compactification to three dimensions," `arXiv:hep-th/9607163`.

[37] T. Gocho and H. Nakajima, "Einstein-Hermitian connections on hyper-Kähler quotients," *J. Math. Soc. Japan* **44** (1992) no. 1, 43–51.

[38] N. S. Manton and B. J. Schroers, "Bundles over moduli spaces and the quantization of BPS monopoles," *Ann. Phys.* **225** (1993) 290–338.

[39] M. Verbitsky, "Hyperholomorphic bundles over a hyper-Kähler manifold," *J. Algebraic Geom.* **5** (1996) no. 4, 633–669.

[40] M. Verbitsky, "Hyperkähler manifolds with torsion obtained from hyperholomorphic bundles," *Math. Res. Lett.* **10** (2003) no. 4, 501–513.

[41] M. Verbitsky and D. Kaledin, *Hyperkahler manifolds*, vol. 12 of *Mathematical Physics (Somerville)*. International Press, Somerville, MA, 1999.

[42] B. Feix, "Hypercomplex manifolds and hyperholomorphic bundles," *Math. Proc. Cambridge Philos. Soc.* **133** (2002) no. 3, 443–457.

[43] E. Witten, "Solutions of four-dimensional field theories via M-theory," *Nucl. Phys.* **B500** (1997) 3–42, `arXiv:hep-th/9703166`.

[44] D. Gaiotto, "$\mathcal{N} = 2$ dualities," `arXiv:0904.2715 [hep-th]`.

[45] A. Klemm, W. Lerche, P. Mayr, C. Vafa, and N. P. Warner, "Self-dual strings and n=2 supersymmetric field theory," *Nucl. Phys.* **B477** (1996) 746–766, `arXiv:hep-th/9604034`.

[46] N. Seiberg, "Modifying the Sum Over Topological Sectors and Constraints on Supergravity," *JHEP* **07** (2010) 070, `arXiv:1005.0002 [hep-th]`.

[47] T. Banks and N. Seiberg, "Symmetries and Strings in Field Theory and Gravity," `arXiv:1011.5120 [hep-th]`.

[48] E. Andriyash, F. Denef, D. L. Jafferis, and G. W. Moore, "Bound state transformation walls," `arXiv:1008.3555 [hep-th]`.

[49] G. W. Moore, "PiTP Lectures on BPS States and Wall-Crossing in d=4, N=2 Theories". `http://www.physics.rutgers.edu/~gmoore/`. Talk 34.

[50] T. Dimofte, S. Gukov, and Y. Soibelman, "Quantum Wall Crossing in N=2 Gauge Theories," *Lett.Math.Phys.* **95** (2011) 1–25, `arXiv:0912.1346 [hep-th]`.

[51] E. Witten, "D-branes and K-theory," *JHEP* **12** (1998) 019, `arXiv:hep-th/9810188`.

[52] N. Nekrasov and E. Witten, "The Omega Deformation, Branes, Integrability, and Liouville Theory," `arXiv:1002.0888 [hep-th]`.

[53] T. Hausel and M. Thaddeus, "Mirror symmetry, Langlands duality, and the Hitchin system," `arXiv:math.AG/0205236`.

[54] G. Thompson, "Holomorphic vector bundles, knots and the Rozansky-Witten invariants," *Adv. Theor. Math. Phys.* **5** (2002) 457–481, `arXiv:hep-th/0002168`.

[55] H. Ooguri and C. Vafa, “Summing up D-instantons,” *Phys. Rev. Lett.* **77** (1996) 3296–3298, `arXiv:hep-th/9608079`.

[56] N. Seiberg and S. H. Shenker, “Hypermultiplet moduli space and string compactification to three dimensions,” *Phys. Lett.* **B388** (1996) 521–523, `arXiv:hep-th/9608086`.

[57] E. Frenkel and E. Witten, “Geometric Endoscopy and Mirror Symmetry,” `arXiv:0710.5939 [math.AG]`.

[58] S. Cecotti and C. Vafa, “Topological-antitopological fusion,” *Nucl. Phys.* **B367** (1991) 359–461.

[59] L. F. Alday, D. Gaiotto, S. Gukov, Y. Tachikawa, and H. Verlinde, “Loop and surface operators in N=2 gauge theory and Liouville modular geometry,” *JHEP* **01** (2010) 113, `arXiv:0909.0945 [hep-th]`.

[60] D. Gaiotto and J. Maldacena, “The gravity duals of N=2 superconformal field theories,” `arXiv:0904.4466 [hep-th]`.

[61] N. J. Hitchin, “The self-duality equations on a Riemann surface,” *Proc. London Math. Soc. (3)* **55** (1987) no. 1, 59–126.

[62] V. Balaji, I. Biswas, O. Gabber, and D. S. Nagaraj, “Brauer obstruction for a universal vector bundle,” *C. R. Math. Acad. Sci. Paris* **345** (2007) no.˜5, 265–268.

[63] I. Biswas and A. Hogadi, “Brauer group of moduli spaces of $PGL(r)$-bundles over a curve,” `arXiv:0904.4640 [math.AG]`.

[64] I. Biswas and A. Dey, “Brauer group of a moduli space of parabolic vector bundles over a curve,” `arXiv:1005.3161 [math.AG]`.

[65] V. Fock and A. Goncharov, “Moduli spaces of local systems and higher Teichmüller theory,” *Publ. Math. Inst. Hautes Études Sci.* (2006) no. 103, 1–211, `arXiv:math/0311149`.

[66] V. Fock, “Dual Teichmuller spaces,” `arXiv:dg-ga/9702018`.

[67] V. V. Fock and A. B. Goncharov, “Dual Teichmüller and lamination spaces,” in *Handbook of Teichmüller theory. Vol. I*, vol. 11 of *IRMA Lect. Math. Theor. Phys.*, pp. 647–684. Eur. Math. Soc., Zürich, 2007. `arXiv:math/0510312`.

[68] J. Teschner, “An analog of a modular functor from quantized Teichmuller theory,” `arXiv:math/0510174`.

[69] A. D. Shapere and C. Vafa, “BPS structure of Argyres-Douglas superconformal theories,” `arXiv:hep-th/9910182`.

[70] A. Bilal and F. Ferrari, “The Strong-Coupling Spectrum of the Seiberg-Witten Theory,” *Nucl. Phys.* **B469** (1996) 387–402, `arXiv:hep-th/9602082`.

[71] A. Bilal and F. Ferrari, “Curves of Marginal Stability and Weak and Strong-Coupling BPS Spectra in $N = 2$ Supersymmetric QCD,” *Nucl. Phys.* **B480** (1996) 589–622, `arXiv:hep-th/9605101`.

[72] A. Kapustin and E. Witten, “Electric-magnetic duality and the geometric Langlands program,” `arXiv:hep-th/0604151`.

[73] M. Herbst, “On higher rank coisotropic A-branes,” `arXiv:1003.3771 [hep-th]`.

[74] E. Witten, “Fivebranes and Knots,” `arXiv:1101.3216 [hep-th]`.

[75] K. Hori, A. Iqbal, and C. Vafa, "D-branes and mirror symmetry," `arXiv:hep-th/0005247 [hep-th]`.

[76] E. J. Martinec and G. W. Moore, "On decay of K theory," `arXiv:hep-th/0212059 [hep-th]`.

[77] G. W. Moore and A. Parnachev, "Localized tachyons and the quantum McKay correspondence," *JHEP* **0411** (2004) 086, `arXiv:hep-th/0403016 [hep-th]`.

[78] G. W. Moore and A. Parnachev, "Profiling the brane drain in a nonsupersymmetric orbifold," *JHEP* **0601** (2006) 024, `arXiv:hep-th/0507190 [hep-th]`.

[79] L. D. Faddeev and L. A. Takhtajan, *Hamiltonian methods in the theory of solitons*. Classics in Mathematics. Springer, Berlin, english ed., 2007. Translated from the 1986 Russian original by Alexey G. Reyman.

[80] O. Babelon, D. Bernard, and M. Talon, *Introduction to classical integrable systems*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 2003.

[81] S. L. Lukyanov and A. B. Zamolodchikov, "Quantum Sine(h)-Gordon Model and Classical Integrable Equations," `arXiv:1003.5333 [math-ph]`.

[82] L. F. Alday and J. Maldacena, "Null polygonal Wilson loops and minimal surfaces in Anti-de-Sitter space," *JHEP* **0911** (2009) 082, `arXiv:0904.0663 [hep-th]`.

[83] L. F. Alday, D. Gaiotto, and J. Maldacena, "Thermodynamic Bubble Ansatz," `arXiv:0911.4708 [hep-th]`.

[84] L. F. Alday, J. Maldacena, A. Sever, and P. Vieira, "Y-system for Scattering Amplitudes," *J.Phys.A* **A43** (2010) 485401, `arXiv:1002.2459 [hep-th]`.

[85] L. F. Alday, D. Gaiotto, J. Maldacena, A. Sever, and P. Vieira, "An Operator Product Expansion for Polygonal null Wilson Loops," `arXiv:1006.2788 [hep-th]`.

[86] H. Ooguri and C. Vafa, "Knot invariants and topological strings," *Nucl. Phys.* **B577** (2000) 419–438, `arXiv:hep-th/9912123`.

[87] A. Neitzke and J. Walcher, "Background independence and the open topological string wavefunction," in *From Hodge theory to integrability and TQFT*. American Mathematical Society, Providence, R.I., 2008. `arXiv:0709.2390 [hep-th]`.

[88] M. Aganagic and M. Yamazaki, "Open BPS Wall Crossing and M-theory," *Nucl. Phys.* **B834** (2010) 258–272, `arXiv:0911.5342 [hep-th]`.

[89] P. Sulkowski, "Wall-crossing, open BPS counting and matrix models," `arXiv:1011.5269 [hep-th]`.

[90] T. Buscher, U. Lindstrom, and M. Rocek, "New supersymmetric sigma models with Wess-Zumino terms," *Phys. Lett.* **B202** (1988) 94.

[91] U. Lindstrom, M. Rocek, R. von Unge, and M. Zabzine, "Generalized Kaehler manifolds and off-shell supersymmetry," *Commun. Math. Phys.* **269** (2007) 833–849, `arXiv:hep-th/0512164`.

[92] U. Lindstrom, M. Rocek, I. Ryb, R. von Unge, and M. Zabzine, "T-duality and Generalized Kahler Geometry," *JHEP* **02** (2008) 056, `arXiv:0707.1696 [hep-th]`.

[93] R. S. Ward, “On selfdual gauge fields,” *Phys. Lett.* **A61** (1977) 81–82.

[94] A. Kapustin, “D-branes in a topologically nontrivial B-field,” *Adv. Theor. Math. Phys.* **4** (2000) 127–154, `arXiv:hep-th/9909089`.